RETRACTED ARTICLE: Advantages of clinical pathways in severity-based treatment of spondylodiscitis

Lars Homagk^{1,2} • T. Jarmuzek³ • N. Homagk¹ • G. O. Hofmann⁴

Received: 28 February 2019 / Revised: 20 July 2019 / Accepted: 20 August 2019 / Published online: 24 August 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

The Editor-in-Chief has retracted this article [1] because, owing to a technical error, it was accidentally published after being rejected. Springer Nature apologizes to the authors and to readers. All authors agree to this retraction.

[1] Homagk, L., Jarmuzek, T., Homagk, N., and Hofmann, G.O.; Neurosurg Rev (2019). <https://doi.org/10.1007/s10143-019-01166-5>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10143-019-01166-5>) contains supplementary material, which is available to authorized users.

✉ Lars Homagk
info@praxisklinik-dr-homagk.de

¹ Praxisklinik Dr. Homagk, Naumburg, Germany

² Weißenfels, Germany

³ Centre for Spinal Cord Injuries, BG -Kliniken Bergmannstrost, Halle/Saale, Germany

⁴ Department of Trauma, Hand and Reconstructive Surgery, Friedrich Schiller University, Jena, Germany