

Nancy A. Schellhorn · David A. Andow

Response of coccinellids to their aphid prey at different spatial scales

Published online: 5 October 2005

© The Society of Population Ecology and Springer-Verlag Tokyo 2005

Popul Ecol (2005) 47:71–76

In Table 2, unnecessary words appeared in two entries in column 1. “Grid plot*plant” should read “Plot*plant,” and “Rid error” should read “Error.” The correct table appears here.

Table 2 Homogeneity-of-slopes model for the effect of aphids at the plant and plot scale on *C. maculata* and *H. convergens* adult density

Source	df	MS	<i>F</i>	<i>P</i>
<i>C. maculata</i>				
Plot	38	0.5599	1.46	0.0002
Plant	1	0.6565	1.71	0.1919
Plot*plant	38	0.5878	1.53	0.0241
Error	597	0.3845		
<i>H. convergens</i>				
Plot	38	0.1199	0.81	0.7832
Plant	1	1.5524	10.51	0.0013
Plot*plant	38	0.2430	1.65	0.0098
Error	598	0.1477		

Aphid density was log-transformed

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10144-004-0204-x>

N. A. Schellhorn · D. A. Andow
Department of Entomology, University of Minnesota,
Saint Paul, MN 55108, USA

N. A. Schellhorn (✉)
CSIRO Entomology, 120 Meiers Road,
Indooroopilly, QLD 4068, Australia
E-mail: nancy.schellhorn@csiro.au
Tel.: +61-7-32142721
Fax: +61-7-32142885