



Correction to: An analysis of tumor-related skin temperature differences in malignant soft-tissue tumors

Akiyoshi Shimatani¹ · Manabu Hoshi¹ · Naoto Oebisu¹ · Naoki Takada¹ · Yoshitaka Ban¹ · Hiroaki Nakamura¹

Published online: 17 October 2021
© Japan Society of Clinical Oncology 2021

Correction to: International Journal of Clinical Oncology
<https://doi.org/10.1007/s10147-021-02044-1>

In the original publication Table 4 has been published with incorrect entries. The incorrect and correct entries are given below:

Incorrect entry: G grade (median IQR) 4 [34]

Correct entry: G grade (median IQR) 4 [3-4]

Incorrect entry: Histological grade (median[IQR]) 2 [23]

Correct entry: Histological grade (median[IQR]) 2 [2-3]

The corrected version of Table 4 is given in this correction.

The original publication is also corrected.

Table 4 Relationship between clinical data and skin temperature differences ≥ 0.2 °C

	Skin temperature differences (°C)		p value
	0.2 ≤	<0.2	
Sex (n)			0.85
Men	28	37	
Women	24	29	
Age (years old)	61.8 ± 19.4	60.9 ± 13.8	0.77
Size (cm)	6.8 ± 5.3	5.2 ± 4.3	0.08
Distance (mm)	5.8 ± 6.7	9.4 ± 9.9	0.03*
CRP (mg/dL)	0.6 ± 1.9	0.5 ± 1.9	0.7
G grade (median [IQR])	4 [3-4]	2 [1-2]	<0.001*
Histological grade (median [IQR])	2 [2-3]	2 [1-2]	0.02*

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10147-021-02044-1>.

✉ Akiyoshi Shimatani
bpcip920@tcct.zaq.ne.jp

¹ Department of Orthopedic Surgery, Osaka City University Graduate School of Medicine, 1-4-3 Asahi-Machi, Abeno-ku, Osaka 545-8585, Japan