Erratum to: Is the lateralization distance important in terms in patients undergoing the modified Limberg flap procedure for treatment of pilonidal sinus?

**B. R. Karakaş · A. Aslaner · U. R. Gündüz ·
H. Çalış · A. N. Öngen · O. Z. Öner ·
N. Bülbüller**

Published online: 24 January 2015
© Springer-Verlag Italia Srl 2015

**Erratum to: Tech Coloproctol
DOI 10.1007/s10151-014-1252-7**

In the original publication of this article, the word “male” has been erroneously included. It should be removed from the “Results” section as female patients were also enrolled in the study (shown in Table 1).

The correct sentence in the beginning of paragraph 1 should now read as “In total, 40 consecutive patients were enrolled in the study and no patient was lost to follow-up.”

The online version of the original article can be found under doi:[10.1007/s10151-014-1252-7](https://doi.org/10.1007/s10151-014-1252-7).

B. R. Karakaş · A. Aslaner · U. R. Gündüz · H. Çalış ·
A. N. Öngen · O. Z. Öner · N. Bülbüller
Department of General Surgery, Antalya Training and Research
Hospital, Antalya 07100, Turkey

B. R. Karakaş (✉)
Department of General Surgery, Antalya Training and Research
Hospital, Antalya 07050, Turkey
e-mail: barisrafetkarakas@yahoo.com