



# Correction to: Index cost comparison of laparoscopic vs robotic surgery in colon and rectal cancer resection: a retrospective financial investigation of surgical methodology innovation at a single institution

E. U. Ezeokoli<sup>1</sup> · R. Hilli<sup>2</sup> · H. J. Wasvary<sup>2</sup>

Published online: 3 November 2022  
© Springer Nature Switzerland AG 2022

**Correction to: Techniques in Coloproctology**  
<https://doi.org/10.1007/s10151-022-02703-z>

The “Results” section was incorrect in the abstract text and it should read as given below.

## Results

Two hundred forty R-CR (mean age  $64.9 \pm 12.4$  years) and 258 L-CR (mean age  $66.4 \pm 15.5$  years) patients met the inclusion criteria. The overall mean direct cost between R-CR and L-CR was significantly higher (\$8756 vs \$7776

respectively,  $p = 0.001$ ) as well as the supply cost per case (\$3789 vs \$2122,  $p < 0.001$ ). Operating time was also higher for R-CR than L-CR (224 min vs 187 min,  $p = 0.066$ ) but LOS was slightly lower (5.08 days vs 5.55 days,  $p = 0.113$ ).

The original article has been corrected.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The original article can be found online at <https://doi.org/10.1007/s10151-022-02703-z>.

---

✉ E. U. Ezeokoli  
ekenex@gmail.com

<sup>1</sup> Oakland University William Beaumont School of Medicine, 586 Pioneer Dr., Rochester, MI 48309, USA

<sup>2</sup> Department of Colorectal Surgery, Beaumont Health Systems, Royal Oak, MI, USA