

## Erratum to: Serum level of soluble (pro)renin receptor is modulated in chronic kidney disease

Kazu Hamada · Yoshinori Taniguchi · Yoshiko Shimamura · Kosuke Inoue · Koji Ogata · Masayuki Ishihara · Taro Horino · Shimpei Fujimoto · Takashi Ohguro · Yukio Yoshimoto · Mika Ikebe · Kenji Yuasa · Eri Hoshino · Tatsuo Iiyama · Atsuhiko Ichihara · Yoshio Terada

Published online: 26 September 2013  
© Japanese Society of Nephrology 2013

**Erratum to: Clin Exp Nephrol**  
DOI 10.1007/s10157-013-0803-y

The original version of this article unfortunately contained errors.

In Table 1, in the first column, for the line “(P)RR”, the unit should be “ng/ml”.

In Figs. 1, 2, 3, 4, 5, 7, and 8, on the vertical axes, the unit for “soluble (P)RR” should be “ng/ml”.

In Fig. 6, on the vertical axis, the unit for “prorenin” should be “ng/ml”.

---

The online version of the original article can be found under doi:[10.1007/s10157-013-0803-y](https://doi.org/10.1007/s10157-013-0803-y).

---

K. Hamada · Y. Taniguchi · Y. Shimamura · K. Inoue · K. Ogata · M. Ishihara · T. Horino · S. Fujimoto · Y. Terada (✉)

Department of Endocrinology, Metabolism and Nephrology, Kochi University School of Medicine, Kohasu, Oko-cho, Nankoku, Kochi 783-8505, Japan  
e-mail: [terada@kochi-u.ac.jp](mailto:terada@kochi-u.ac.jp)

T. Ohguro · Y. Yoshimoto  
Department of Internal Medicine, Kochi Red Cross Hospital, Kochi, Japan

M. Ikebe · K. Yuasa  
Department of Urology and Internal Medicine, Kochi-Takasu Hospital, Kochi, Japan

E. Hoshino · T. Iiyama  
Clinical Trial Center, Kochi University School of Medicine, Kochi, Japan

A. Ichihara  
Department of Endocrinology and Hypertension, Tokyo Women's Medical University, Tokyo, Japan