



Correction to: Impaired endogenous nighttime melatonin secretion relates to intrarenal renin–angiotensin system activation and renal damage in patients with chronic kidney disease

Sayaka Ishigaki¹ · Naro Ohashi¹ · Shinsuke Isobe¹ · Naoko Tsuji¹ · Takamasa Iwakura¹ · Masafumi Ono¹ · Yukitoshi Sakao² · Takayuki Tsuji¹ · Akihiko Kato² · Hiroaki Miyajima¹ · Hideo Yasuda¹

Published online: 13 December 2018
© Japanese Society of Nephrology 2018

Correction to: Clin Exp Nephrol (2016) 20:878–884
<https://doi.org/10.1007/s10157-015-1224-x>

In the original publication, an error occurred in Table 4(B), under Nighttime group. The value of “Nighttime Log U-AGT/Cr” for model 3 (under $R=0.68$) was incorrectly published as 0.11. The correct value should read as -0.31 .

The corrected table is given below.

Table 4 Multiple regression analyses for urinary 6-sulphatoxymelatonin per hour (U-aMT6s/hr) levels by age, gender, estimated glomerular filtration rate (eGFR), and one of the parameters {blood pressures (BPs), urinary angiotensinogen/creatinine ratio (U-AGT/Cr), or urinary albumin/creatinine ratio (U-Alb/Cr)} during daytime and nighttime

The original article can be found online at <https://doi.org/10.1007/s10157-015-1224-x>.

✉ Naro Ohashi
ohashi-n@hama-med.ac.jp

¹ Internal Medicine 1, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu 431-3192, Japan

² Blood Purification Unit, Hamamatsu University School of Medicine, 1-20-1 Handayama, Higashi-ku, Hamamatsu 431-3192, Japan

B. Nighttime

	Model 1		Model 2		Model 3		Model 4	
	$R=0.66$	$P<0.001$	$R=0.65$	$P<0.001$	$R=0.68$	$P<0.001$	$R=0.69$	$P<0.001$
	β	P value	β	P value	β	P value	β	P value
Age	- 0.069	0.63	- 0.11	0.42	- 0.075	0.58	- 0.083	0.53
Gender	- 0.009	0.94	- 0.041	0.72	- 0.11	0.32	- 0.025	0.82
eGFR	0.50	0.002	0.55	0.001	0.38	0.024	0.52	< 0.001
Nighttime systolic BP	- 0.16	0.27						
Nighttime diastolic BP			- 0.046	0.72				
Nighttime log U-AGT/Cr					- 0.31	0.044		
Nighttime log U-Alb/Cr							- 0.25	0.025