Correction to: A nationwide survey on clinical practice patterns and bleeding complications of percutaneous native kidney biopsy in Japan

Takehiko Kawaguchi¹ · Tasuku Nagasawsa² · Kazuhiko Tsuruya³ · Kenichiro Miura⁴ · Takayuki Katsuno⁵ · Takashi Morikawa⁶ · Eiji Ishikawa⁷ · Masao Ogura⁸ · Hideki Matsumura⁹ · Ryota Kurayama¹⁰ · Shinsuke Matsumoto¹¹ · Yuhji Marui¹² · Shigeo Hara¹³ · Shoichi Maruyama¹⁴ · Ichiei Narita¹⁵ · Hirokazu Okada¹⁶ · Yoshifumi Ubara¹⁷ on behalf of Committee of Practical Guide for Kidney Biopsy 2019

Published online: 7 April 2020
 © Japanese Society of Nephrology 2020

Correction to: Clinical and Experimental Nephrology <https://doi.org/10.1007/s10157-020-01869-w>

In the original publication, some errors have been found in the alignment of Table 9. The corrected table is given below

Table 9 Biopsy procedures and timing of bleeding after biopsy for patients undergoing transcatheter arterial embolization (TAE)

Biopsy procedures	(%)	
Needle gauge	14	8
	15	62
	16	3

Biopsy procedures	(%)	
	17	27
Needle passes for sampling specimens	1	3
	2	32
	3	35
	4	14
	≥ 5	16
Use of hemostatic agents	Yes	57
	No	43
Manual compression on the biopsy site after biopsy	< 15 min	61
	15–30 min	31
	≥ 30 min	8

The original article can be found online at <https://doi.org/10.1007/s10157-020-01869-w>.

✉ Takehiko Kawaguchi
 kawatake45@gmail.com

- ¹ Department of Nephrology, National Hospital Organization Chibahigashi National Hospital, 673 Nitona-cho, Chuo-ku, Chiba 260-8712, Japan
- ² Division of Nephrology, Endocrinology, and Vascular Medicine, Tohoku University Hospital, Sendai, Japan
- ³ Department of Nephrology, Nara Medical University, Nara, Japan
- ⁴ Department of Pediatric Nephrology, Tokyo Women's Medical University, Tokyo, Japan
- ⁵ Department of Nephrology and Rheumatology, Aichi Medical University, Aichi, Japan
- ⁶ Department of Nephrology and Hypertension, Osaka City General Hospital, Osaka, Japan
- ⁷ Department of Nephrology, Mie University, Mie, Japan
- ⁸ Department of Nephrology and Rheumatology, National Center for Child Health and Development, Tokyo, Japan

- ⁹ Department of Pediatrics, Osaka Medical College, Osaka, Japan
- ¹⁰ Department of Pediatrics, Kosei General Hospital, Tokyo, Japan
- ¹¹ Department of Pediatrics, Matsudo City General Hospital, Chiba, Japan
- ¹² Department of Urology, St. Marianna University School of Medicine, Kawasaki, Japan
- ¹³ Department of Pathology, Kobe City Medical Center General Hospital, Kobe, Japan
- ¹⁴ Department of Nephrology, Nagoya University, Nagoya, Japan
- ¹⁵ Division of Clinical Nephrology and Rheumatology, Niigata University, Niigata, Japan
- ¹⁶ Department of Nephrology, Saitama Medical University, Saitama, Japan
- ¹⁷ Department of Nephrology, Toranomon Hospital, Tokyo, Japan

Biopsy procedures		(%)
Strict bed rest in a supine position after biopsy	< 2 h	26
	2–4 h	28
	4–8 h	30
	8–12 h	13
	≥ 12 h	3
Total bed rest after biopsy	< 8 h	13
	8–16 h	30
	16–24 h	48
	≥ 24 h	8
Number of kidney biopsies the operators had experienced before the biopsy for the patient with TAE	< 50	32
	50–200	39
	≥ 200	29
Timing of bleeding requiring TAE after biopsy		(%)
Cumulative timing of bleeding	< 6 h	53
	< 12 h	60
	< 24 h	65
	< 7 days	80
	≥ 7 days	20

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.