

© The Ichthyological Society of Japan 2011

### 2011 ISJ Meeting, Hirosaki, 29 September–2 October 2011

The 44th annual meeting of the Ichthyological Society of Japan (ISJ) will be held at the Bunkyo-cho Campus of Hirosaki University ([http://www.hirosaki-u.ac.jp/index\\_e.html](http://www.hirosaki-u.ac.jp/index_e.html)) in Hirosaki, Aomori, Japan, from 29 September to 2 October 2011. Following meetings of the board of editors and trustees on 29 September 2011, scientific sessions including oral and poster presentations will be held on 30 September and 1 October. One symposium entitled “Behavioral ecology of fishes: recent studies and future perspective” will be held on 2 October. Contact Yuji Sawara (Hirosaki University, e-mail: [isj-nenkai2011@bunken.co.jp](mailto:isj-nenkai2011@bunken.co.jp)) for further information about the annual meeting. Detailed information will be provided on the ISJ homepage (<http://www.fish-isj.jp/index-e.html>).

### Papers Published in *Japanese Journal of Ichthyology*

The following papers were published in *Japanese Journal of Ichthyology* Volume 58, Number 1 on 25 April 2011, and English abstracts are available on the ISJ website (<http://www.fish-isj.jp/english/journal.html>).

#### Volume 58, Number 1

##### Full Papers

Annual reproductive cycle of a wild population of oriental weather loach *Misgurnus anguillicaudatus* in Gifu Prefecture, central Japan. *Atsuko Kimura and Yasunori Koya.*

Longitudinal distribution of fishes in the Inohzawa River watershed, southern Izu Peninsula, Japan. *Noriyuki Takai, Akifumi Kato, Masaomi Uekusa, Yuko Kimura, Keitaro Dairiki, Shiro Itoi, Haruo Sugita, and Kiyoshi Yoshihara.*

First records of a lionfish, *Pterois mombasae* (Scorpaenidae: Pteroinae), from Japan, and morphological comparisons with *P. antennata*. *Mizuki Matsunuma and Hiroyuki Motomura.*

Reproductive ecology and life cycle of the invasive alien bitterling *Acheilognathus macropterus* (Cyprinidae, Acheilognathinae) in Lake Kasumigaura, Japan. *Tomiji Hagiwara.*

Seasonal changes in fish assemblages in an area of hermatypic corals in Yokonami, Tosa Bay, Japan. *Tomonori Hirata, Sosuke Oguri, Shiori Hirata, Hironobu Fukami, Yohei Nakamura, and Kosaku Yamaoka.*

Introduction of the dark chub *Zacco temminckii* from the Yoshino River system into the Yamato River system via the Yoshino diversion (Yoshino Bunsui). *Fumiko Ishii, Yukiko Anzai, Reika Ito, Naoto Koyama, and Tadao Kitagawa.*

##### Short Reports

Ultrastructure of the surface morphology of eggs in the terrestrial spawning blenny, *Andamia tetradactyla* (Blenniidae). *Norio Shimizu, Masako Hara, Yoichi Sakai, Hiroaki Hashimoto, and Kenji Gushima.*

Genetic composition of commercial strains of *Oryzias latipes* revealed by mtDNA analyses. *Naoto Koyama, Tomohiro Mori, Koji Nakai, and Tadao Kitagawa.*

First records of a silverside (Atheriniformes: Atherinidae), *Hypoatherina temminckii*, from Japan. *Daichi Sasaki and Seishi Kimura.*

First record of a deep-sea ophidiid fish, *Bassozetus glutinosus*, from Japan. *Shinichi Tomiyama, Munehiro Takami, and Atsushi Fukui.*