

© The Ichthyological Society of Japan 2016

ISJ meeting, Gifu, 23–26 September 2016

The 49th annual meeting of the Ichthyological Society of Japan (ISJ) will be held at Gifu University (<http://www.gifu-u.ac.jp/en/access/>) in Gifu City, Japan from 23 to 26 September 2016. The board of editors and trustees will be held on 23 September, and scientific sessions including oral and poster presentations on 24 and 25 September. One symposium entitled “Theory and practice of sustainable lotic ecosystem conservation in ecology of bitterling fishes (Cyprinidae, Acheilognathinae) and freshwater mussels (Unionidae)” will be held on 26 September. Contact Prof. Yasunori Koya (Faculty of Education, Gifu University, e-mail: isj-nenkai2016@bunken.co.jp) for further information on the annual meeting. Information on registration and mixer fees, and other detailed information will be provided on the ISJ official website (<http://www.fish-isj.jp/index-e.html>).

Papers published in *Japanese Journal of Ichthyology*

The following papers were published in Japanese Journal of Ichthyology Volume 63, Number 1 on 25 April 2016, and English abstracts are available on the ISJ website (<http://www.fish-isj.jp/english/journal.html>).

Volume 63, Number 1

Short reports

First record of *Conocara krefftii* (Alepocephalidae) collected from off Okinawa Island, Japan. *Munehiro Takami, Hiromitsu Endo and Atsushi Fukui.*

Upstream migration of adult amago salmon, *Oncorhynchus masou ishikawae* in the Nagara River, central Japan. *Tetsuya Tokuhara, Tomonori Kuwada, Ryouji Fujii, Toru Hara, Tetsuji Kariya and Daisuke Kishi.*

Distribution and origin of bagrid catfish *Tachysurus nudiceps* established in the Oita River system, Kyushu Island, Japan. *Hiroki Takano, Kazuo Hoshino, Tetsuya Okura, Toshio Matsuo and Katsutoshi Watanabe.*

Effect of dark chub *Nipponocypris temminckii* on feeding behavior of pale chub *Zacco platypus* in experimental ponds influenced by water currents. *Hiromu Ogawa and Osamu Katano.*

High levels of genetic diversity and gene flow in the endangered goby *Hypseleotris cyprinoides* on Okinawajima Island, Ishigakijima Island and Luzon Island. *Shumpei Tomita, Shohei Matsuzaki, Shin-ichiro Oka, Minoru Toda and Hideyuki Imai.*

A first record of a jawfish (Perciformes: Opistognathidae) *Opistognathus variabilis* from Yonaguni Island, Yaeyama Islands, Japan. *Yusuke Hibino, Daichi Sasaki and Seishi Kimura.*