Correction to: A new atherinomorine genus *Doboatherina* (Atheriniformes: Atherinidae) with a review of included species

Daichi Sasaki^{1,2} · Seishi Kimura^{1,3}

Published online: 9 January 2020
© The Ichthyological Society of Japan 2020

Correction to: Ichthyological Research
<https://doi.org/10.1007/s10228-019-00718-5>

The original publication of this article contains erroneous description and figure, which are corrected as below.

In “Comparison” of *Doboatherina aetholepis* (Kimura, Iwatsuki and Yoshino 2002), the second sentence reads,

“Although *D. aetholepis* shares a long spatular outgrowth posteriorly on the predorsal scales (upper jaw length 32–41% SL vs. 29–34%)”. However, the proportion of upper jaw length should be a ratio to head length (HL), not standard length (SL).

Figure 8 is incorrect. The correct figure is as follows.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10228-019-00718-5>.

✉ Seishi Kimura
kimura-s@bio.mie-u.ac.jp

¹ Fisheries Research Laboratory, Mie University, 4190-172 Wagu, Shima-cho, Shima, Mie 517-0703, Japan

² Present Address: Mie Prefecture Fisheries Research Institute, 3564-3 Hamajima, Hamajima-cho, Shima, Mie 517-0404, Japan

³ Present Address: Graduate School of Bioresources, Mie University, 1577 Kurimamachiya-cho, Tsu, Mie 514-8507, Japan

Fig. 8 Midlateral band and lateral scale laws at the level of second dorsal- and anal-fin origins in *Doboatherina aetholepis* (a), *D. balabacensis* (b), *D. bleekeri* (c), *D. duodecimalis* (d), *D. iwatsukii* sp. nov. (e), *D. magnidentata* sp. nov. (f), *D. salangensis* sp. nov. (g), *D. valenciennesi* (h), *D. woodwardi* (i), *D. yoshinoi* sp. nov. (j). a FRLM 14644, paratype, 69 mm SL, Ambon, Indonesia; b USNM 136802, neotype, 64 mm SL, Ticao I., Philippines; c FRLM 36718, 66 mm SL, Owase, Mie, Japan; d FRLM 36073, 73 mm SL, Ambon,

Indonesia; e MUFS 14670, paratype, 44 mm SL, Phuket, Thailand; f KAUM-I. 24170, paratype, 64 mm SL, Bang Saen, Chonburi, Thailand; g FRLM 47561, paratype, 48 mm SL, Phuket, Thailand; h FRLM 35363, 70 mm SL, Songkhla, Thailand; i URM-P 17407, 55 mm SL, Uruma, Okinawa, Japan; j FRLM 39132, paratype, 60 mm SL, Iriomote I., Okinawa, Japan. OD2, second dorsal-fin origin; OA, anal-fin origin; P, central pit of midlateral scales