Maximal estimates for a generalized spherical mean Radon transform acting on radial functions

Óscar Ciaurri¹ · Adam Nowak² · Luz Roncal^{3,4}

Received: 3 May 2019 / Accepted: 15 November 2019 / Published online: 26 November 2019
© Fondazione Annali di Matematica Pura ed Applicata and Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

We study a generalized spherical means operator, viz., generalized spherical mean Radon transform, acting on radial functions. As the main results, we find conditions for the associated maximal operator and its local variant to be bounded on power-weighted Lebesgue spaces. This translates, in particular, into almost everywhere convergence to radial initial data results for solutions to certain Cauchy problems for classical Euler–Poisson–Darboux and wave equations. Moreover, our results shed some new light on the interesting and important question of optimality of the yet known L^p boundedness results for the maximal operator in the general non-radial case. It appears that these could still be notably improved, as indicated by our conjecture of the ultimate sharp result.

Keywords Spherical Radon transform · Spherical mean · Maximal operator · Radial function · Weighted estimate · Wave equation · Euler–Poisson–Darboux equation · Axially symmetric solution · Convergence to initial data

The first-named author was supported by the Grant PGC2018-096504-B-C32 from Spanish Government. The second-named author was supported by the National Science Centre of Poland within the research project OPUS 2017/27/B/ST1/01623. The third-named author was supported by the Basque Government through BERC 2018–2021 program, by Spanish Ministry of Economy and Competitiveness MINECO through BCAM Severo Ochoa excellence accreditation SEV-2017-2018 and through the project MTM2017-82160-C2-1-P funded by (AEI/FEDER, UE) and acronym “HAQMEC,” and by 2017 Leonardo grant for Researchers and Cultural Creators, BBVA Foundation. The Foundation accepts no responsibility for the opinions, statements and contents included in the project and/or the results thereof, which are entirely the responsibility of the authors.

✉ Luz Roncal
lroncal@bcamath.org

Óscar Ciaurri
oscar.ciaurri@unirioja.es

Adam Nowak
adam.nowak@impan.pl

¹ Departamento de Matemáticas y Computación, Universidad de La Rioja, 26004 Logroño, Spain

² Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, 00-656 Warsaw, Poland

³ BCAM - Basque Center for Applied Mathematics, 48009 Bilbao, Spain

⁴ Ikerbasque, Basque Foundation for Science, 48011 Bilbao, Spain

Mathematics Subject Classification Primary 44A12; Secondary 42B37 · 35L15 · 35B07 · 35L05 · 35Q05

1 Preliminaries and statement of results

This paper is a natural continuation of our recent research from [2]. We study a generalized spherical means operator acting on radial functions. In [2], we viewed this operator as a family of integral transforms $\{M_t^{\alpha,\beta} : t > 0\}$ acting on profile functions on $\mathbb{R}_+$ and found fairly precise estimates of the associated integral kernels $K_t^{\alpha,\beta}(x, z)$. This enabled us to prove two-weight $L^p - L^q(L_t^r)$ estimates for $f \mapsto M_t^{\alpha,\beta} f$, with $1 \leq p, q \leq \infty$ and $1 \leq r < \infty$. In the present work, we focus on the more subtle limiting case $r = \infty$ and restrict to $p = q$, which in the above language of mixed norm estimates corresponds to weighted L^p -boundedness of the maximal operator $f \mapsto \sup_{t>0} |M_t^{\alpha,\beta} f|$. Obtaining such results requires a different, in fact more tricky, approach from that used in [2]. It is worth emphasizing that both our works, [2] and this one, were to large extent motivated by connections of the generalized spherical means with solutions to a number of classical initial-value PDE problems being of physical and practical importance; see, e.g., [2, Section 7] and references given there.

Let $n \geq 2$ and consider the generalized spherical means transformation

$$M^\beta f(x, t) = \mathcal{F}^{-1}(m_\beta(t|\cdot|)\mathcal{F}f)(x),$$

where $\mathcal{F}$ is the Fourier transform in $\mathbb{R}^n$ and the radial multiplier is given via

$$m_\beta(s) = 2^{\beta+n/2-1} \Gamma(\beta + n/2) \frac{J_{\beta+n/2-1}(s)}{s^{\beta+n/2-1}}, \quad s > 0,$$

with J_ν denoting the Bessel function of the first kind and order ν . The parameter β can, in general, be a complex number excluding $\beta = -n/2, -n/2 - 1, -n/2 - 2, \dots$. For $\beta = 0$, one recovers the classical spherical means

$$M^0 f(x, t) = \int_{S^{n-1}} f(x - ty) d\sigma(y), \quad (x, t) \in \mathbb{R}^n \times \mathbb{R}_+,$$

where $d\sigma$ is the normalized uniform measure on the unit sphere $S^{n-1} \subset \mathbb{R}^n$. Clearly, $M^0 f(x, t)$ returns the mean value of f on the sphere centered at x and of radius t .

For the maximal operator $M_*^\beta f = \sup_{t>0} |M^\beta f(\cdot, t)|$, Stein [8] proved the following.

Theorem 1.1 [8] *Let $n \geq 3$. Then, M_*^β is bounded on $L^p(\mathbb{R}^n)$ provided that*

$$1 < p \leq 2 \text{ and } \beta > 1 - n + \frac{n}{p}, \text{ or } p > 2 \text{ and } \beta > \frac{2-n}{p}. \quad (1.1)$$

This result was enhanced in the sense of admitted parameters and dimensions by subsequent authors: Bourgain [1], Mockenhaupt et al. [7] and recently by Miao et al. [6], see the historical comments in [6, p. 4272]. All these refinements can be stated altogether as follows, cf. [6, Theorem 1.1].

Theorem 1.2 [6] *Let $n \geq 2$. Then, M_*^β is bounded on $L^p(\mathbb{R}^n)$ provided that*

$$2 < p \leq \frac{2n+2}{n-1} \text{ and } \beta > \frac{1-n}{4} + \frac{3-n}{2p}, \text{ or } p > \frac{2n+2}{n-1} \text{ and } \beta > \frac{1-n}{p}. \quad (1.2)$$

The range of β in Theorem 1.2, for $p > 2$, is strictly wider than in Theorem 1.1; see Fig. 1 as follows. However, according to our best knowledge, it is not known whether it is already optimal. We strongly believe it is not, see our Conjecture 1.9 as follows. We remark that both Theorems 1.1 and 1.2 were originally proved for complex β , but for our purposes it is enough to state them for real values of the parameter.

A restriction of M^β to radially symmetric functions is still of interest and, moreover, admits a more explicit finer analysis that potentially leads to more general or stronger theorems. In this connection, we invoke two quite recent results of Duoandikoetxea et al. [4,5]. The first of them is a characterization of weighted L_{rad}^p -boundedness of M_*^0 with radial power weights involved; here and elsewhere the subscript “rad” indicates the subspace of radial functions.

Theorem 1.3 [4] *Let $n \geq 2$. The operator M_*^0 is bounded on $L_{\text{rad}}^p(\mathbb{R}^n, |x|^\gamma dx)$ for $1 < p < \infty$ (for $2 < p < \infty$ in case $n = 2$) if and only if*

$$1 - n \leq \gamma < p(n - 1) - n.$$

Another result in this spirit provides sufficient conditions for weighted L_{rad}^p -boundedness of $M_*^{(3-n)/2}$, again with radial power weights involved.

Theorem 1.4 [5] *Let $n \geq 2$ and $1 < p < \infty$. Then, the operator $M_*^{(3-n)/2}$ is bounded on $L_{\text{rad}}^p(\mathbb{R}^n, |x|^\gamma dx)$ provided that*

$$\frac{n-3}{2}p + 1 - n < \gamma < \frac{n+1}{2}p - n,$$

where in case $n = 2$ the lower bound for γ should be replaced by $-3/2$, and with the first inequality weakened for odd dimensions n .

Here and elsewhere by weakening a strict inequality, we mean replacing “ $<$ ” by “ $\leq$.” We take this opportunity to note that the statement of [5, Theorem 1.1] (an unweighted specification of Theorem 1.4) contains a small error, the upper bound for p in even dimensions higher than 3 should be given by strict inequality, cf. [5, Lemma 4.3]. This problem affects the abstract of [5] as well. The radial improvement for the unweighted estimates occurs only in odd dimensions higher than 3.

We omit here discussion of more sophisticated mapping properties of M_*^β , like boundedness from L^1 to weak L^1 , or from the Lorentz space $L^{1,1}$ to weak L^1 (which correspond to weak and restricted weak-type estimates for the maximal operator, respectively). For this kind of results, especially in the radial case, we refer to [4,5] and references given there.

The maximal operator $M_*^{\alpha,\beta}$ we will study generalizes the restriction of M_*^β to radial functions, since, in a sense, it covers a continuous range of dimensions $n = 2\alpha + 2$, $\alpha > -1$. Our main result, Theorem 1.5 as follows, contains Theorems 1.3 and 1.4 as special cases; in particular, we deliver a partly alternative and seemingly simpler proof of Theorem 1.4. Moreover, our more general perspective sheds new light on the discrepancy between $n = 2$ and higher dimensions in Theorems 1.3 and 1.4, as well as on the discrepancy between odd and even dimensions in Theorem 1.4. Finally, we gain some intuition that enables us to conjecture an optimal result in the spirit of Theorems 1.1 and 1.2, see Conjecture 1.9 as follows.

Let $\alpha > -1$ and $\alpha + \beta > -1/2$. For each $t > 0$, consider the integral operator

$$M_t^{\alpha,\beta} f(x) = \int_0^\infty K_t^{\alpha,\beta}(x, z) f(z) d\mu_\alpha(z), \quad x \in \mathbb{R}_+,$$

with the measure $d\mu_\alpha(x) = x^{2\alpha+1} dx$ and the kernel given by

$$K_t^{\alpha,\beta}(x, z) = \frac{2^{\alpha+\beta} \Gamma(\alpha + \beta + 1)}{t^{\alpha+\beta} (xz)^\alpha} \int_0^\infty J_{\alpha+\beta}(ty) J_\alpha(xy) J_\alpha(zy) y^{1-\alpha-\beta} dy.$$

This kernel is well defined for $(t, x, z) \in \mathbb{R}_+^3$ such that, in general, $t \neq |x - z|$ and $t \neq x + z$. Note that the integral here converges absolutely when $\alpha + \beta > 1/2$, but otherwise the convergence at ∞ is only conditional, in the Riemann sense.

For a radial function $f = f_0(|\cdot|)$ in $L^2(\mathbb{R}^n)$, $n \geq 2$, $M^\beta f(x, t)$ is for each $t > 0$ a radial function in $x \in \mathbb{R}^n$ whose profile is given by $M_t^{n/2-1, \beta} f_0$; see [2, Corollary 4.2]. Clearly, the maximal operators M_*^β and

$$M_*^{\alpha,\beta} f(x) = \sup_{t>0} |M_t^{\alpha,\beta} f(x)|$$

are connected in the same way. Thus, M_*^β is bounded on $L_{\text{rad}}^p(\mathbb{R}^n, |x|^\gamma dx)$ if and only if $M_*^{n/2-1, \beta}$ is bounded on $L^p(\mathbb{R}_+, x^\gamma d\mu_{n/2-1})$.

We now formulate the main result of this paper.

Theorem 1.5 *Assume that $\alpha > -1$ and $\alpha + \beta > -1/2$. Let $1 < p < \infty$ and $\delta \in \mathbb{R}$. Then, the maximal operator $M_*^{\alpha,\beta}$ is bounded on $L^p(\mathbb{R}_+, x^\delta dx)$ if*

$$\frac{1}{p} < \alpha + \beta + \frac{1}{2} \quad (1.3)$$

and

(a) in case $\beta \geq 1$,

$$-1 < \delta < (2\alpha + 2)p - 1, \quad (1.4)$$

(b) in case $0 < \beta < 1$,

$$\frac{-\beta}{[\beta \vee (\alpha + \beta + 1/2)] \wedge 1} < \delta < (2\alpha + \beta + 1)p - 1, \quad (1.5)$$

with the first inequality weakened when $\alpha + \beta > 1/2$,

(c) in case $\beta \leq 0$,

$$-\beta p < \delta < (2\alpha + \beta + 1)p - 1, \quad (1.6)$$

with the first inequality weakened when $-\beta \in \mathbb{N}$ or $\alpha + \beta > 1/2$.

We presume that Theorem 1.5 is sharp, but at the moment we are not able to give a complete justification of this statement. Nevertheless, we can show that Theorem 1.5 is optimal, or very close to optimal, for most choices of the parameters α, β .

Proposition 1.6 *Assume that $\alpha > -1$ and $\alpha + \beta > -1/2$. Let $1 < p < \infty$ and $\delta \in \mathbb{R}$. If $M_*^{\alpha,\beta}$ is bounded on $L^p(\mathbb{R}_+, x^\delta dx)$, then (1.3) holds and, moreover,*

$$-1 < \delta, \quad -\beta p \leq \delta, \quad \delta < (2\alpha + 2)p - 1, \quad \delta < (2\alpha + \beta + 1)p - 1.$$

It follows that Theorem 1.5 gives sharp conditions when $\beta \geq 1$ or when $\beta \leq 0$ and $[-\beta \in \mathbb{N}$ or $\alpha + \beta > 1/2]$ or when $\beta \in (0, 1)$ and $\alpha \leq -1/2$. For $\beta \leq 0$ and $-\beta \notin \mathbb{N}$ and $\alpha + \beta \leq 1/2$, Theorem 1.5 is sharp up to a possible weakening the strict inequality in the lower bound for δ in (1.6); this remains to be settled. In case $\beta \in (0, 1)$ and $\alpha > -1/2$, the upper bound for δ in (1.5) is sharp, while optimality of the lower one remains an open question. See also Remark 4.3 at the end.

Note that in Theorem 1.5 condition (1.3) is meaningful only when $\alpha + \beta < 1/2$. Further, considering the lower bound in (1.5), notice that

$$\frac{-\beta}{[\beta \vee (\alpha + \beta + 1/2)] \wedge 1} = \begin{cases} -\beta, & \alpha + \beta \geq 1/2, \\ \frac{-\beta}{\alpha + \beta + 1/2}, & \alpha + \beta < 1/2 \text{ and } \alpha > -1/2, \\ -1, & \alpha + \beta < 1/2 \text{ and } \alpha \leq -1/2. \end{cases}$$

Observe also that (1.4) is in fact the A_p condition for the power weight $x^{\delta-(2\alpha+1)}$ in the context of the space of homogeneous type $(\mathbb{R}_+, d\mu_\alpha, |\cdot|)$.

In the circumstances of Theorem 1.5, assuming in addition that $2\alpha + \beta > -1$, for each p satisfying (1.3) there is always a non-trivial interval of δ for which the $L^p(x^\delta dx)$ boundedness holds. On the other hand, when $2\alpha + \beta \leq -1$ (i.e., (α, β) is inside or on the rightmost side of the small triangle with vertices $(-1/2, 0)$, $(-1, 1/2)$, $(-1, 1)$) Theorem 1.5 gives no $L^p(x^\delta dx)$ boundedness (the range of δ is empty for each p).

Taking $\delta = \gamma + 2\alpha + 1$ and $\alpha = n/2 - 1$, $n \geq 2$ in Theorem 1.5, and either $\beta = 0$ or $\beta = (3 - n)/2$, one recovers Theorems 1.3 and 1.4, respectively. Furthermore, specifying Theorem 1.5 to the natural weight we set $\delta = 2\alpha + 1$ and get the following.

Corollary 1.7 *Assume that $\alpha > -1$ and $\alpha + \beta > -1/2$. Let $1 < p < \infty$. Then, $M_*^{\alpha, \beta}$ is bounded on $L^p(\mathbb{R}_+, d\mu_\alpha)$ provided that condition (1.3) is satisfied and*

$$\frac{-\beta}{2\alpha + 1} < \frac{1}{p} < 1 - \frac{1 - \beta}{2\alpha + 2}, \quad (1.7)$$

with the first inequality weakened if $-\beta \in \mathbb{N}$ or $\alpha + \beta > 1/2$. These conditions are optimal up to a possible weakening of the strict inequality in the lower bound in (1.7) when $0 < -\beta \notin \mathbb{N}$ and $\alpha + \beta \leq 1/2$.

Observe that, in the context of Corollary 1.7, condition (1.7) is always satisfied in case $\beta \geq 1$, and for $0 \leq \beta < 1$ the lower bound in (1.7) is always true. Note also that there is no p for which the boundedness holds if either $\beta \in (0, 1)$ and $2\alpha + \beta + 1 \leq 0$ or $\beta < 0$ and $\alpha + \beta \leq -1/4 - 1/(16\alpha + 12)$. Otherwise, there is always a non-trivial interval of p for which $M_*^{\alpha, \beta}$ is bounded on $L^p(d\mu_\alpha)$.

Taking $\alpha = n/2 - 1$, $n \geq 2$, in Corollary 1.7 we obtain

Corollary 1.8 *Let $n \geq 2$ and $1 < p < \infty$. Then, M_*^β is bounded on $L_{rad}^p(\mathbb{R}^n)$ provided that*

$$p \leq \frac{2n-1}{n-1} \text{ and } \beta > 1 - n + n/p, \text{ or } p > \frac{2n-1}{n-1} \text{ and } \beta > \frac{1-n}{p}, \quad (1.8)$$

with the last inequality weakened when $-\beta \in \mathbb{N}$ or $\beta > \frac{3-n}{2}$. This condition is optimal up to a possible weakening of the last inequality in (1.8) in case $-\beta \notin \mathbb{N}$ and $\beta \leq \frac{3-n}{2}$.

We believe that weakening the inequality in Corollary 1.8 reflects radial improvement in mapping properties of M_*^β . On the other hand, the fact that for $2 < p < 2(n+1)/(n-1)$ condition (1.8) is strictly less restrictive than condition (1.2), see Fig. 1 as follows, most probably indicates non-optimality of the yet known results on L^p -boundedness of M_*^β stated in Theorem 1.2; cf. [6, Section 3, Problem (1)].

Taking into account Corollary 1.8 and known sharp results on boundedness of M_*^0 on $L^p(\mathbb{R}^n)$, we state the following.

Conjecture 1.9 *Let $n \geq 2$ and $1 < p < \infty$. The operator M_*^β is bounded on $L^p(\mathbb{R}^n)$ if and only if condition (1.8) is satisfied.*

Corollary 1.11 *Let α, β and p be as in Proposition 1.10. Then,*

$$M_t^{\alpha, \beta} f(x) \rightarrow f(x) \text{ for a.a. } x \in \mathbb{R}_+ \text{ as } t \rightarrow 0^+$$

for any locally integrable function f on $\mathbb{R}_+$ in case $\alpha + \beta \geq 1/2$, or for any f locally in $L^p(\mathbb{R}_+, dx)$ with p satisfying condition (1.3) in case $\alpha + \beta < 1/2$.

Consequently, we obtain almost everywhere convergence to initial data results in contexts of differential problems whose solutions express via $M_t^{\alpha, \beta}$. In particular, this pertains to the following Cauchy initial-value problems, see, e.g., [2, Section 7].

- (i) Euler–Poisson–Darboux equation in $\mathbb{R}^n$ with a locally integrable/locally in L^p radial initial position and null initial speed.
- (ii) Wave equation in $\mathbb{R}^n$ with null initial position and a locally integrable radial initial speed (this case was already treated before, see [5, Section 5]).
- (iii) Euler–Poisson–Darboux equation based on the one-dimensional Bessel operator with a locally integrable/locally in L^p initial position and null initial speed.
- (iv) Wave equation based on the one-dimensional Bessel operator with null initial position and a locally integrable/locally in L^p initial speed.

In (i), (iii) and (iv) by “locally integrable/locally in L^p ” we mean local integrability in case $\alpha + \beta \geq 1/2$, and being locally in L^p for some p satisfying condition (1.3) in case $-1/2 < \alpha + \beta < 1/2$.

Structure of the paper

The rest of the paper is organized as follows. Section 2 is a technical preparation for the proof of Theorem 1.5. It provides estimates of the integral kernel $K_t^{\alpha, \beta}(x, z)$, decomposition of the (t, x, z) space into suitable regions and definitions of special auxiliary operators together with description of their L^p -behavior. Section 3 contains the proof of Theorem 1.5. The main part is preceded by establishing suitable control in terms of the special operators. At the end of Sect. 3, a brief justification of Proposition 1.10 is given. Finally, in Sect. 4 we construct suitable counterexamples in order to prove Proposition 1.6.

Notation

Throughout the paper, we use a fairly standard notation. Thus, $\mathbb{N} = \{0, 1, 2, \dots\}$ and $\mathbb{R}_+ = (0, \infty)$. The symbols “ $\vee$ ” and “ $\wedge$ ” mean the operations of taking maximum and minimum, respectively. We write $X \lesssim Y$ to indicate that $X \leq CY$ with a positive constant C independent of significant quantities. We shall write $X \simeq Y$ when simultaneously $X \lesssim Y$ and $Y \lesssim X$.

For the sake of brevity, we often omit $\mathbb{R}_+$ when denoting L^p spaces related to the measure spaces $(\mathbb{R}_+, x^\delta dx)$ and $(\mathbb{R}_+, d\mu_\alpha)$. We write $L_{\text{rad}}^p(\dots)$ for the subspace of $L^p(\dots)$ consisting of radial functions.

2 Technical preparation

Define the main regions

$$\begin{aligned} E &= \{(t, x, z) \in \mathbb{R}_+^3 : |x - z| < t < x + z\}, \\ F &= \{(t, x, z) \in \mathbb{R}_+^3 : x + z < t\}. \end{aligned}$$

Following the strategy used in [5], see also [4], we shall estimate the kernel $K_t^{\alpha,\beta}(x, z)$ and then split the bounds according to suitably defined sub-regions of E and F . Then, we will estimate the resulting maximal operators independently and get control in terms of a number of special operators whose mapping properties are essentially known. Altogether, this will give an overall control of the maximal operator $M_*^{\alpha,\beta}$.

In preparatory Sects. 2.1–2.3 as follows, we gather the kernel estimates, suitable decompositions of E and F and definitions of the special operators, respectively.

2.1 Kernel estimates

We first rephrase [2, Theorem 3.3] in a way more convenient for our present purposes; see also [2, Section 2.3]. In particular, we neglect parts concerning sharpness of the bounds for certain α and β .

Theorem 2.1 [2] *Assume that $\alpha > -1$ and $\alpha + \beta > -1/2$. Let $t, x, z > 0$. The kernel $K_t^{\alpha,\beta}(x, z)$ vanishes when $t < |x - z|$, whereas in $E \cup F$ the following uniform estimates hold.*

(1) *If $-\beta \in \mathbb{N}$, then the kernel vanishes in F and*

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \frac{(xz)^{-2\alpha-\beta}}{t^{2\alpha+2\beta}} \left([t^2 - (x-z)^2][(x+z)^2 - t^2] \right)^{\alpha+\beta-1/2} \quad \text{in } E.$$

(2) *If $\alpha + \beta > 1/2$ and neither $-\beta \in \mathbb{N}$ nor $2\alpha + \beta = 0$, then*

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \begin{cases} \frac{(xz)^{-\alpha-1/2}}{t^{2\alpha+2\beta}} [t^2 - (x-z)^2]^{\alpha+\beta-1/2} & \text{in } E, \\ \frac{1}{t^{2\alpha+2\beta}} [t^2 - (x-z)^2]^{\beta-1} & \text{in } F. \end{cases}$$

(3) *If $\alpha + \beta > 1/2$ and $2\alpha + \beta = 0$, then*

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \frac{1}{t^{-2\alpha}} \left([t^2 - (x-z)^2] |t^2 - (x+z)^2| \right)^{-\alpha-1/2} \quad \text{in } E \cup F.$$

(4) *If $\alpha + \beta = 1/2$ and neither $-\beta \in \mathbb{N}$ nor $\beta = 1$, then*

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \begin{cases} \frac{(xz)^{-\alpha-1/2}}{t} \log \frac{8xz}{(x+z)^2 - t^2} & \text{in } E, \\ \frac{1}{t} [t^2 - (x-z)^2]^{-\alpha-1/2} \log \left(2 \frac{t^2 - (x-z)^2}{t^2 - (x+z)^2} \right) & \text{in } F. \end{cases}$$

(5) *If $\alpha + \beta = 1/2$ and $\beta = 1$, then*

$$K_t^{\alpha,\beta}(x, z) \simeq \frac{1}{t} \quad \text{in } E \cup F.$$

(6) *If $\alpha + \beta < 1/2$ and neither $-\beta \in \mathbb{N}$ nor $\beta = 1$ nor $\alpha + 1/2 \in \mathbb{N}$, then*

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \begin{cases} \frac{(xz)^{-2\alpha-\beta}}{t^{2\alpha+2\beta}} \left([t^2 - (x-z)^2][(x+z)^2 - t^2] \right)^{\alpha+\beta-1/2} & \text{in } E, \\ \frac{1}{t^{2\alpha+2\beta}} [t^2 - (x-z)^2]^{-\alpha-1/2} [t^2 - (x+z)^2]^{\alpha+\beta-1/2} & \text{in } F. \end{cases}$$

(7) *If $\alpha + \beta < 1/2$ and $\beta = 1$, then*

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \begin{cases} \frac{(xz)^{-2\alpha-1}}{t^{2\alpha+2}} \left([t^2 - (x-z)^2][(x+z)^2 - t^2] \right)^{\alpha+1/2} & \text{in } E, \\ \frac{1}{t^{2\alpha+2}} & \text{in } F. \end{cases}$$

(8) If $\alpha + \beta < 1/2$ and $\alpha + 1/2 \in \mathbb{N}$, then

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \begin{cases} \frac{(xz)^{-\alpha-1/2}}{t^{2\alpha+2\beta}} [t^2 - (x-z)^2]^{\alpha+\beta-1/2} & \text{in } E, \\ \frac{1}{t^{2\alpha+2\beta}} [t^2 - (x-z)^2]^{-\alpha-1/2} [t^2 - (x+z)^2]^{\alpha+\beta-1/2} & \text{in } F. \end{cases}$$

The above theorem distinguishes cases, or rather segments and lines in the (α, β) plane, where the estimates are better comparing to other neighboring (α, β) . However, this does not seem to be reflected in power-weighted L^p , $1 < p < \infty$, boundedness of $M_*^{\alpha,\beta}$. This claim is based on our detailed analysis of the maximal operator in each of the cases related to items (1)–(8) of Theorem 2.1. Even though we could often obtain seemingly better control of $K_*^{\alpha,\beta}$ in terms of the auxiliary special operators, it led us to the same mapping properties as stated in Theorem 1.5. Therefore, we simplify things already at this stage and derive less accurate kernel bounds, but with simpler structure (in particular, containing no logarithmic expressions), which will be sufficient for our purpose. However, this strategy might not be appropriate for studying more subtle mapping properties of $M_*^{\alpha,\beta}$, like weak or restricted weak-type estimates.

For notational convenience, define auxiliary kernels

$$\begin{aligned} \Phi_t^{\alpha,\beta}(x, z) &= \chi_E(t, x, z) \frac{(xz)^{-\alpha-1/2}}{t^{2\alpha+2\beta}} [t^2 - (x-z)^2]^{\alpha+\beta-1/2} \\ &\quad + \chi_F(t, x, z) \frac{1}{t^{2\alpha+2\beta}} [t^2 - (x-z)^2]^{\beta-1}, \\ \Psi_t^{\alpha,\beta}(x, z) &= \chi_E(t, x, z) \frac{(xz)^{-2\alpha-\beta}}{t^{2\alpha+2\beta}} \left([t^2 - (x-z)^2] [(x+z)^2 - t^2] \right)^{\alpha+\beta-1/2} \\ &\quad + \chi_F(t, x, z) \frac{1}{t^{2\alpha+2\beta}} [t^2 - (x-z)^2]^{-\alpha-1/2} [t^2 - (x+z)^2]^{\alpha+\beta-1/2}. \end{aligned}$$

Theorem 2.2 Assume that $\alpha > -1$ and $\alpha + \beta > -1/2$. Fix an arbitrary $\varepsilon > 0$. The following estimates hold uniformly in $(t, x, z) \in E \cup F$:

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \begin{cases} \Phi_t^{\alpha,\beta}(x, z), & \text{if } \alpha + \beta > 1/2, \\ \Psi_t^{\alpha,\beta-\varepsilon}(x, z), & \text{if } \alpha + \beta = 1/2, \\ \Psi_t^{\alpha,\beta}(x, z), & \text{if } \alpha + \beta < 1/2, \\ \Psi_t^{\alpha,\beta}(x, z) \chi_E(t, x, z), & \text{if } -\beta \in \mathbb{N}. \end{cases}$$

Moreover, $\chi_E(t, x, z) \Psi_t^{\alpha,\beta}(x, z) \lesssim \chi_E(t, x, z) \Phi_t^{\alpha,\beta}(x, z)$ when $\alpha + \beta \geq 1/2$.

Proof All the asserted bounds are straightforward consequences of Theorem 2.1. Clearly,

$$|K_t^{\alpha,\beta}(x, z)| \lesssim \chi_E(t, x, z) \Psi_t^{\alpha,\beta}(x, z) \quad \text{if } -\beta \in \mathbb{N},$$

for $(t, x, z) \in E \cup F$, is just Theorem 2.1(1). We can write

$$\chi_E(t, x, z) \Psi_t^{\alpha,\beta}(x, z) = \chi_E(t, x, z) \Phi_t^{\alpha,\beta}(x, z) \left[\frac{(x+z)^2 - t^2}{xz} \right]^{\alpha+\beta-1/2}$$

and, since the expression in square brackets is bounded from above by a constant, we infer that

$$\chi_E(t, x, z) \Psi_t^{\alpha,\beta}(x, z) \lesssim \chi_E(t, x, z) \Phi_t^{\alpha,\beta}(x, z) \quad \text{when } \alpha + \beta \geq 1/2. \quad (2.1)$$

Next, assume that $\alpha + \beta > 1/2$ and $2\alpha + \beta = 0$ (see Theorem 2.1(3)), which forces $\alpha < -1/2$. For $(t, x, z) \in E \cup F$, we have

$$\begin{aligned} & \frac{1}{t^{-2\alpha}} \left([t^2 - (x - z)^2] |t^2 - (x + z)^2| \right)^{-\alpha-1/2} \\ &= \chi_E(t, x, z) \Phi_t^{\alpha, \beta}(x, z) \left[\frac{xz}{(x + z)^2 - t^2} \right]^{\alpha+1/2} \\ & \quad + \chi_F(t, x, z) \Phi_t^{\alpha, \beta}(x, z) \left[\frac{t^2 - (x - z)^2}{t^2 - (x + z)^2} \right]^{\alpha+1/2}. \end{aligned}$$

Here, the expressions in square brackets are bounded from below by a positive constant. Hence, in view of Theorem 2.1(1)–(3) and (2.1), we see that

$$|K_t^{\alpha, \beta}(x, z)| \lesssim \Phi_t^{\alpha, \beta}(x, z) \quad \text{if } \alpha + \beta > 1/2.$$

Let now $\alpha + \beta = 1/2$ and fix $\varepsilon > 0$. For $(t, x, z) \in E$, we have (see Theorem 2.1(4))

$$\begin{aligned} & \frac{(xz)^{-\alpha-1/2}}{t} \log \frac{8xz}{(x + z)^2 - t^2} \\ & \lesssim \frac{(xz)^{-\alpha-1/2}}{t} \left[\frac{(x + z)^2 - t^2}{xz} \right]^{\alpha+\beta-\varepsilon-1/2} \\ & = \Psi_t^{\alpha, \beta-\varepsilon}(x, z) \left[\frac{t^2}{t^2 - (x - z)^2} \right]^{\alpha+\beta-\varepsilon-1/2} \leq \Psi_t^{\alpha, \beta-\varepsilon}(x, z), \end{aligned}$$

where we used the fact that $\log y$ grows slower than the positive power $y^{-(\alpha+\beta-\varepsilon-1/2)}$ (here y positive and separated from zero). Further, for $(t, x, z) \in F$,

$$\begin{aligned} & \frac{1}{t} [t^2 - (x - z)^2]^{-\alpha-1/2} \log \left(2 \frac{t^2 - (x - z)^2}{t^2 - (x + z)^2} \right) \\ & \lesssim \frac{1}{t} [t^2 - (x - z)^2]^{-\alpha-1/2} \left[\frac{t^2 - (x + z)^2}{t^2 - (x - z)^2} \right]^{\alpha+\beta-\varepsilon-1/2} \\ & = \Psi_t^{\alpha, \beta-\varepsilon}(x, z) \left[\frac{t^2}{t^2 - (x - z)^2} \right]^{\alpha+\beta-\varepsilon-1/2} \leq \Psi_t^{\alpha, \beta-\varepsilon}(x, z). \end{aligned}$$

This shows that (see Theorem 2.1(4), (5))

$$|K_t^{\alpha, \beta}(x, z)| \lesssim \Psi_t^{\alpha, \beta-\varepsilon}(x, z) \quad \text{if } \alpha + \beta = 1/2$$

for $(t, x, z) \in E \cup F$ (to be precise, the case coming from Theorem 2.1(5), i.e., $(\alpha, \beta) = (-1/2, 1)$, is trivially included with no logarithms involved).

Finally, we consider $\alpha + \beta < 1/2$, see Theorem 2.1(6)–(8). If $\beta = 1$, then automatically $\alpha < -1/2$ and for $(t, x, z) \in F$ we have (see Theorem 2.1(7))

$$\frac{1}{t^{2\alpha+2}} = \Psi_t^{\alpha, \beta}(x, z) \left[\frac{t^2 - (x - z)^2}{t^2 - (x + z)^2} \right]^{\alpha+1/2} \leq \Psi_t^{\alpha, \beta}(x, z).$$

If $\alpha + 1/2 \in \mathbb{N}$, then for $(t, x, z) \in E$ (see Theorem 2.1(8))

$$\frac{(xz)^{-\alpha-1/2}}{t^{2\alpha+2\beta}} [t^2 - (x - z)^2]^{\alpha+\beta-1/2} = \Psi_t^{\alpha, \beta}(x, z) \left[\frac{(x + z)^2 - t^2}{xz} \right]^{-(\alpha+\beta-1/2)}.$$

Altogether, this implies for $(t, x, z) \in E \cup F$

$$|K_t^{\alpha, \beta}(x, z)| \lesssim \Psi_t^{\alpha, \beta}(x, z) \quad \text{if } \alpha + \beta < 1/2.$$

The proof of Theorem 2.2 is complete. $\square$

2.2 Decompositions of E and F

Inspired by the analysis presented in [5], we define the following sub-regions of E and F , see Fig. 2 as follows,

$$\begin{aligned} E_1 &= E \cap \{(t, x, z) \in \mathbb{R}_+^3 : t < x/2\}, \\ E_2 &= E \cap \{(t, x, z) \in \mathbb{R}_+^3 : x/2 \leq t < 3x\}, \\ E_3 &= E \cap \{(t, x, z) \in \mathbb{R}_+^3 : 3x \leq t\} \end{aligned}$$

and

$$\begin{aligned} F_1 &= F \cap \{(t, x, z) \in \mathbb{R}_+^3 : x < t < 3x\}, \\ F_2 &= F \cap \{(t, x, z) \in \mathbb{R}_+^3 : 3x \leq t\}, \\ F'_2 &= F_2 \cap \{(t, x, z) \in \mathbb{R}_+^3 : z < (t-x)/2\}, \\ F''_2 &= F_2 \cap \{(t, x, z) \in \mathbb{R}_+^3 : (t-x)/2 \leq z < t-x\}. \end{aligned}$$

Clearly, $E = E_1 \cup E_2 \cup E_3$, $F = F_1 \cup F_2$, $F_2 = F'_2 \cup F''_2$, all the sums being disjoint.

In what follows, we will use uniform estimates of expressions in t, x, z that hold specifically in one (or more) of the above regions, cf. [5]. We list them as follows for an easy further reference. Verification of these relations is straightforward and left to the reader.

$$\text{in } E_1 : \quad x \simeq z, \quad (2.2)$$

$$\text{in } E_2 : \quad t^2 - (x - z)^2 \lesssim tz, \quad (2.3)$$

Fig. 2 Sections of regions E_i and F_i given $x > 0$ fixed

$$\text{in } E_3 : \quad t^2 - (x - z)^2 \lesssim tx, \quad z \simeq t, \quad (2.4)$$

$$\text{in } F_1 : \quad t^2 - (x - z)^2 \simeq x(t - x) \simeq x(t - x + z), \quad t^2 - (x + z)^2 \simeq x(t - x - z), \quad (2.5)$$

$$\text{in } F_2 : \quad t^2 - (x - z)^2 \simeq t(t + x - z), \quad t^2 - (x + z)^2 \simeq t(t - x - z), \quad (2.6)$$

$$\text{in } F'_2 : \quad t^2 - (x \pm z)^2 \simeq t^2, \quad (2.7)$$

$$\text{in } F''_2 : \quad z \simeq t \simeq t - x \simeq t - (x - z), \quad t + x - z \simeq t - z. \quad (2.8)$$

2.3 Special operators

We will use the following auxiliary operators whose mapping properties are essentially known, see [4,5] and also references given there.

A variant of the Hardy–Littlewood maximal operator (denoted by E_2 in [4])

$$Df(x) = \sup_{0 \leq a < x < b} \frac{1}{b^2 - a^2} \int_a^b z |f(z)| \, dz.$$

The operator D is bounded on $L^p(x^\gamma dx)$, $1 < p < \infty$, if and only if $-1 < \gamma < 2p - 1$.

The local Hardy–Littlewood maximal operator

$$Lf(x) = \sup_{t < x/2} \frac{1}{2t} \int_{x-t}^{x+t} |f(z)| \, dz.$$

The operator L is bounded on $L^p(x^\gamma dx)$, $1 < p < \infty$, for any $\gamma \in \mathbb{R}$.

The Hardy-type operator

$$H_\eta f(x) = \frac{1}{x^\eta} \int_0^x z^{\eta-1} f(z) \, dz.$$

The operator H_η is bounded on $L^p(x^\gamma dx)$, $1 < p < \infty$, if and only if $\gamma < \eta p - 1$.

The remote maximal operator

$$Rf(x) = \sup_{t > 2x} \frac{1}{2x} \int_{t-x}^{t+x} |f(z)| \, dz.$$

The operator R is bounded on $L^p(x^\gamma dx)$, $1 < p < \infty$, if and only if $\gamma \geq 0$.

The maximal operator N . For positive η , let

$$N_\eta f(x) = \sup_{t > x} \frac{1}{t^\eta} \int_0^t z^{\eta-1} |f(z)| \, dz.$$

The operator N_η is bounded on $L^p(x^\gamma dx)$, $1 < p < \infty$, if $-1 < \gamma < \eta p - 1$. At this point, it is perhaps in order to note that the range given in [5, p. 1545] for the boundedness of N_η is wrong; consequently some statements in the proof of [5, Theorem 4.3] are not correct, nevertheless the result claimed there is true.

The maximal operator T . For any $\eta \in \mathbb{R}$, consider the operator

$$T_\eta f(x) = \sup_{t > 2x} \int_{t/2}^t \frac{z^{\eta-1} |f(z)|}{(t - z + x)^\eta} \, dz. \quad (2.9)$$

Boundedness of T_η on power-weighted L^p spaces is shown in the following lemma, which is a simple extension of [5, Lemma 2.1]. Actually, this is even more than we need; we state item (b) only for the sake of completeness.

Lemma 2.3 *Let $1 < p < \infty$.*

- (a) *For $\eta > 1$, T_η is bounded on $L^p(x^\gamma dx)$ for $\gamma \geq (\eta - 1)p$.*
- (b) *For $\eta = 1$, T_η is bounded on $L^p(x^\gamma dx)$ for $\gamma > 0$.*
- (c) *For $0 < \eta < 1$, T_η is bounded on $L^p(x^\gamma dx)$ for $\gamma \geq \eta - 1$.*
- (d) *For $\eta \leq 0$, T_η is bounded on $L^p(x^\gamma dx)$ for $\gamma > -1$.*

Proof Items (a) and (c) are proved in [5], while (b) is commented there.

To show (d), assume that $\eta \leq 0$ and observe that

$$T_\eta f(x) = \sup_{t > 2x} \int_{t/2}^t \left(\frac{t - z + x}{z} \right)^{-\eta} \frac{|f(z)|}{z} dz \lesssim \sup_{t > 2x} \int_{t/2}^t \frac{|f(z)|}{z} dz \leq \int_x^\infty \frac{|f(z)|}{z} dz.$$

Thus, T_η is controlled in terms of the dual Hardy operator $f(x) \mapsto \int_x^\infty \frac{f(z)}{z} dz$, which is well known to be bounded on $L^p(x^\gamma dx)$ if and only if $\gamma > -1$; see, e.g., [2, Lemma 6.3]. $\square$

3 Proof of the main theorem

From now on, we will consider only nonnegative functions f . This will be enough, since our approach is based on absolute estimates of the kernel.

Let $G \subset \mathbb{R}_+^3$. We denote

$$\begin{aligned} \Phi_{*,G}^{\alpha,\beta} f(x) &= \sup_{t>0} \int_0^\infty \chi_G(t, x, z) \Phi_t^{\alpha,\beta}(x, z) f(z) d\mu_\alpha(z), \\ \Phi_*^{\alpha,\beta} f(x) &= \Phi_{*,\mathbb{R}_+^3}^{\alpha,\beta} f(x) \end{aligned}$$

and analogously for the Ψ counterparts. Clearly, $\Phi_*^{\alpha,\beta} f \leq \Phi_{*,E}^{\alpha,\beta} f + \Phi_{*,F}^{\alpha,\beta} f$ and similarly in case of the Ψ operators.

3.1 Control of $\Phi_*^{\alpha,\beta}$

In this subsection, we will establish a control of $\Phi_*^{\alpha,\beta}$ in terms of the special operators defined in Sect. 2.3. This will be done under the assumption $\alpha + \beta \geq 1/2$.

Lemma 3.1 *Let $\alpha > -1$ and $\alpha + \beta \geq 1/2$. Then,*

$$\begin{aligned} \Phi_{*,E}^{\alpha,\beta} f(x) &\lesssim Lf(x) + H_{2\alpha+\beta+1} f(4x) + x^\beta R(z^{-\beta} f)(x), \\ \Phi_{*,F}^{\alpha,\beta} f(x) &\lesssim \chi_{\{\beta \geq 1\}} H_{2\alpha+2} f(2x) + \chi_{\{\beta < 1\}} H_{2\alpha+\beta+1} f(2x) + N_{2\alpha+2} f(x) + T_{1-\beta} f(x), \end{aligned}$$

uniformly in $f \geq 0$ and $x > 0$.

Proof We will bound the maximal operators related to restrictions of $\Phi_t^{\alpha,\beta}(x, z)$ to the subregions of E and F specified in Sect. 2.2.

Region E_1 . In view of (2.2), taking into account that $\alpha + \beta - 1/2 \geq 0$, we have

$$\begin{aligned} \Phi_{*,E_1}^{\alpha,\beta} f(x) &\lesssim \sup_{t < x/2} \frac{x^{-\alpha-1/2}}{t^{2\alpha+2\beta}} \int_{x-t}^{x+t} [t^2 - (x-z)^2]^{\alpha+\beta-1/2} f(z) z^{\alpha+1/2} dz \\ &\lesssim \sup_{t < x/2} \frac{1}{t} \int_{x-t}^{x+t} f(z) dz = 2Lf(x). \end{aligned}$$

Region E_2 . In this region $t \simeq x$, so using (2.3) and $\alpha + \beta - 1/2 \geq 0$, we get

$$\Phi_{*,E_2}^{\alpha,\beta} f(x) \lesssim \sup_{x/2 \leq t < 3x} \frac{1}{x^{2\alpha+\beta+1}} \int_{|x-t|}^{x+t} z^{2\alpha+\beta} f(z) dz \leq H_{2\alpha+\beta+1} f(4x).$$

Region E_3 . Using (2.4) and $\alpha + \beta - 1/2 \geq 0$ gives

$$\Phi_{*,E_3}^{\alpha,\beta} f(x) \lesssim \sup_{t \geq 3x} x^\beta \frac{1}{x} \int_{t-x}^{t+x} z^{-\alpha-\beta-1/2} f(z) z^{\alpha+1/2} dz \lesssim x^\beta R(z^{-\beta} f)(x).$$

Region F_1 . Now, $t \simeq x$. Using the fact that $t + x - z \simeq x$ in F_1 , see (2.5), we can write

$$\Phi_{*,F_1}^{\alpha,\beta} f(x) \lesssim \sup_{x < t < 3x} \frac{1}{x^{2\alpha+\beta+1}} \int_0^{t-x} (t-x+z)^{\beta-1} f(z) z^{2\alpha+1} dz.$$

Here, the factor $(t-x+z)^{\beta-1}$ is controlled by $(t-x)^{\beta-1} \lesssim x^{\beta-1}$ when $\beta \geq 1$, and by $z^{\beta-1}$ otherwise. This leads directly to the bound

$$\Phi_{*,F_1}^{\alpha,\beta} f(x) \lesssim \chi_{\{\beta \geq 1\}} H_{2\alpha+2} f(2x) + \chi_{\{\beta < 1\}} H_{2\alpha+\beta+1} f(2x).$$

Region F'_2 . In view of (2.7),

$$\Phi_{*,F'_2}^{\alpha,\beta} f(x) \lesssim \sup_{t \geq 3x} \frac{1}{t^{2\alpha+2}} \int_0^{(t-x)/2} f(z) z^{2\alpha+1} dz \leq N_{2\alpha+2} f(3x) \leq N_{2\alpha+2} f(x).$$

Region F''_2 . Here, we use (2.8) and (2.6) obtaining

$$\Phi_{*,F''_2}^{\alpha,\beta} f(x) \lesssim \sup_{t \geq 3x} \int_{(t-x)/2}^{t-x} \frac{z^{-\beta} f(z)}{(t-z)^{1-\beta}} dz = \sup_{t \geq 2x} \int_{t/2}^t \frac{z^{-\beta} f(z)}{(t+x-z)^{1-\beta}} dz = T_{1-\beta} f(x).$$

The bounds of Lemma 3.1 follow. $\square$

3.2 Control of $\Psi_{*,E}^{\alpha,\beta}$

We will prove the following.

Lemma 3.2 *Let $\alpha > -1$ and $-1/2 < \alpha + \beta < 1/2$. Then, given any $\theta > 0$ such that $\frac{1}{1+\theta} < \alpha + \beta + 1/2$,*

$$\Psi_{*,E}^{\alpha,\beta} f(x) \lesssim x^\beta [D(z^{-\beta} f)^{1+\theta}(x)]^{\frac{1}{1+\theta}} + x^\beta [R(z^{-\beta} f)^{1+\theta}(x)]^{\frac{1}{1+\theta}},$$

uniformly in $f \geq 0$ and $x > 0$.

Proof We first reduce the task to the special case $\beta = 0$. Observe that

$$\Psi_{*,E}^{\alpha,\beta} f(x) = \sup_{t>0} \frac{x^\beta}{(tx)^{2\alpha+2\beta}} \int_{|x-t|}^{x+t} \left([t^2 - (x-z)^2][(x+z)^2 - t^2] \right)^{\alpha+\beta-1/2} z^{-\beta} f(z) z dz.$$

Thus,

$$\Psi_{*,E}^{\alpha,\beta} f(x) = x^\beta \Psi_{*,E}^{\alpha+\beta,0} (z^{-\beta} f)(x)$$

and hence it is enough to show that, given $-1/2 < \alpha < 1/2$,

$$\Psi_{*,E}^{\alpha,0} f(x) \lesssim [Df^{1+\theta}(x)]^{\frac{1}{1+\theta}} + [Rf^{1+\theta}(x)]^{\frac{1}{1+\theta}} \quad (3.1)$$

with any fixed $\theta > 0$ such that $\frac{1}{1+\theta} < \alpha + 1/2$.

The proof of (3.1) is a straightforward generalization of the reasoning from [4], see the proof of [4, Theorem 3.1(b)]. We present briefly some details for the reader's convenience. Using the identity

$$[t^2 - (x - z)^2][(x + z)^2 - t^2] = [(t + x)^2 - z^2][z^2 - (x - t)^2] \quad (3.2)$$

and then Hölder's inequality, one gets the bound

$$\begin{aligned} \Psi_{*,E}^{\alpha,0} f(x) &\leq \sup_{t>0} \left[\frac{1}{(xt)^{2\alpha}} \left(\int_{|t-x|}^{t+x} z f^{1+\theta}(z) dz \right)^{\frac{1}{1+\theta}} \right. \\ &\quad \times \left. \left(\int_{|t-x|}^{t+x} \left[(t+x)^2 - z^2 \right] [z^2 - (x-t)^2] \right)^{(2\alpha-1)\frac{1+\theta}{2\theta}} z dz \right]^{\frac{\theta}{1+\theta}}. \end{aligned}$$

The second integral here converges when $(2\alpha - 1)\frac{1+\theta}{2\theta} > -1$, i.e., when $\frac{1}{1+\theta} < \alpha + 1/2$. In such case, it is comparable to $(xt)^{(2\alpha-1)\frac{1+\theta}{\theta}+1}$, see Lemma 4.2 in Sect. 4 as follows.

Consequently, we arrive at the bound

$$\Psi_{*,E}^{\alpha,0} f(x) \lesssim \sup_{t>0} \left(\frac{1}{xt} \int_{|t-x|}^{t+x} z f^{1+\theta}(z) dz \right)^{\frac{1}{1+\theta}}.$$

From here, one proceeds by splitting the supremum into $t \leq 2x$ and $t > 2x$ (which corresponds essentially to estimating separately $\Psi_{*,E_1 \cup E_2}^{\alpha,0}$ and $\Psi_{*,E_3}^{\alpha,0}$). In the first case $|t - x| \leq x$ and we get the control by $[Df^{1+\theta}(x)]^{1/(1+\theta)}$. On the other hand, if $t > 2x$, then $z \simeq t$ and this part of the maximal operator is controlled by $[Rf^{1+\theta}(x)]^{1/(1+\theta)}$. $\square$

3.3 Control of $\Psi_{*,F}^{\alpha,\beta}$

We finally establish a control of $\Psi_{*,F}^{\alpha,\beta}$ in terms of the special operators.

Lemma 3.3 *Let $\alpha > -1$ and $-1/2 < \alpha + \beta < 1/2$. Then, given any $\theta > 0$ such that $\frac{1}{1+\theta} < \alpha + \beta + 1/2$,*

$$\begin{aligned} \Psi_{*,F}^{\alpha,\beta} f(x) &\lesssim \chi_{\{\beta \geq 1\}} [H_{(\alpha+1/2+1-\beta)(1+\theta)+1}(f^{1+\theta})(2x)]^{\frac{1}{1+\theta}} \\ &\quad + \chi_{\{\beta < 1\}} [H_{(\alpha+1/2)(1+\theta)+1}(f^{1+\theta})(2x)]^{\frac{1}{1+\theta}} \\ &\quad + N_{2\alpha+2} f(x) + [T_{(\alpha+1/2)(1+\theta)}(f^{1+\theta})(x)]^{\frac{1}{1+\theta}}, \end{aligned}$$

uniformly in $f \geq 0$ and $x > 0$.

Proof We will bound suitably the restricted maximal operators $\Psi_{*,F_1}^{\alpha,\beta}$, $\Psi_{*,F_2}^{\alpha,\beta}$ and $\Psi_{*,F_2'}^{\alpha,\beta}$.

Region F_1 . Here, $t \simeq x$. Then, by (2.5), we have

$$\Psi_{*,F_1}^{\alpha,\beta} f(x) \lesssim \sup_{x < t < 3x} \frac{(t-x)^{-\alpha-1/2}}{x^{2\alpha+\beta+1}} \int_0^{t-x} (t-x-z)^{\alpha+\beta-1/2} f(z) z^{2\alpha+1} dz.$$

By Hölder's inequality, the last integral is not bigger than

$$\left(\int_0^{t-x} [z^{\alpha+1/2} f(z)]^{1+\theta} dz \right)^{\frac{1}{1+\theta}} \left(\int_0^{t-x} [z^{\alpha+1/2} (t-x-z)^{\alpha+\beta-1/2}]^{\frac{1+\theta}{\theta}} dz \right)^{\frac{\theta}{1+\theta}}.$$

Assuming that $\beta < 1$, the second integral here is finite if and only if $\frac{1}{1+\theta} < \alpha + \beta + 1/2$. If this is the case, it is comparable with $(t-x)^{(2\alpha+\beta)\frac{1+\theta}{\theta}+1}$, as verified by changing the variable $z \mapsto (t-x)z$. This gives

$$\begin{aligned}\Psi_{*,F_1}^{\alpha,\beta} f(x) &\lesssim \sup_{x < t < 3x} \frac{(t-x)^{\alpha+\beta-1/2+\frac{\theta}{1+\theta}}}{x^{2\alpha+\beta+1}} \left(\int_0^{t-x} [z^{\alpha+1/2} f(z)]^{1+\theta} dz \right)^{\frac{1}{1+\theta}} \\ &\lesssim \frac{1}{x^{\alpha+1/2+\frac{1}{1+\theta}}} \left(\int_0^{2x} [z^{\alpha+1/2} f(z)]^{1+\theta} dz \right)^{\frac{1}{1+\theta}} \\ &= [H_{(\alpha+1/2)(1+\theta)+1}(f^{1+\theta})(2x)]^{\frac{1}{1+\theta}},\end{aligned}\quad (3.3)$$

where in the second relation we used the assumption $\alpha + \beta + 1/2 > \frac{1}{1+\theta}$.

When $\beta \geq 1$ (then automatically $\alpha \leq -1/2$), we use Hölder's inequality with the splitting of $z^{2\alpha+1}$ so that $z^{\alpha+1/2+1-\beta}$ is attached to f . This leads to the bound

$$\Psi_{*,F_1}^{\alpha,\beta} f(x) \lesssim [H_{(\alpha+1/2+1-\beta)(1+\theta)+1}(f^{1+\theta})(2x)]^{\frac{1}{1+\theta}}$$

under the condition $\frac{1}{1+\theta} < \alpha + \beta + 1/2$. The assumption $\beta \geq 1$ is used in the analog of estimate (3.3), to assure that a power of $t-x$ occurring there is nonnegative.

Region F'_2 . In view of (2.7),

$$\Psi_{*,F'_2}^{\alpha,\beta} f(x) \lesssim \sup_{t \geq 3x} \frac{1}{t^{2\alpha+2}} \int_0^{(t-x)/2} f(z) z^{2\alpha+1} dz \leq N_{2\alpha+2} f(x).$$

Region F''_2 . Now, we use (2.6) and get

$$\Psi_{*,F''_2}^{\alpha,\beta} f(x) \lesssim \sup_{t \geq 3x} \frac{1}{t^{2\alpha+\beta+1}} \int_{(t-x)/2}^{t-x} (t+x-z)^{-\alpha-1/2} (t-x-z)^{\alpha+\beta-1/2} f(z) z^{2\alpha+1} dz.$$

By means of Hölder's inequality, we can control the above integral by

$$\begin{aligned}&\left(\int_{(t-x)/2}^{t-x} \left[\left(\frac{z}{t+x-z} \right)^{\alpha+1/2} f(z) \right]^{1+\theta} dz \right)^{\frac{1}{1+\theta}} \\ &\quad \times \left(\int_{(t-x)/2}^{t-x} [(t-x-z)^{\alpha+\beta-1/2} z^{\alpha+1/2}]^{\frac{1+\theta}{\theta}} dz \right)^{\frac{\theta}{1+\theta}}.\end{aligned}$$

The second integral here is finite if and only if $\frac{1}{1+\theta} < \alpha + \beta + 1/2$. In this case, it is comparable with $(t-x)^{(2\alpha+\beta)\frac{1+\theta}{\theta}+1}$. This implies, see (2.8),

$$\begin{aligned}\Psi_{*,F''_2}^{\alpha,\beta} f(x) &\lesssim \sup_{t \geq 3x} \frac{(t-x)^{2\alpha+\beta+\frac{\theta}{1+\theta}}}{t^{2\alpha+\beta+1}} \left(\int_{(t-x)/2}^{t-x} \left[\left(\frac{z}{t+x-z} \right)^{\alpha+1/2} f(z) \right]^{1+\theta} dz \right)^{\frac{1}{1+\theta}} \\ &\lesssim \sup_{u \geq 2x} \left(\int_{u/2}^u \frac{z^{(\alpha+1/2)(1+\theta)-1}}{(u+x-z)^{(\alpha+1/2)(1+\theta)}} f^{1+\theta}(z) dz \right)^{\frac{1}{1+\theta}} \\ &= [T_{(\alpha+1/2)(1+\theta)}(f^{1+\theta})(x)]^{\frac{1}{1+\theta}}.\end{aligned}$$

Now, Lemma 3.3 follows. $\square$

3.4 Proof of Theorem 1.5

We distinguish several cases described by conditions on α and β . Altogether, they imply Theorem 1.5. In what follows, we always assume that $\alpha > -1$ and $\alpha + \beta > -1/2$, $1 < p < \infty$, and $x \in \mathbb{R}_+$.

Case 1 $\alpha + \beta > 1/2$ and $-\beta \notin \mathbb{N}$. By Theorem 2.2, $M_*^{\alpha, \beta} f(x) \lesssim \Phi_*^{\alpha, \beta} f(x)$. Thus, from Lemma 3.1 we have the control

$$M_*^{\alpha, \beta} f(x) \lesssim Lf(x) + H_{2\alpha+\beta+1}f(4x) + H_{2\alpha+2}f(2x) + N_{2\alpha+2}f(x) \\ + x^\beta R(z^{-\beta}f)(x) + T_{1-\beta}f(x).$$

The operator L is bounded on $L^p(x^\delta dx)$ for any $\delta \in \mathbb{R}$. The conditions for $L^p(x^\delta dx)$ boundedness of the remaining operators controlling $M_*^{\alpha, \beta}$ are as follows, see Sect. 2.3:

$$\begin{aligned} H_{2\alpha+\beta+1} : & \quad \delta < (2\alpha + \beta + 1)p - 1, \\ H_{2\alpha+2} : & \quad \delta < (2\alpha + 2)p - 1, \\ N_{2\alpha+2} : & \quad -1 < \delta < (2\alpha + 2)p - 1, \\ f \mapsto x^\beta R(z^{-\beta}f)(x) : & \quad -\beta p \leq \delta, \\ T_{1-\beta} : & \quad \begin{cases} -1 < \delta, & \text{if } \beta \geq 1, \\ -\beta \leq \delta, & \text{if } \beta \in (0, 1), \\ -\beta p \leq \delta, & \text{if } \beta < 0. \end{cases} \end{aligned}$$

We conclude that $M_*^{\alpha, \beta}$ is bounded on $L^p(x^\delta dx)$ if δ satisfies

$$-\min(1, \beta, \beta p) < \delta < \min((2\alpha + \beta + 1)p - 1, (2\alpha + 2)p - 1),$$

with the first inequality weakened when $\beta < 1$. Here,

$$\min(1, \beta, \beta p) = \begin{cases} 1, & \text{if } \beta \geq 1, \\ \beta, & \text{if } \beta \in (0, 1), \\ \beta p, & \text{if } \beta < 0, \end{cases}$$

while

$$\min((2\alpha + \beta + 1)p - 1, (2\alpha + 2)p - 1) = \begin{cases} (2\alpha + \beta + 1)p - 1, & \text{if } \beta < 1, \\ (2\alpha + 2)p - 1, & \text{if } \beta \geq 1. \end{cases}$$

Case 2 $\alpha + \beta < 1/2$ and $-\beta \notin \mathbb{N}$. In view of Theorem 2.2, $M_*^{\alpha, \beta} f(x) \lesssim \Psi_*^{\alpha, \beta} f(x)$. Consequently, from Lemmas 3.2 and 3.3 we get the control

$$M_*^{\alpha, \beta} f(x) \lesssim x^\beta [D(z^{-\beta}f)^{1+\theta}(x)]^{\frac{1}{1+\theta}} + x^\beta [R(z^{-\beta}f)^{1+\theta}(x)]^{\frac{1}{1+\theta}} \\ + \chi_{\{\beta < 1\}} [H_{(\alpha+1/2)(1+\theta)+1}(f^{1+\theta})(2x)]^{\frac{1}{1+\theta}} \\ + \chi_{\{\beta \geq 1\}} [H_{(\alpha+1/2+1-\beta)(1+\theta)+1}(f^{1+\theta})(2x)]^{\frac{1}{1+\theta}} \\ + N_{2\alpha+2}f(x) + [T_{(\alpha+1/2)(1+\theta)}(f^{1+\theta})(x)]^{\frac{1}{1+\theta}},$$

with any fixed $\theta > 0$ satisfying $\frac{1}{1+\theta} < \alpha + \beta + 1/2$.

Assuming that $1/p < \alpha + \beta + 1/2$, conditions for $L^p(x^\delta dx)$ boundedness of the operators controlling $M_*^{\alpha, \beta}$ are as follows, see Sect. 2.3:

$$\begin{aligned} f &\mapsto x^\beta [D(z^{-\beta} f)^{1+\theta}(x)]^{\frac{1}{1+\theta}} : & -\beta p - 1 < \delta < (2\alpha + \beta + 1)p - 1, \\ f &\mapsto x^\beta [R(z^{-\beta} f)^{1+\theta}(x)]^{\frac{1}{1+\theta}} : & -\beta p \leq \delta, \\ f &\mapsto \chi_{\{\beta < 1\}} [H_{(\alpha+1/2)(1+\theta)+1}(f^{1+\theta})(x)]^{\frac{1}{1+\theta}} : & \delta < (2\alpha + \beta + 1)p - 1, \\ f &\mapsto \chi_{\{\beta \geq 1\}} [H_{(\alpha+3/2-\beta)(1+\theta)+1}(f^{1+\theta})(x)]^{\frac{1}{1+\theta}} : & \delta < (2\alpha + 2)p - 1, \\ N_{2\alpha+2} : & & -1 < \delta < (2\alpha + 2)p - 1, \\ f &\mapsto [T_{(\alpha+1/2)(1+\theta)}(f^{1+\theta})(x)]^{\frac{1}{1+\theta}} : & \begin{cases} -\beta p < \delta, & \text{if } \beta < 0, \\ -1 < \delta, & \text{if } \beta > 0 \text{ and } \alpha \leq -1/2, \\ \frac{-\beta}{\alpha+\beta+1/2} < \delta, & \text{if } \beta > 0 \text{ and } \alpha > -1/2. \end{cases} \end{aligned}$$

Treatment of the last operator will be explained in a moment; the cases of the other θ -operators are simpler and left to the reader. In all these conditions we understand that given δ from the indicated range, there exists $\theta > 0$ satisfying $\frac{1}{1+\theta} < \alpha + \beta + 1/2$ such that the $L^p(x^\delta dx)$ boundedness holds.

Intersecting the above conditions, we see that $M_*^{\alpha, \beta}$ is bounded on $L^p(x^\delta dx)$ if $1/p < \alpha + \beta + 1/2$ and any of the following three sets of conditions holds:

- (a) $\beta \geq 1$ and $-1 < \delta < (2\alpha + 2)p - 1$;
- (b) $0 < \beta < 1$ and $\frac{-\beta}{\max(0, \alpha+1/2)+\beta} < \delta < (2\alpha + \beta + 1)p - 1$;
- (c) $\beta < 0$ and $-\beta p < \delta < (2\alpha + \beta + 1)p - 1$.

We now explain the analysis related to T in greater detail. Observe that, for $p > 1 + \theta$, the operator $f \mapsto [T_{(\alpha+1/2)(1+\theta)}(f^{1+\theta})(x)]^{\frac{1}{1+\theta}}$ is bounded on $L^p(x^\delta dx)$ if and only if $T_{(\alpha+1/2)(1+\theta)}$ is bounded on $L^{\frac{p}{1+\theta}}(x^\delta dx)$. If $\beta < 0$ (then automatically $\alpha > -1/2$), consider $\theta > 0$ such that $1/p < \frac{1}{1+\theta} < \alpha + \beta + 1/2$ (recall that we assume $1/p < \alpha + \beta + 1/2$). Then also $\frac{1}{1+\theta} < \alpha + 1/2$ and $T_{(\alpha+1/2)(1+\theta)}$ falls under Lemma 2.3(a): It is bounded on $L^{\frac{p}{1+\theta}}(x^\delta dx)$ when

$$\delta \geq [(\alpha + 1/2)(1 + \theta) - 1] \frac{p}{1 + \theta} = (\alpha + 1/2)p - \frac{p}{1 + \theta}.$$

Choosing θ so that $\frac{1}{1+\theta}$ is sufficiently close to $\alpha + \beta + 1/2$, we can approach with the lower bound for δ arbitrarily close to $(\alpha + 1/2)p - (\alpha + \beta + 1/2)p = -\beta p$.

Let now $\beta > 0$. If $\alpha \leq -1/2$, then we choose any $\theta > 0$ such that $1/p < \frac{1}{1+\theta} < \alpha + \beta + 1/2$ and apply Lemma 2.3(d). On the other hand, when $\alpha > -1/2$ (in fact $\alpha \in (-1/2, 1/2)$), we consider $\theta > 0$ such that $0 < \alpha + 1/2 < \frac{1}{1+\theta} < \alpha + \beta + 1/2$. By Lemma 2.3(c), the condition for the boundedness is $\delta \geq (\alpha + 1/2)(1 + \theta) - 1$. One can choose θ so that $\frac{1}{1+\theta}$ is arbitrarily close to $\alpha + \beta + 1/2$, which covers all $\delta > \frac{\alpha+1/2}{\alpha+\beta+1/2} - 1 = \frac{-\beta}{\alpha+\beta+1/2}$.

Case 3 $\alpha + \beta = 1/2$ and $-\beta \notin \mathbb{N}$. By Theorem 2.2, $M_*^{\alpha, \beta} f(x) \lesssim \Psi_*^{\alpha, \beta-\varepsilon} f(x)$ with any fixed $\varepsilon > 0$. From the analysis in Case 2, with β replaced by $\beta - \varepsilon$, we infer that $M_*^{\alpha, \beta}$ is bounded on $L^p(x^\delta dx)$ if $1/p < \alpha + \beta - \varepsilon + 1/2$ and any of the following sets of conditions holds:

- (a) $\beta - \varepsilon \geq 1$ and $-1 < \delta < (2\alpha + 2)p - 1$;
- (b) $0 < \beta - \varepsilon < 1$ and $\frac{-(\beta-\varepsilon)}{\max(0, \alpha+1/2)+\beta-\varepsilon} < \delta < (2\alpha + \beta - \varepsilon + 1)p - 1$;

(c) $\beta - \varepsilon < 0$ and $-(\beta - \varepsilon)p < \delta < (2\alpha + \beta - \varepsilon + 1)p - 1$.

Since ε can be chosen arbitrarily small, the above conditions with $\varepsilon = 0$ also imply $L^p(x^\delta dx)$ boundedness of $M_*^{\alpha, \beta}$, as can be easily verified.

Case 4 $-\beta \in \mathbb{N}$. In view of Theorem 2.2, we have the bound $M_*^{\alpha, \beta} f(x) \lesssim \Phi_{*, E}^{\alpha, \beta} f(x)$ in case $\alpha + \beta \geq 1/2$, and $M_*^{\alpha, \beta} f(x) \lesssim \Psi_{*, E}^{\alpha, \beta} f(x)$ when $\alpha + \beta < 1/2$. Therefore, Lemmas 3.1 and 3.2 provide the control

$$M_*^{\alpha, \beta} f(x) \lesssim \begin{cases} Lf(x) + H_{2\alpha+\beta+1} f(4x) + x^\beta R(z^{-\beta} f)(x), & \text{if } \alpha + \beta \geq 1/2, \\ x^\beta [D(z^{-\beta} f)^{1+\theta}(x)]^{\frac{1}{1+\theta}} + x^\beta [R(z^{-\beta} f)^{1+\theta}(x)]^{\frac{1}{1+\theta}}, & \text{if } \alpha + \beta < 1/2, \end{cases}$$

with any fixed $\theta > 0$ such that $\frac{1}{1+\theta} < \alpha + \beta + 1/2$.

Taking now into account conditions for $L^p(x^\delta dx)$ boundedness of the component operators involved, see Cases 1 and 2 above, we conclude that $M_*^{\alpha, \beta}$ is bounded on $L^p(x^\delta dx)$ if $1/p < \alpha + \beta + 1/2$ (this condition is automatically satisfied when $\alpha + \beta \geq 1/2$) and

$$-\beta p \leq \delta < (2\alpha + \beta + 1)p - 1.$$

The proof of Theorem 1.5 is complete. $\square$

3.5 Proof of Proposition 1.10

To begin with, observe that $M_{*, \text{tru}}^{\alpha, \beta}$ arises from $M_*^{\alpha, \beta}$ by restricting the kernel to the region E_1 .

When $\alpha + \beta \geq 1/2$, we have $\Phi_{*, E_1}^{\alpha, \beta} f(x) \lesssim Lf(x)$, see the proof of Lemma 3.1. Further, in case $\alpha + \beta < 1/2$, $\Psi_{*, E_1}^{\alpha, \beta} f(x) \lesssim x^\beta [L(z^{-\beta} f)^{1+\theta}(x)]^{1/(1+\theta)}$ with any fixed $\theta > 0$ such that $\frac{1}{1+\theta} < \alpha + \beta + 1/2$. This is implicitly contained in the proof of Lemma 3.2. The more precise bounds of Theorem 2.1(4),(5) lead easily to the control $M_{*, \text{tru}}^{\alpha, \beta} f(x) \lesssim Lf(x)$ in case $\alpha + \beta = 1/2$ and $-\beta \notin \mathbb{N}$ (when $t < x/2$ the logarithm occurring in Theorem 2.1(4) has no effect). All this together with Theorem 2.2 justifies (1.9).

The remaining part of Proposition 1.10 follows from the well-known fact that for any $\delta \in \mathbb{R}$, L is bounded on $L^p(x^\delta dx)$, $1 < p < \infty$, and from $L^1(x^\delta dx)$ to weak $L^1(x^\delta dx)$. $\square$

4 Counterexamples

Our aim in this section is to prove Proposition 1.6. This will be done by constructing suitable counterexamples, in the first step for certain auxiliary maximal operators.

4.1 Counterexamples for auxiliary operators

For $\alpha > -1$, $\alpha + \beta > -1/2$ and $t > 0$ consider the following integral operators, with positive kernels, acting on functions on $\mathbb{R}_+$:

$$U_{t,1}^{\alpha, \beta} f(x) = \frac{x^{-2\alpha-\beta}}{t^{2\alpha+2\beta}} \int_{|t-x|}^{t+x} \left([(x+t)^2 - z^2][z^2 - (x-t)^2] \right)^{\alpha+\beta-1/2} z^{1-\beta} f(z) dz,$$

$$U_{t,2}^{\alpha, \beta} f(x) = \frac{x^{-\alpha-1/2}}{t^{2\alpha+2\beta}} \int_{|t-x|}^{t+x} [t^2 - (x-z)^2]^{\alpha+\beta-1/2} z^{\alpha+1/2} f(z) dz,$$

$$\begin{aligned}\tilde{U}_{t,2}^{\alpha,\beta} f(x) &= \frac{x^{-\alpha-1/2}}{t^{2\alpha+2\beta}} \int_{|t-x|}^{t+x} [t^2 - (x-z)^2]^{\alpha+\beta-1/2} \log \left(\frac{8xz}{(x+z)^2 - t^2} \right) z^{\alpha+1/2} f(z) dz, \\ V_{t,1}^{\alpha,\beta} f(x) &= \frac{\chi_{\{x < t\}}}{t^{2\alpha+2\beta}} \int_0^{t-x} [t^2 - (x-z)^2]^{-\alpha-1/2} [t^2 - (x+z)^2]^{\alpha+\beta-1/2} z^{2\alpha+1} f(z) dz, \\ V_{t,2}^{\alpha,\beta} f(x) &= \frac{\chi_{\{x < t\}}}{t^{2\alpha+2\beta}} \int_0^{t-x} [t^2 - (x-z)^2]^{\beta-1} z^{2\alpha+1} f(z) dz, \\ \tilde{V}_{t,2}^{\alpha,\beta} f(x) &= \frac{\chi_{\{x < t\}}}{t^{2\alpha+2\beta}} \int_0^{t-x} [t^2 - (x-z)^2]^{\beta-1} \log \left(2 \frac{t^2 - (x-z)^2}{t^2 - (x+z)^2} \right) z^{2\alpha+1} f(z) dz.\end{aligned}$$

These operators correspond to right-hand sides of the bounds (1)–(8) in Theorem 2.1. More precisely, in view of (3.2), $U_{t,1}^{\alpha,\beta}$ matches the E part in (1) and (3), (5), (6), (7), $U_{t,2}^{\alpha,\beta}$ matches the E part in (2), (5), (8), while $\tilde{U}_{t,2}^{\alpha,\beta}$ matches the E part in (4). Similarly, $V_{t,1}^{\alpha,\beta}$ matches the F part in (3), (5), (6), (8), $V_{t,2}^{\alpha,\beta}$ matches the F part in (2), (5), (7), while $\tilde{V}_{t,2}^{\alpha,\beta}$ matches the F part in (4). Observe that

$$U_{t,2}^{\alpha,\beta} f(x) \lesssim \tilde{U}_{t,2}^{\alpha,\beta} f(x), \quad V_{t,2}^{\alpha,\beta} f(x) \lesssim \tilde{V}_{t,2}^{\alpha,\beta} f(x), \quad f \geq 0, \quad x, t > 0.$$

Denote $U_{*,1}^{\alpha,\beta} f = \sup_{t>0} |U_{t,1}^{\alpha,\beta} f|$ and similarly for the remaining operators in question replacing t by $*$ in the subscript indicates the corresponding maximal operator.

Lemma 4.1 *Let $\alpha > -1$, $\alpha + \beta > -1/2$, $\delta \in \mathbb{R}$ and $1 < p < \infty$.*

(a) *The following are necessary conditions for each of $U_{*,1}^{\alpha,\beta}$, $U_{*,2}^{\alpha,\beta}$, $\tilde{U}_{*,2}^{\alpha,\beta}$ to be well defined and bounded on $L^p(\mathbb{R}_+, x^\delta dx)$:*

- (a1) $\delta < (2\alpha + \beta + 1)p - 1$,
- (a2) $\frac{1}{p} < \alpha + \beta + 1/2$,
- (a3) $-\beta p \leq \delta$.

(b) *The following are necessary conditions for each of $V_{*,1}^{\alpha,\beta}$, $V_{*,2}^{\alpha,\beta}$, $\tilde{V}_{*,2}^{\alpha,\beta}$ to be well defined and bounded on $L^p(\mathbb{R}_+, x^\delta dx)$:*

- (b1) $\delta < (2\alpha + 2)p - 1$,
- (b2) $-1 < \delta$.

In the proof of Lemma 4.1 we will need a simple technical result.

Lemma 4.2 *Let $\gamma > -1$ and $0 \leq A < B$. Then*

$$\int_A^B \left([B^2 - z^2][z^2 - A^2] \right)^\gamma z dz = \frac{\Gamma(\gamma+1)^2}{2\Gamma(2\gamma+2)} (B^2 - A^2)^{2\gamma+1}.$$

Proof Changing the variable $(z^2 - A^2)/(B^2 - A^2) = s$, we immediately see that the integral in question is equal to

$$\frac{1}{2} (B^2 - A^2)^{2\gamma+1} \int_0^1 s^\gamma (1-s)^\gamma ds.$$

The last integral is the well-known Euler beta integral. The conclusion follows. $\square$

Proof of Lemma 4.1 We will construct suitable counterexamples.

Part (a) Take $f(z) = \chi_{(1,2)}(z)$. Then, for large x we have the bounds

$$U_{x-1,1}^{\alpha,\beta} f(x) \gtrsim \frac{x^{-2\alpha-\beta}}{x^{2\alpha+2\beta}} \int_1^2 x^{2\alpha+2\beta-1} (z-1)^{\alpha+\beta-1/2} dz \simeq \frac{1}{x^{2\alpha+\beta+1}},$$

$$U_{x-1,2}^{\alpha,\beta} f(x) \gtrsim \frac{x^{-\alpha-1/2}}{x^{2\alpha+2\beta}} \int_1^2 x^{\alpha+\beta-1/2} (z-1)^{\alpha+\beta-1/2} dz \simeq \frac{1}{x^{2\alpha+\beta+1}}.$$

Thus, a necessary condition for each of $U_{*,1}^{\alpha,\beta}$, $U_{*,2}^{\alpha,\beta}$ and $\tilde{U}_{*,2}^{\alpha,\beta}$ to map $L^p(x^\delta dx)$ into itself is that the function $x \mapsto x^{-2\alpha-\beta-1}$ is in $L^p(x^\delta dx)$ for large x , which is equivalent to (a1).

Next, choose $f(z) = \chi_{(1,2)}(z)(z-1)^{-\alpha-\beta-1/2}/\log \frac{2}{z-1}$. This function belongs to $L^p(x^\delta dx)$ when $\frac{1}{p} \geq \alpha + \beta + 1/2$. However, estimating similarly as above, for large x we get

$$U_{x-1,1}^{\alpha,\beta} f(x) \gtrsim \frac{1}{x^{2\alpha+\beta+1}} \int_1^2 \frac{1}{(z-1) \log(\frac{2}{z-1})} dz = \infty,$$

and in the same way $U_{x-1,2}^{\alpha,\beta} f(x) = \infty$. This shows that (a2) is necessary for each of $U_{*,1}^{\alpha,\beta}$, $U_{*,2}^{\alpha,\beta}$ and $\tilde{U}_{*,2}^{\alpha,\beta}$ to be well defined on $L^p(x^\delta dx)$.

Now, let $f_N(z) = \chi_{(N-1,N+1)}(z)z^\beta$ with N large. Then, for $x \in (1, 2)$ we have the estimates

$$U_{N,1}^{\alpha,\beta} f_N(x) \simeq \frac{1}{N^{2\alpha+2\beta}} \int_{N-1}^{N+1} \left([(x+N)^2 - z^2][z^2 - (x-N)^2] \right)^{\alpha+\beta-1/2} z dz \simeq 1,$$

where the last relation follows from Lemma 4.2. Furthermore, still for $x \in (1, 2)$,

$$U_{N,2}^{\alpha,\beta} f_N(x) \simeq \frac{1}{N^{2\alpha+2\beta}} \int_{N-1}^{N+1} [(N-x+z)(N+x-z)]^{\alpha+\beta-1/2} N^{\alpha+1/2} N^\beta dz$$

$$\simeq \int_{N-1}^{N+1} (N+x-z)^{\alpha+\beta-1/2} dz \simeq 1.$$

All these bounds hold uniformly in x and N . On the other hand, when $\delta < -\beta p$ it is straightforward to check that the norms of f_N in $L^p(x^\delta dx)$ tend to zero as $N \rightarrow \infty$. This gives the necessity of (a3).

Part (b) Observe that for $x > 0$ and $f \geq 0$, we have the bounds

$$V_{2x,1}^{\alpha,\beta} f(x) \simeq \frac{1}{x^{2\alpha+2\beta}} \int_0^x [4x^2 - (x-z)^2]^{-\alpha-1/2} [4x^2 - (x+z)^2]^{\alpha+\beta-1/2} z^{2\alpha+1} f(z) dz$$

$$\gtrsim \frac{1}{x^{2\alpha+2\beta}} x^{-2\alpha-1} \int_0^{x/2} [4x^2 - (x+z)^2]^{\alpha+\beta-1/2} z^{2\alpha+1} f(z) dz$$

$$\simeq \frac{1}{x^{2\alpha+2}} \int_0^{x/2} z^{2\alpha+1} f(z) dz,$$

$$V_{2x,2}^{\alpha,\beta} f(x) \gtrsim \frac{1}{x^{2\alpha+2\beta}} \int_0^{x/2} [4x^2 - (x-z)^2]^{\beta-1} z^{2\alpha+1} f(z) dz$$

$$\simeq \frac{1}{x^{2\alpha+2}} \int_0^{x/2} z^{2\alpha+1} f(z) dz.$$

Take now $f(z) = \chi_{(0,1)}(z)z^{-(\delta+1)/p}/\log \frac{2}{z}$. As easily verified, this function belongs to $L^p(x^\delta dx)$. Hence, in view of the above estimates, a necessary condition for each of $V_{*,1}^{\alpha,\beta}$,

$V_{*,2}^{\alpha,\beta}$ and $\tilde{V}_{*,2}^{\alpha,\beta}$ to be well defined on $L^p(x^\delta dx)$ is that the function $z \mapsto z^{2\alpha+1} z^{-(\delta+1)/p} / \log \frac{2}{z}$ is integrable at 0^+ , which is equivalent to (b1).

Finally, consider $f(z) = \chi_{(0,1)}(z)z^{-\delta}$. Assume that $\delta \leq -1$. Then, $f \in L^p(x^\delta dx)$. Further, choosing $t = 2$, for $x \in (0, 1)$ we have

$$\begin{aligned} V_{2,1}^{\alpha,\beta} f(x) &\simeq \int_0^{2-x} [4 - (x-z)^2]^{-\alpha-1/2} [4 - (x+z)^2]^{\alpha+\beta-1/2} z^{2\alpha+1-\delta} \chi_{(0,1)}(z) dz \\ &\gtrsim \int_0^{1/2} [4 - (x-z)^2]^{-\alpha-1/2} [4 - (x+z)^2]^{\alpha+\beta-1/2} z^{2\alpha+1-\delta} dz \\ &\simeq \int_0^{1/2} z^{2\alpha+1-\delta} dz \simeq 1, \\ V_{2,2}^{\alpha,\beta} f(x) &\simeq \int_0^{2-x} [4 - (x-z)^2]^{\beta-1} z^{2\alpha+1-\delta} \chi_{(0,1)}(z) dz \simeq \int_0^1 z^{2\alpha+1-\delta} dz \simeq 1. \end{aligned}$$

This implies that none of $V_{*,1}^{\alpha,\beta}$, $V_{*,2}^{\alpha,\beta}$ and $\tilde{V}_{*,2}^{\alpha,\beta}$ maps $L^p(x^\delta dx)$ into itself. The necessity of (b2) follows.

The proof of Lemma 4.1 is complete. $\square$

4.2 Proof of Proposition 1.6

Let $\alpha > -1$ and $\alpha + \beta > -1/2$. For $\varepsilon > 0$, denote

$$\begin{aligned} E_\varepsilon &= \{(t, x, z) \in E : t - |x - z| < \varepsilon\sqrt{xz} \text{ or } x + z - t < \varepsilon\sqrt{xz}\}, \\ F_\varepsilon &= \{(t, x, z) \in F : t - (x + z) < \varepsilon\sqrt{xz} \text{ or } t > \varepsilon^{-1}\sqrt{xz}\}. \end{aligned}$$

From the results of [2], see [2, Theorem 3.3] and the accompanying comments, it follows that for each pair α, β there exists $\varepsilon = \varepsilon(\alpha, \beta) > 0$ such that the kernel $K_t^{\alpha,\beta}(x, z)$ does not change sign in E_ε and in F_ε and, moreover, $|K_t^{\alpha,\beta}(x, z)|$ is comparable in E_ε and in F_ε with the corresponding expressions on the right-hand side of the relevant bound from (1)–(8) in Theorem 2.1.

Therefore, the unboundedness results obtained for $U_{*,1}^{\alpha,\beta}$, $U_{*,2}^{\alpha,\beta}$, $\tilde{U}_{*,2}^{\alpha,\beta}$, $V_{*,1}^{\alpha,\beta}$, $V_{*,2}^{\alpha,\beta}$, $\tilde{V}_{*,2}^{\alpha,\beta}$ in Lemma 4.1 will also be valid for $M_*^{\alpha,\beta}$ (with α, β suitably restricted according to the splitting in (1)–(8) in Theorem 2.1) provided that we assure that (t, x, z) involved in the counterexamples are located either in E_ε or in F_ε .

In the counterexamples from part (a) of the proof of Lemma 4.1, one can consider either x large enough (in the first and the second counterexamples) or N large enough (in the third counterexample) so that all (t, x, z) involved are contained in E_ε . In part (b), in the first counterexample one considers x sufficiently large, while in the second one x in a sufficiently small neighborhood of 0, so that all (t, x, z) involved belong to F_ε .

We conclude that necessary conditions for $M_*^{\alpha,\beta}$ to be well defined and bounded on $L^p(x^\delta dx)$ are (a1), (a2), (a3) of Lemma 4.1, and in case $-\beta \notin \mathbb{N}$ also (b1) and (b2). However, since when $-\beta \in \mathbb{N}$ conditions (b1) and (b2) are less restrictive than (a1) and (a3), respectively, actually all (a1), (a2), (a3), (b1), (b2) are always the necessary conditions. $\square$

Remark 4.3 The discrepancy between the sufficient conditions in Theorem 1.5 and the necessary conditions from Proposition 1.6 is related to region F_2'' and the special maximal operator T_η . We do not know whether the restriction of $M_*^{\alpha,\beta}$ to F_2'' is always controlled by means of

T_η in an optimal way. Neither we know whether the description of mapping properties of T_η from Lemma 2.3 is sharp. These issues remain to be investigated.

Acknowledgements We thank the referee for a constructive criticism that improved the presentation and led us to enrich the paper with the sharpness results.

References

1. Bourgain, J.: Averages in the plane over convex curves and maximal operators. *J. Anal. Math.* **47**, 69–85 (1986)
2. Ciaurri, Ó., Nowak, A., Roncal, L.: Two-weight mixed norm estimates for a generalized spherical mean Radon transform acting on radial functions. *SIAM J. Math. Anal.* **49**, 4402–4439 (2017)
3. Colzani, L., Cominardi, A., Stempak, K.: Radial solutions to the wave equation. *Ann. Mat. Pura Appl.* **181**, 25–54 (2002)
4. Duoandikoetxea, J., Moyua, A., Oruetebarria, O.: The spherical maximal operator on radial functions. *J. Math. Anal. Appl.* **387**, 655–666 (2012)
5. Duoandikoetxea, J., Moyua, A., Oruetebarria, O.: Estimates for radial solutions to the wave equation. *Proc. Am. Math. Soc.* **144**, 1543–1552 (2016)
6. Miao, C., Yang, J., Zheng, J.: On local smoothing problems and Stein’s maximal spherical means. *Proc. Am. Math. Soc.* **145**, 4269–4282 (2017)
7. Mockenhaupt, G., Seeger, A., Sogge, C.: Wave front sets, local smoothing and Bourgain’s circular maximal theorem. *Ann. Math.* **136**, 207–218 (1992)
8. Stein, E.M.: Maximal functions. I. Spherical means. *Proc. Natl. Acad. Sci. USA* **73**, 2174–2175 (1976)

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.