



Blow-up versus global existence of solutions for reaction–diffusion equations on classes of Riemannian manifolds

Gabriele Grillo¹ · Giulia Meglioli¹ · Fabio Punzo¹

Received: 31 May 2022 / Accepted: 7 October 2022 / Published online: 7 December 2022
© The Author(s) 2022

Abstract

It is well known from the work of Bandle et al. (J Differ Equ 251:2143–2163, 2011) that the Fujita phenomenon for reaction–diffusion evolution equations with power nonlinearities does not occur on the hyperbolic space \mathbb{H}^N , thus marking a striking difference with the Euclidean situation. We show that, on classes of manifolds in which the bottom Λ of the L^2 spectrum of $-\Delta$ is strictly positive (the hyperbolic space being thus included), a different version of the Fujita phenomenon occurs for other kinds of nonlinearities, in which the role of the critical Fujita exponent in the Euclidean case is taken by Λ . Such nonlinearities are time-independent, in contrast to the ones studied in Bandle et al. (2011). As a consequence of our results we show that, on a class of manifolds much larger than the case $M = \mathbb{H}^N$ considered in Bandle et al. (2011), solutions to (1.1) with power nonlinearity $f(u) = u^p$, $p > 1$, and corresponding to sufficiently small data, are global in time. Though qualitative similarities with similar problems in bounded, Euclidean domains can be seen in the results, the methods are significantly different because of noncompact setting dealt with.

Keywords Reaction diffusion equations · Riemannian manifolds · Blow-up · Global existence · Fujita type exponent

Mathematics Subject Classification Primary 35K57 · Secondary 35B44 · 58J35 · 35K65 · 35R01

✉ Giulia Meglioli
giulia.meglioli@polimi.it
Gabriele Grillo
gabriele.grillo@polimi.it
Fabio Punzo
fabio.punzo@polimi.it

¹ Dipartimento di Matematica, Politecnico di Milano, Piazza Leonardo da Vinci 32, 20133 Milan, Italy

1 Introduction

We investigate existence of global in time solutions, versus blow-up in finite time, to nonlinear reaction–diffusion problems of the following type:

$$\begin{cases} u_t = \Delta u + f(u) & \text{in } M \times (0, T) \\ u = u_0 \geq 0 & \text{in } M \times \{0\}, \end{cases} \tag{1.1}$$

where Δ is the Laplace–Beltrami operator on a Riemannian manifold M , $T \in (0, \infty]$ and $f : [0, +\infty) \rightarrow [0, +\infty)$ is e.g., a locally Lipschitz, increasing function. Further specification of the assumptions on the manifold M and on the nonlinearity f will be made later on. A crucial parameter, according to whose value the behaviour of solutions will change, will be the value of $f'(0)$ when f is differentiable in $x = 0$. It will be usually required that $u_0 \in C(M) \cap L^\infty(M)$ to ensure the existence of classical solutions at least up to a certain time.

The analogue of (1.1) in the Euclidean setting has a long history especially in the particularly important case of power nonlinearities, i.e., for the problem

$$\begin{cases} u_t = \Delta u + |u|^{p-1}u & \text{in } \mathbb{R}^n \times (0, T) \\ u = u_0 & \text{in } \mathbb{R}^n \times \{0\}, \end{cases} \tag{1.2}$$

where it is assumed that $u_0 \in L^\infty(\mathbb{R}^n)$. It has been shown by Fujita in [5], and in [12] and [14] for the critical case, that for problem (1.2) the following facts hold:

- a) If $1 < p \leq p^* := \frac{N+2}{N}$, (1.2) does not possess nontrivial global solutions.
- b) If $p > p^*$ solutions corresponding to data that are sufficiently small in a suitable sense are global in time.

It should be noticed that, by a generalization of a result of Kaplan [13], solutions corresponding to *sufficiently large* data blow up for any $p > 1$. A complete account of results concerning blow-up and global existence of solutions to semilinear parabolic equations posed in \mathbb{R}^n can be found, e.g., in [1, 4, 16, 23] and in references therein.

In the case of evolution equation posed on Riemannian manifolds the situation may change completely. In fact, an analogue of (1.2) has been studied in [2] in the important context of the *hyperbolic space* \mathbb{H}^n , namely on the simply connected manifold of constant sectional curvature equal to -1 . It is shown there that for *all* $p > 1$ sufficiently small initial data give rise to global in time solutions. A kind of Fujita phenomenon nonetheless takes place but when a *time dependent* modification of (1.2) is taken into account. Namely, if the reaction term u^p (considering nonnegative data, hence nonnegative solutions) is replaced by $e^{\alpha t}u^p$, $\alpha > 0$ being a fixed parameter, then a Fujita-type phenomenon then takes place, the threshold value being $p^\sharp := 1 + \frac{\alpha}{\Lambda}$ where $\Lambda := (N - 1)^2/4$ is the bottom of the L^2 spectrum of $-\Delta$ on \mathbb{H}^n . Informally, one might say that for this to hold the nonlinearity must be amplified exponentially as time grows, in fact it is also shown in [2] that if the exponential factor in time is replaced by a power of time the Fujita phenomenon still does not occur. See also [24] for a careful analysis of the critical case $p = p^\sharp$, in which the authors show, combining their results with the ones of [2], that in such case global in time solutions exist for all values of α , thus marking a further difference with the Euclidean case. Further precise results on the lifespan of solutions that are not global are given in [25].

Recently, a number of results concerning blow-up and global existence for solutions of nonlinear parabolic equations with power-like reaction term and *nonlinear, slow diffusion of porous medium type* has also been obtained, on some classes of Riemannian manifolds, in [9–11, 17, 26]. The results for this class of equations show usually several differences with the case involving a linear diffusion, already in the Euclidean case, see [22].

Our goal here will be to present some result for different type of *time independent* nonlinearities in which a new kind of Fujita phenomenon takes place, in a wide class of manifolds that includes the hyperbolic space. We shall consider noncompact complete Riemannian manifold M of infinite volume, with dimension $N \geq 3$, and require that some further conditions on M , to be described below, hold. In our first result, Theorem 3.1, we require the following additional condition on M :

- M is stochastically complete, and $\lambda_1(M) := \inf \text{spec}(-\Delta) > 0$, where $\text{spec}(-\Delta)$ is the L^2 spectrum of the Laplace–Beltrami operator $-\Delta$.

Under such assumptions, we prove *nonexistence of global solutions for problem (1.1) e.g., if f is convex and increasing, $f'(0) > \lambda_1(M)$, and $1/f$ is integrable at infinity.*

Stochastic completeness is a well-studied property of Riemannian manifolds, and it amounts to requiring that $T_t 1 = 1$ for all $t > 0$, or equivalently that

$$\int_M p(x, y, t) \, d\mu(y) = 1, \quad \text{for all } x \in M, t > 0. \tag{1.3}$$

where $p(x, y, t)$ is the heat kernel of the manifold M and μ the Riemannian measure. See e.g., [6, 7] for a number of conditions on M ensuring that stochastic completeness holds. For example, it suffices that, for some $o \in M$, the function $r \mapsto \frac{r}{\log \mathcal{V}(o, r)}$ is not integrable at infinity, where $\mathcal{V}(o, r)$ is the volume of the geodesic ball of radius r centered at o . Note that this is true in particular if $\mathcal{V}(o, r) \leq Ce^{ar^2}$ for suitable $C, a > 0$. This allows e.g., sectional curvatures to tend to $-\infty$ at the infinity of M (namely when $\rho(o, x) \rightarrow +\infty$, ρ being the Riemannian distance and o being fixed in M), at most quadratically, in a suitable precise sense, see also [8] for relations to nonlinear elliptic and parabolic equations on M . As for the assumption $\lambda_1(M) > 0$ we comment that a well-known sufficient condition for this to hold is that $\text{sec} \leq c < 0$, sec denoting sectional curvatures. Thus, the class of manifolds on which the above result works is large, as it includes e.g., all those manifolds whose sectional curvatures are pinched between two strictly negative constants, and in particular the hyperbolic space.

In our second result, Theorem 3.2, the additional assumption we require on M beside the previous ones is the following:

- the following Faber–Krahn inequality holds: for some $c > 0$, for any non-empty relatively compact open subset $\Omega \subset M$,

$$\lambda_1(\Omega) \geq \frac{c}{[\mu(\Omega)]^{\frac{2}{N}}}, \tag{1.4}$$

where $\lambda_1(\Omega)$ is the first eigenvalue of the Laplace operator on Ω completed with homogeneous Dirichlet boundary conditions, and μ is the Riemannian measure.

Under such assumptions, we show *existence of global solutions for small data e.g., if f is differentiable at $x = 0$ with $f'(0) < \lambda_1(M)$.*

We refer e.g., to [7, Cor. 14.23, Cor. 15.17] for equivalent conditions for the Faber–Krahn inequality to hold. In particular, its validity is implied by on-diagonal bounds for the heat kernel in the form $p(t, x, x) \leq c t^{-N/2}$ for all $x \in M, t > 0$, or by the validity of the Sobolev (or the Nash) inequality on M , which is in turn satisfied e.g., when $\sec \leq 0$ on M , \sec denoting sectional curvatures.

Observe that our blow-up result is similar in character to the well-known blow-up result for bounded domains of \mathbb{R}^n (see, e.g., [1, Section 3.2]). However, the methods of proof exploited for bounded domains do not work on general Riemannian manifolds, thus our arguments are completely different. Indeed, in bounded domains the blow-up result is usually obtained by means of the Kaplan method (see [1, 13]), which makes use of the first eigenvalue and of the first eigenfunction of the Laplacian. In order to extend that argument to a general Riemannian manifold, it would be necessary to know precisely the behaviour at infinity of the positive solution ϕ to

$$\Delta \phi + \lambda_1(M)\phi = 0 \quad \text{in } M$$

and in particular its integrability properties w.r.t. the Riemannian measure, which are not known in general, e.g., on the hyperbolic space \mathbb{H}^n ϕ belongs just to $L^{2+\varepsilon}$ for all $\varepsilon > 0$. The Kaplan method yields indeed partial results on the subclass of Cartan–Hadamard manifolds, i.e., simply connected Riemannian manifolds with nonpositive sectional curvatures, with Ricci curvature bounded from below, see [20] when $f(u) = u^p$, but the method can not be pushed to get the sharp threshold value $\lambda_1(M)$ provided in Theorem 3.1, 3.2, even in the special case of Cartan–Hadamard manifolds and, in fact, even on the special case $M = \mathbb{H}^n$.

Let us mention that some blow-up results in bounded domains have also been established in [18, 19], for more general operators, by means of the method of sub- and super-solutions. Those results seem to be quite implicit in character, and they are based on the asymptotic behaviour for large times of solutions to the associated linear problem.

As a further comment, we mention that it is easy to show that, on a wide class of manifolds, characterized by the validity of the parabolic maximum principle, a sufficient condition for which is e.g., the very general condition (6.2) below, *blow-up of solutions corresponding to large data occurs*, provided the nonlinearity f is convex and $1/f$ is integrable at infinity. See the end of this section for some more detail and Sect. 6 for a concise proof.

Let us now give briefly some more precise detail on the conditions on the nonlinearities to be verified in order to prove our results. In the first one, Theorem 3.1, we shall assume that f is continuous. Besides, we assume that $f \geq h$ where h is increasing and convex in $[0, +\infty)$, that it satisfies $\int_0^{+\infty} \frac{1}{h(s)} ds < +\infty$, that $h(0) = 0$ and finally that the condition $h'(0) > \lambda_1(M)$ holds. Of course, a sufficient condition for this to hold is that the conditions satisfied by h are satisfied by the nonlinearity f itself, as mentioned above. Then we show that *all* solutions blow up in finite time. In Theorem 3.2 we shall show that, for any locally Lipschitz nonlinearity f such that $f(x) \leq \lambda x$ in a neighbourhood of $x = 0$, with $\lambda \leq \lambda_1(M)$, then *sufficiently small* data give rise to solutions existing for all times. Of course, if $f'(0)$ exists, a sufficient condition for the above condition on f to hold for some $\lambda < \lambda_1(M)$ is $f'(0) < \lambda_1(M)$, as mentioned above. The combination of the two results thus shows the version of the Fujita phenomenon we aim at.

It is important to mention that the above results provide, as immediate consequences, new results w.r.t. the ones proved in [2] for (1.1) even in the classical case $f(u) = u^p$. In fact, Theorem 3.2 shows in particular that *solutions corresponding to sufficiently small data are global*, on a much wider class of manifolds than the hyperbolic space considered in one of the main result of [2], see Corollary 3.3 for a precise statement. In particular the

results holds e.g., on all those manifolds whose sectional curvatures are pinched between two strictly negative constants everywhere.

The paper is organized as follows. In Sect. 2 we collect some preliminary material on the class of manifolds considered and on the concept of solution. In Sect. 3 we state all our main results. Section 4 contains the proof of Theorem 3.1, whereas Theorem 3.2 is proved in Sect. 5. In Sect. 6 we complement our main results by considering a large class of manifolds, e.g., those ones in which the radial Ricci curvature does not diverge at infinity faster than $-cr^2$, where r is the Riemannian distance from a given pole $o \in M$. We give a concise proof of the fact that on such manifolds, if f is a locally Lipschitz, increasing, convex function such that $1/f$ is integrable at infinity, large data give rise to solutions blowing up in finite time.

2 Preliminaries

2.1 Heat semigroup on M

Let $\{e^{t\Delta}\}_{t \geq 0}$ be the heat semigroup of M , acting on $L^p(M)$ for all $p \in [1, +\infty]$. It admits a (minimal) *heat kernel*, namely a function $p \in C^\infty(M \times M \times (0, +\infty))$, $p > 0$ in $M \times M \times (0, +\infty)$ such that

$$(e^{t\Delta}u_0)(x) = \int_M p(x, y, t) u_0(y) \, d\mu(y), \quad x \in M, \, t > 0,$$

for any $u_0 \in L^p(M)$. It is well known that

$$\int_M p(x, y, t) \, d\mu(y) \leq 1, \quad \text{for all } x \in M, \, t > 0. \tag{2.1}$$

As recalled in the Introduction, we say that a manifold M is stochastically complete if the following condition holds:

$$\int_M p(x, y, t) \, d\mu(y) = 1, \quad \text{for all } x \in M, \, t > 0.$$

See the considerations and the references after (1.3) for sufficient conditions for this fact to hold.

Furthermore, it is known that if M is a noncompact Riemannian manifold, then (see [3, Corollary 1])

$$\lim_{t \rightarrow +\infty} \frac{\log p(x, y, t)}{t} = -\lambda_1(M) \quad \text{locally uniformly in } M \times M, \tag{2.2}$$

where $\lambda_1(M)$ is the infimum of the L^2 spectrum of $-\Delta$. We also recall that from the Faber–Krahn inequality (1.4) it follows that (see [7, Cor. 15.17 (b)]), for some $\bar{C} > 0$:

$$p(x, y, t) \leq \bar{C} e^{-\lambda_1 t} \quad \text{for any } x, y \in M, \, t \geq 1. \tag{2.3}$$

2.2 On the concept of solution

We shall always deal with bounded initial data. Solutions will be meant in the classical sense. More precisely, setting $Q_T = M \times (0, T]$, we require that $u \in C^{2,1}(Q_T) \cap C(\overline{Q_T}) \cap L^\infty(\overline{Q_T})$ and that (1.1) holds in the classical sense.

We shall use in the sequel two different concepts of solution. On the one hand a function $u \in C(M \times (0, \tau]) \cap L^\infty(M \times (0, \tau])$, for every $\tau \in (0, T]$ is called a *mild solution* of problem (1.1) if

$$u(x, t) = (e^{t\Delta}u_0)(x) + \int_0^t (e^{(t-s)\Delta}f(u))(x) ds, \tag{2.4}$$

for any $t \in [0, \tau]$.

We notice that, by adapting the methods of [2, Prop. 2.1, Lemma 2.1], for *bounded* initial data u_0 and up to a time T such that $u(t)$ is bounded for all $t \in [0, T)$ (blow-up might occur at some positive time), the two concepts of solutions coincide provided f is locally Lipschitz, as required in our main results. Hence we shall use them indifferently when needed.

3 Statements of main results

In this section, we state our results concerning solutions to problem (1.1). We say that a solution *blows up in finite time*, whenever there exists $\tau > 0$ such that

$$\lim_{t \rightarrow \tau^-} \|u(t)\|_{L^\infty(M)} = +\infty.$$

Otherwise, if a solution $u(t) \in L^\infty(M)$ for all $t > 0$, we say that it is *global*. Our first results involves nonexistence of global solutions. Notice that the assumptions on the auxiliary function h below entail that h is differentiable in $x = 0$.

Theorem 3.1 *Let M be a complete, non compact, stochastically complete Riemannian manifold with $\lambda_1(M) > 0$. Let $u_0 \in C(M) \cap L^\infty(M)$, $u_0 \geq 0$, $u_0 \not\equiv 0$ in M . Let f be locally Lipschitz in $[0, +\infty)$. Assume that $f \geq h$ where h is increasing and convex in $[0, +\infty)$ and $h(0) = 0$. Moreover, suppose that*

$$\int_0^{+\infty} \frac{1}{h(s)} ds < +\infty, \tag{3.1}$$

and finally that $h'(0) > \lambda_1(M)$. Then any solution to problem (1.1) blows up in finite time.

Notice that in the above Theorem the fact that h is assumed to be increasing and convex implies the existence of $h'(0)$.

Theorem 3.2 *Let M be a complete, non compact, stochastically complete Riemannian manifold with $\lambda_1(M) > 0$ and such that the Faber–Krahn inequality (1.4) holds. Assume also that f is increasing, locally Lipschitz and $f(0) = 0$. Moreover, suppose that, for some $\delta > 0$ and $0 < \alpha \leq \lambda_1(M)$,*

$$f(x) \leq \alpha x \quad \text{for all } x \in [0, \delta]. \tag{3.2}$$

Furthermore, assume that $u_0 \in C(M) \cap L^\infty(M) \cap L^1(M)$, $u_0 \geq 0$ in M is small enough. Then there exists a global solution to problem (1.1); in addition, $u \in L^\infty(M \times (0, +\infty))$.

The smallness condition on u_0 in Theorem 3.2 can be precisely formulated. Indeed, our hypothesis is that

$$\|u_0\|_{L^\infty(M)} \leq \delta e^{-\alpha}, \tag{3.3}$$

and

$$\|u_0\|_{L^1(M)} \leq \frac{\delta}{C_2}, \tag{3.4}$$

where $C_2 = \bar{C} e^{-(\lambda_1 - \alpha)}$, \bar{C} being defined in (2.3), while α and δ are given by (3.2).

As a consequence, we can generalize one of the main results of [2] (see also [21]) to a class of manifolds much wider than \mathbb{H}^n .

Corollary 3.3 *Let M be a complete, non compact, stochastically complete Riemannian manifold with $\lambda_1(M) > 0$ and such that the Faber–Krahn inequality (1.4) holds. Assume $f(x) = x^p$ for all $x \geq 0$ with $p > 1$. Assume also that $u_0 \in C(M) \cap L^\infty(M) \cap L^1(M)$, $u_0 \geq 0$ in M is small enough. Then there exists a global solution to problem (1.1); in addition, $u \in L^\infty(M \times (0, +\infty))$.*

Remark 3.4

- For any $p > 1$, let

$$f(u) = \begin{cases} \alpha u, & u \in [0, 1], \\ \alpha u^p & u \in (1, +\infty). \end{cases}$$

If $\alpha > \lambda_1(M)$, then by Theorem 3.1, the solution to problem (1.1) blows up in finite time for any nontrivial u_0 . On the other hand, if $\alpha \leq \lambda_1(M)$, then the solution exists globally in time, provided that u_0 is sufficiently small.

- Let $f(u) = e^{\beta u} - 1$ with $\beta > 0$. By Theorem 3.1, if $\beta > \lambda_1(M)$, then the solution to problem (1.1) blows up in finite time. On the contrary, if $\beta < \lambda_1(M)$, then condition (3.2) is satisfied with $\beta < \alpha \leq \lambda_1(M)$. Therefore, by Theorem 3.2, the solution exists globally in time, whenever u_0 is sufficiently small.

By standard methods, on suitable manifolds and for a wide class of nonlinearities f , it is possible to show that whenever u_0 is large enough, blow-up of solutions occurs. We defer the discussion of this fact to Sect. 6.

4 Finite time blow-up for any initial datum

4.1 Two key estimates

Let us first prove a preliminary lemma.

Lemma 4.1 *Let M be a complete, non-compact Riemannian manifold with $\lambda_1(M) > 0$. Let $u_0 \in C(M) \cap L^\infty(M)$, $u_0 \geq 0$, $u_0 \not\equiv 0$ in M . Let $\varepsilon \in (0, \lambda_1(M))$. Then there exist $\Omega \subset M$, $t_0 > 0$, $C_1 > 0$ such that*

$$(e^{t\Delta}u_0)(x) \geq C_1 e^{-[\lambda_1(M)+\varepsilon]t}, \quad \text{for any } x \in \Omega, \quad t > t_0. \tag{4.1}$$

Proof Let $\Omega \subset M$ be such that $\mu(\Omega) < +\infty$, $\int_\Omega u_0 \, d\mu > 0$. From (2.2), there exists $t_0 > 0$ such that, for every $x, y \in \Omega$,

$$p(x, y, t) \geq e^{-[\lambda_1(M)+\varepsilon]t} \quad \text{for every } t > t_0.$$

Hence

$$\begin{aligned} (e^{t\Delta}u_0)(x) &\geq \int_M p(x, y, t)u_0(y) \, d\mu(y) \\ &\geq e^{-[\lambda_1(M)+\varepsilon]t} \int_\Omega u_0(y) \, d\mu(y). \end{aligned}$$

Consequently, we obtain (4.1) with $C_1 := \int_\Omega u_0(y) \, d\mu(y) > 0$. □

Let u be a mild solution of equation (1.1), so that it fulfills (2.4). Then, for any $x \in M$ and for any $T > 0$, we define

$$\Phi_x^T(t) \equiv \Phi_x(t) := \int_M p(x, z, T-t)u(z, t) \, d\mu(z) \quad \text{for any } t \in [0, T]. \tag{4.2}$$

Observe that

$$\Phi_x(0) = \int_M p(x, z, T)u_0(z) \, d\mu(z) = (e^{T\Delta}u_0)(x), \quad x \in M. \tag{4.3}$$

Suppose that $u_0 \in L^\infty(M)$. Choose any

$$\delta > \|u_0\|_{L^\infty(M)}. \tag{4.4}$$

From (4.4) and (2.1) we obtain that, for any $x \in M$,

$$\begin{aligned} \Phi_x(0) &= \int_M p(x, z, T)u_0(z) \, d\mu(z) \\ &\leq \|u_0\|_{L^\infty(M)} \int_M p(x, z, T) \, d\mu(z) \\ &< \delta. \end{aligned} \tag{4.5}$$

We now state the following lemma.

Lemma 4.2 *Let M, f, h, u_0 be as in Theorem 3.1. Let $x \in M$ and $\Phi_x(t)$ be as in (4.2). Set $\alpha := h'(0)$. Then*

$$\Phi_x(0) \leq C e^{-\alpha T}, \quad \text{for any } T \geq \bar{t}, \tag{4.6}$$

for suitable $\bar{t} > 0$ and $C > 0$, depending on x .

Note that \bar{t} and C are given by (4.18) and (4.27) below, respectively, with δ as in (4.4).

Proof Let u be a solution to problem (1.1). So (2.4) holds; hence

$$u(z, t) = \int_M p(z, y, t)u_0(y) \, d\mu(y) + \int_0^t \int_M p(z, y, t - s)f(u) \, d\mu(y) \, ds. \tag{4.7}$$

In the definition of $\Phi_x^T(t) \equiv \Phi_x(t)$ (see (4.2)) fix any

$$T > \frac{1}{\alpha} [\log \delta - \log \Phi_x(0)]. \tag{4.8}$$

We multiply (4.7) by $p(x, z, T - t)$ and integrate over M . Therefore, we get

$$\begin{aligned} \int_M p(x, z, T - t) u(z, t) \, d\mu(z) &= \int_M \int_M p(z, y, t) u_0(y) p(x, z, T - t) \, d\mu(z) \, d\mu(y) \\ &\quad + \int_0^t \int_M \int_M p(z, y, t - s)f(u) p(x, z, T - t) \, d\mu(z) \, d\mu(y) \, ds. \end{aligned} \tag{4.9}$$

Now, due to (4.2), for all $t \in (0, T)$, equality (4.9) reads

$$\begin{aligned} \Phi_x(t) &= \int_M \int_M p(z, y, t) u_0(y) p(x, z, T - t) \, d\mu(z) \, d\mu(y) \\ &\quad + \int_0^t \int_M \int_M p(z, y, t - s)f(u) p(x, z, T - t) \, d\mu(z) \, d\mu(y) \, ds. \end{aligned}$$

By (4.3), for all $t \in (0, T)$,

$$\begin{aligned} \Phi_x(t) &= \int_M p(x, y, T) u_0(y) \, d\mu(y) + \int_0^t \int_M f(u) p(x, y, T - s) \, d\mu(y) \, ds \\ &= \Phi_x(0) + \int_0^t \int_M f(u) p(x, y, T - s) \, d\mu(y) \, ds. \end{aligned}$$

Since $f \geq h$ in $[0, +\infty)$,

$$\begin{aligned} \Phi'_x(t) &= \int_M f(u) p(x, y, T - t) \, d\mu(y) \\ &\geq \int_M h(u) p(x, y, T - t) \, d\mu(y). \end{aligned} \tag{4.10}$$

Since h is an increasing convex function, due to (1.3), by using Jensen inequality, we get

$$\begin{aligned} \int_M p(x, y, T - t) h(u(y, t)) \, d\mu(y) &\geq h\left(\int_M p(x, y, T - t) u(y, t) \, d\mu(y)\right) \\ &= h(\Phi_x(t)). \end{aligned} \tag{4.11}$$

Combining together (4.10) and (4.11), we obtain

$$\Phi'_x(t) \geq h(\Phi_x(t)) \quad \text{for all } t \in (0, T). \tag{4.12}$$

Fix any $x \in M$. We first observe that (4.12) implies that $\Phi_x(t)$ is an increasing function w.r.t the time variable t , since

$$\Phi'_x(t) > 0 \quad \text{for any } t \in (0, T). \tag{4.13}$$

Moreover, due to (4.4) and (4.5), by continuity of $t \mapsto \Phi_x(t)$, we can infer that there exists $t_1 > 0$ such that

$$\Phi_x(t) < \delta \quad \text{for all } t \in (0, t_1). \tag{4.14}$$

Since h is convex, increasing in $[0, +\infty)$, $h(0) = 0, h'(0) = \alpha$, then

$$h(s) \geq \alpha s \quad \text{for all } s \geq 0. \tag{4.15}$$

Due to (4.15) and to (4.14), we get

$$\begin{cases} \Phi'_x(t) \geq \alpha \Phi_x(t) & \text{for any } t \in (0, t_1), \\ \Phi_x(0) < \delta \end{cases}.$$

Let

$$\bar{t} := \sup\{t > 0 : \Phi_x(t) < \delta\}.$$

We claim that

$$0 < \bar{t} \leq -\frac{1}{\alpha} \log(\Phi_x(0)) + \frac{\log \delta}{\alpha}. \tag{4.16}$$

In order to show (4.16), consider the Cauchy problem

$$\begin{cases} y'(t) = \alpha y(t), & t > 0 \\ y(0) = \Phi_x(0). \end{cases}$$

Clearly,

$$y(t) = \Phi_x(0)e^{\alpha t}, \quad t > 0.$$

Hence

$$y(\tau) = \delta \quad \text{whenever } \tau = \frac{1}{\alpha} [\log(\delta) - \log(\Phi_x(0))].$$

Furthermore, note that, in view of (4.5) and (4.8),

$$0 < \tau < T. \tag{4.17}$$

By comparison,

$$\Phi_x(t) \geq y(t) \quad \text{for all } t \in (0, T).$$

Thus, we can infer that there exists $\bar{t} \in (0, \tau]$ such that

$$\Phi_x(\bar{t}) = \delta. \tag{4.18}$$

In particular, from (4.17) it follows that $\bar{t} < T$.

Due to (4.13) and (4.18), we obtain that

$$\Phi'_x(t) > \delta \quad \text{for any } \bar{t} < t < T. \tag{4.19}$$

By (4.12), in particular we have

$$\Phi'_x(t) \geq h(\Phi_x(t)) \quad \text{for any } \bar{t} < t < T. \tag{4.20}$$

Define

$$G(t) := \int_{\Phi_x(t)}^{+\infty} \frac{1}{h(z)} dz \quad \text{for all } \bar{t} < t < T.$$

Note that G is well-defined thanks to hypothesis (3.1) and to (4.19). Furthermore,

$$G'(t) = -\frac{\Phi'_x(t)}{h(\Phi_x(t))} \quad \text{for any } \bar{t} < t < T. \tag{4.21}$$

We now define

$$w(t) := \exp\{G(t)\} \quad \text{for any } \bar{t} < t < T. \tag{4.22}$$

Then, due to (4.21) and (4.20),

$$\begin{aligned} w'(t) &= -\frac{\Phi'_x(t)}{h(\Phi_x(t))}w(t) \\ &\leq -w(t) \quad \text{for any } \bar{t} < t < T. \end{aligned} \tag{4.23}$$

By integrating (4.23) we get:

$$\begin{aligned} w(t) &\leq w(\bar{t}) \exp\left\{-\int_{\bar{t}}^t ds\right\} \\ &\leq w(\bar{t}) \exp\{-(t-\bar{t})\} \quad \text{for any } \bar{t} < t < T. \end{aligned} \tag{4.24}$$

We substitute (4.22) into (4.24), so we have

$$\exp\{G(t)\} \leq \exp\{G(\bar{t}) - (t-\bar{t})\} \quad \text{for any } \bar{t} < t < T.$$

Thus, for any $\bar{t} < t < T$,

$$G(t) \leq G(\bar{t}) - (t-\bar{t}). \tag{4.25}$$

We now combine (4.25) together with (4.16), hence

$$0 \leq G(t) \leq G(\bar{t}) - \left(t + \frac{1}{\alpha} \log(\Phi_x(0)) - \frac{\log \delta}{\alpha}\right) \quad \text{for any } \bar{t} < t < T.$$

Hence

$$\log(\Phi_x(0)) \leq \alpha G(\bar{t}) + \log \delta - \alpha t \quad \text{for any } \bar{t} < t < T. \tag{4.26}$$

We now take the exponential of both sides of (4.26). Thus we get, for any $\bar{t} < t < T$,

$$\begin{aligned} \Phi_x(0) &\leq \exp\{\alpha G(\bar{t}) + \log \delta - \alpha t\} \\ &= C \exp\{-\alpha t\}, \end{aligned}$$

where

$$C := \exp\{\alpha G(\bar{t}) + \log \delta\}. \tag{4.27}$$

This is the inequality (4.6). □

4.2 Proof of Theorem 3.1

Proof Take any $\delta > 0$ fulfilling (4.4). We suppose, by contradiction, that u is a global solution of problem (1.1). Since $\alpha := h'(0) > \lambda_1(M)$, there exists $\varepsilon \in (0, \alpha - \lambda_1(M))$ such that

$$\alpha > \lambda_1(M) + \varepsilon.$$

Let $\Omega \subset M$ be such that

$$\mu(\Omega) < +\infty \quad \text{and} \quad \int_{\Omega} u_0 \, d\mu > 0.$$

Then

$$\lim_{T \rightarrow +\infty} e^{\alpha - (\lambda_1(M) + \varepsilon)T} = +\infty. \tag{4.28}$$

Fix any arbitrary $x \in M$. By Lemmas 4.1 and 4.2,

$$C_1 e^{-[\lambda_1(M) + \varepsilon]T} \leq (e^{T\Delta} u_0)(x) \leq C e^{-\alpha T}, \quad \text{for any } T > \max\{t_0, \bar{t}\},$$

where $t_0 > 0, C_1 > 0$ are given in Lemma 4.1, while $\bar{t} > 0, C > 0$ in Lemma 4.2. Hence, if u exists globally in time, we would have

$$e^{[\alpha - (\lambda_1(M) + \varepsilon)]T} \leq \frac{C}{C_1} \quad \text{for any } T > \max\{t_0, \bar{t}\}. \tag{4.29}$$

Nonetheless, due to (4.28), the left hand side of (4.29) tends to $+\infty$ as $T \rightarrow \infty$. Thus, we have a contradiction. Hence the thesis follows. □

5 Global existence

Consider the linear Cauchy problem for the heat equation

$$\begin{cases} v_t = \Delta v & \text{in } M \times (0, +\infty) \\ v = u_0 & \text{in } M \times \{0\}, \end{cases} \tag{5.1}$$

with u_0 as in Theorem 3.2. Observe that problem (5.1) admits the classical solution

$$v(x, t) = \int_M p(x, y, t) u_0(y) \, dy, \quad x \in M, t \geq 0. \tag{5.2}$$

Hence, since $u_0 \in L^\infty(M)$,

$$\|v(t)\|_{L^\infty(M)} \leq \|u_0\|_{L^\infty(M)} \quad \text{for any } t > 0. \tag{5.3}$$

Moreover, since $u_0 \in L^1(M)$, if the Faber–Krahn inequality holds, then, due to (2.3),

$$v(x, t) \leq \bar{C} \|u_0\|_{L^1(M)} e^{-\lambda_1 t} \quad \text{for any } x \in M, t > 1, \tag{5.4}$$

where \bar{C} has been defined in (2.3).

Let $\{\Omega_j\}_{j \in \mathbb{N}} \subset M$ be a sequence of domains such that

$$\begin{aligned} \Omega_j &\subset \Omega_{j+1} \quad \text{for any } j \in \mathbb{N}, \\ \bigcup_{j \in \mathbb{N}} \Omega_j &= M, \\ \partial\Omega_j &\text{ is smooth for every } j \in \mathbb{N}. \end{aligned}$$

Furthermore, for every $j \in \mathbb{N}$ let $\zeta_j \in C_c^\infty(\Omega_j)$ be such that $0 \leq \zeta_j \leq 1$, $\zeta_j \equiv 1$ in $\Omega_{j/2}$.

Proof of Theorem 3.2 We consider initial data u_0 satisfying (3.3) and (3.4). Define

$$\bar{u}(x, t) := e^{\alpha t} v(x, t), \quad x \in M, t \geq 0,$$

with v and α given by (5.2) and (3.2), respectively.

Note that, due to (3.3) and (5.3), for any $x \in M, t \in (0, 1]$,

$$0 \leq \bar{u}(x, t) \leq e^{\alpha t} \|v(t)\|_{L^\infty(M)} \leq e^{\alpha t} \|u_0\|_{L^\infty(M)} \leq \delta. \tag{5.5}$$

Moreover, due to (3.4), (5.4), since $\alpha \leq \lambda_1(M)$, for any $t > 1$ we get

$$0 \leq \bar{u}(x, t) \leq e^{\alpha t} \|v(t)\|_{L^\infty(M)} \leq \bar{C} \|u_0\|_{L^1(M)} e^{-(\lambda_1 - \alpha)t} \leq \delta. \tag{5.6}$$

Inequalities (5.5) and (5.6) yield

$$0 \leq \bar{u}(x, t) \leq \delta \quad \text{for any } x \in M, t > 0. \tag{5.7}$$

Furthermore, we have

$$\bar{u}_t - \Delta \bar{u} - f(\bar{u}) = \alpha e^{\alpha t} v + e^{\alpha t} v_t - e^{\alpha t} \Delta v - f(\bar{u}).$$

Now, by using the fact that v is a classical solution to problem (5.1), due to (3.2) and (5.7), we get

$$\bar{u}_t - \Delta \bar{u} - f(\bar{u}) \geq \alpha \bar{u} - f(\bar{u}) \geq 0. \tag{5.8}$$

Hence \bar{u} is a weak supersolution to problem (1.1) in $M \times (0, \infty)$.

For any $j \in \mathbb{N}$ there exists a unique classical solution u_j to problem

$$\begin{cases} \partial_t u = \Delta u + f(u) & \text{in } \Omega_j \times (0, T) \\ u = 0 & \text{in } \partial\Omega_j \times (0, T) \\ u = \zeta_j u_0 & \text{in } \Omega_j \times \{0\}. \end{cases} \tag{5.9}$$

Clearly, $u_j \not\equiv 0$ because $u_0 \zeta_j \not\equiv 0$ in Ω_j . Moreover, for any $j \in \mathbb{N}$, in view of (5.8), since

$$v = u_0 \geq \zeta_j u_0 \quad \text{in } M \times \{0\},$$

\bar{u} is a bounded weak supersolution of problem (5.9). Obviously, for any $j \in \mathbb{N}$, $\bar{u} \equiv 0$ is a subsolution to problem (5.9). Hence, by the comparison principle, for every $j \in \mathbb{N}$ we obtain

$$0 \leq u_j \leq \bar{u} \quad \text{for any } (x, t) \in \Omega_j \times (0, +\infty). \tag{5.10}$$

By standard a priori estimates (see, e.g., [15, Chapter 5]), we can infer that there exists a subsequence $\{u_{j_k}\}$ of $\{u_j\}$, which converges in $C_{x,t}^{2,1}(K \times [\varepsilon, T])$ as $k \rightarrow +\infty$, for each compact subset $K \subset M$ and for each $\varepsilon \in (0, T)$, and in $C_{loc}(M \times [0, T])$, to some function $u \in C_{x,t}^{2,1}(M \times (0, T]) \cap C(M \times [0, T])$, which is a classical solution to problem (1.1). Moreover, from (5.10) we get

$$0 \leq u \leq \bar{u} \quad \text{in } M \times (0, +\infty).$$

Hence the thesis follows. □

6 On blow-up of solutions for large data

In this section we discuss a blow-up result that can be obtained by standard tools. More precisely, we show that the solution to problem (1.1) blows up, provided that u_0 is large enough, and $f : [0, +\infty) \rightarrow [0, +\infty)$ is a locally Lipschitz, increasing, convex function fulfilling

$$\int^{+\infty} \frac{1}{f(s)} ds < +\infty.$$

We need to introduce some preliminary material. Let $o \in M$ be a reference point and $r(x)$ be the geodesic distance between x and o . For any $x \in M \setminus \{o\}$, denote by Ric_o the Ricci curvature at x in the radial direction $\frac{\partial}{\partial r}$. We assume that

$$\text{Ric}_o(x) \geq -(N - 1) \frac{\psi''(r(x))}{\psi(r(x))} \quad \text{for all } x \in M \setminus \{o\},$$

for some $\psi \in C^\infty((0, +\infty)) \cap C^1([0, +\infty))$ such that $\psi'(0) = 1, \psi(0) = 0, \psi > 0$ in $(0, +\infty)$ and

$$\int^{+\infty} \frac{\int_0^r \psi^{N-1}(\xi) d\xi}{\psi^{N-1}(r)} dr = +\infty. \tag{6.1}$$

In view of such hypothesis, for problem (1.1) comparison principle for bounded sub- and supersolutions holds (see, e.g., [7, 21]). Condition (6.1) may be stated informally in a quite simpler way: a sufficient condition for this to hold is that

$$\text{Ric}_o(x) \geq -cr(x)^2 \quad \text{for all } x \in M \setminus \{B_1(o)\} \tag{6.2}$$

and a suitable $c > 0$, as can be seen by choosing ψ to be e^{kr^2} in a neighborhood of infinity, for a suitable $k > 0$.

Let D be an open precompact subset of M with smooth boundary. By Kaplan’s method (see [13]) it can be proved that, for some $v_0 \in C(\bar{D}), v_0 \geq 0$ large enough, any solution v to problem

$$\begin{cases} v_t = \Delta v + f(v) & \text{in } D \times (0, T) \\ v = 0 & \text{in } \partial D \times (0, T) \\ v = v_0 & \text{in } D \times \{0\} \end{cases} \tag{6.3}$$

blows up in finite time. Now, consider $u_0 \in C(M), u_0 \geq 0$ with compact support. Take any D as above containing the support of u_0 and set $v_0 := u_{0|D}$. By choosing u_0 big enough, and so v_0 , the solution v to (6.3), corresponding to such v_0 , blows up in a finite time, say $\tau > 0$. Let

$$\underline{u} := \begin{cases} v & \text{in } D \times (0, \tau) \\ 0 & \text{in } (M \setminus D) \times (0, \tau). \end{cases}$$

By the maximum principle,

$$v \geq 0 \quad \text{in } D \times (0, \tau).$$

Hence,

$$\frac{\partial v}{\partial n} \leq 0 \quad \text{in } \partial D \times (0, \tau),$$

n being the outer unit normal vector to ∂D . This easily implies that, for any $0 < T < \tau$, \underline{u} is a bounded weak subsolution to problem (1.1). So, by comparison principle, for any solution u to problem (1.1),

$$u \geq \underline{u} \quad \text{in } M \times (0, T).$$

Since v blows in finite time, the same holds for \bar{u} and so for u .

Acknowledgements The authors are members of the Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA, Italy) of the Istituto Nazionale di Alta Matematica (INdAM, Italy) and are partially supported by the PRIN project 201758MTR2: “Direct and Inverse Problems for Partial Differential Equations: Theoretical Aspects and Applications” (Italy).

Funding Open access funding provided by Politecnico di Milano within the CRUI-CARE Agreement.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bandle, C., Brunner, H.: Blowup in diffusion equations: a survey. *J. Comput. Appl. Math.* **97**, 3–22 (1998)
2. Bandle, C., Pozio, M.A., Tesi, A.: The Fujita exponent for the Cauchy problem in the Hyperbolic Space. *J. Differ. Equ.* **251**, 2143–2163 (2011)
3. Chavel, I., Karp, L.: Large time behavior of the heat kernel: the parabolic λ -potential alternative. *Comment. Math. Helv.* **66**, 541–556 (1991)

4. Deng, K., Levine, H.A.: The role of critical exponents in blow-up theorems: the sequel. *J. Math. Anal. Appl.* **243**, 85–126 (2000)
5. Fujita, H.: On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$. *J. Fac. Sci. Univ. Tokyo Sect. I*(13), 109–124 (1966)
6. Grigor'yan, A.: Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds. *Bull. Am. Math. Soc.* **36**, 135–249 (1999)
7. Grigor'yan, A.: “Heat Kernel and Analysis on Manifolds”, *AMS/IP Studies in Advanced Mathematics*, vol. 47. American Mathematical Society, Providence (2009)
8. Grillo, G., Ishige, K., Muratori, M.: Nonlinear characterizations of stochastic completeness. *J. Math. Pures Appl.* **139**, 63–82 (2020)
9. Grillo, G., Meglioli, G., Punzo, F.: Smoothing effects and infinite time blowup for reaction–diffusion equations: an approach via Sobolev and Poincaré inequalities. *J. Math. Pures Appl.* **151**, 99–131 (2021)
10. Grillo, G., Meglioli, G., Punzo, F.: Global existence of solutions and smoothing effects for classes of reaction–diffusion equations on manifolds. *J. Evol. Equ.* **21**, 2339–2375 (2021)
11. Gu, Q., Sun, Y., Xiao, J., Xu, F.: Global positive solution to a semi-linear parabolic equation with potential on Riemannian manifold. *Calc. Var. PDEs* **59**, 170 (2020)
12. Hayakawa, K.: On nonexistence of global solutions of some semilinear parabolic differential equations. *Proc. Jpn. Acad.* **49**, 503–505 (1973)
13. Kaplan, S.: On the growth of solutions of quasilinear parabolic equations. *Commun. Pure Appl. Math.* **16**, 305–330 (1963)
14. Kobayashi, K., Sirao, T., Tanaka, H.: On the growing up problem for semilinear heat equations. *J. Math. Soc. Jpn.* **29**, 407–424 (1977)
15. Ladyzhenskaya, O.A., Solonnikov, V.A., Ural'tseva, N.A.: *Linear and Quasilinear Equations of Parabolic Type*, Nauka, Moscow (1967) (English translation: series *Transl. Math. Monographs*, 23 AMS, Providence, RI, 1968)
16. Levine, H.: The role of critical exponent in blowup theorems. *SIAM Rev.* **32**, 262–288 (1990)
17. Mastrolia, P., Monticelli, D.D., Punzo, F.: Nonexistence of solutions to parabolic differential inequalities with a potential on Riemannian manifolds. *Math. Ann.* **367**, 929–963 (2017)
18. Meier, P.: On the critical exponent for reaction–diffusion equations. *ARMA* **109**, 63–71 (1990)
19. Meier, P.: Blow-up of solutions of semilinear parabolic differential equations. *ZAMP* **39**, 135–149 (1998)
20. Punzo, F. Global solutions of semilinear parabolic equations with drift on Riemannian manifolds, *DCDS-A (to appear)*. <https://doi.org/10.3934/dcds.2022030>
21. Punzo, F.: Blow-up of solutions to semilinear parabolic equations on Riemannian manifolds with negative sectional curvature. *J. Math. Anal. Appl.* **387**, 815–827 (2012)
22. Samarskii, A.A., Galaktionov, V.A., Kurdyumov, S.P., Mikhailov, A.P.: “Blow-up in Quasilinear Parabolic Equations”, *De Gruyter Expositions in Mathematics*, vol. 19. de Gruyter, Berlin (1995)
23. Vázquez, J.L.: The problems of blow-up for nonlinear heat equations. Complete blow-up and avalanche formation. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei Mat. Appl.* **15**, 281–300 (2004)
24. Wang, Z., Yin, J.: A note on semilinear heat equation in hyperbolic space. *J. Differ. Equ.* **256**, 1151–1156 (2014)
25. Wang, Z., Yin, J.: Asymptotic behaviour of the lifespan of solutions for a semilinear heat equation in hyperbolic space. *Proc. R. Soc. Edinb. Sect. A* **146**, 1091–1114 (2016)
26. Zhang, Q.S.: Blow-up results for nonlinear parabolic equations on manifolds. *Duke Math. J.* **97**, 515–539 (1999)