CORRECTION

Correction to: Population genetic structure and demography of *Magnolia kobus*: variety *borealis* is not supported genetically

Ichiro Tamaki¹ · Naomichi Kawashima^{2,6} · Suzuki Setsuko³ · Jung-Hyun Lee⁴ · Akemi Itaya⁵ · Kyohei Yukitoshi² · Nobuhiro Tomaru²

Published online: 29 April 2020
© The Author(s) 2020

Correction to:

Journal of Plant Research (2019) 132:741–758
<https://doi.org/10.1007/s10265-019-01134-6>

The article Population genetic structure and demography of *Magnolia kobus*: variety *borealis* is not supported genetically, written by Kohma Ichiro Tamaki, Naomichi Kawashima, Suzuki Setsuko, Jung Hyun Lee, Akemi Itaya, Kyohei Yukitoshi and Nobuhiro Tomaru, was originally published Online First without Open Access. After publication in volume 132, issue 6, page 741–758 the author decided to opt for Open Choice and to make the article an Open Access publication. Therefore, the copyright of the article has been changed to © The Author(s) 2020 and the article is forthwith distributed under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits use, duplication, adaptation, distribution and reproduction in any

medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

The original article was updated.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10265-019-01134-6>.

✉ Nobuhiro Tomaru
tomaru@agr.nagoya-u.ac.jp

¹ Gifu Academy of Forest Science and Culture, 88 Sodai, Mino, Gifu 501-3714, Japan

² Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8601, Japan

³ Department of Forest Molecular Genetics and Biotechnology, Forestry and Forest Products Research Institute, Forest Research and Management Organization, 1 Matsunosato, Tsukuba, Ibaraki 305-8687, Japan

⁴ Department of Biology Education, Chonnam National University, 77 Yongbong-ro, Buk-gu, Gwangju 500-757, Republic of Korea

⁵ Graduate School of Bioresources, Mie University, 1577 Kurimamachiya, Tsu, Mie 514-8507, Japan

⁶ Present Address: Mie Prefecture Forestry Research Institute, 3769-1 Nihongi, Hakusan-cho, Tsu, Mie 515-2602, Japan