## CORRECTION





## Correction to: Morphological and antioxidant responses of *Nopalea cochenillifera* cv. Maya (edible *Opuntia* sp. "Kasugai Saboten") to chilling acclimatization

Ayumu Kondo<sup>1</sup> · Masashi Ito<sup>1</sup> · Yusaku Takeda<sup>1</sup> · Yuka Kurahashi<sup>1</sup> · Shigeo Toh<sup>1</sup> · Toru Funaguma<sup>1</sup>

Published online: 7 March 2023 © The Author(s) 2023

## **Correction to:**

Journal of Plant Research (2023) 136:211–225 https://doi.org/10.1007/s10265-023-01437-9

The article Morphological and antioxidant responses of Nopalea cochenillifera cv. Maya (edible Opuntia sp. "Kasugai Saboten") to chilling acclimatization, written by Ayumu Kondo, Masashi Ito, Yusaku Takeda, Yuka Kurahashi, Shigeo Toh, Toru Funaguma, was originally published Online First without Open Access. After publication in volume 136, issue 2, page 211–225 the author decided to opt for Open Choice and to make the article an Open Access publication. Therefore, the copyright of the article has been changed to © The Author(s) 2023 and the article is forthwith distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons

licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

The original article has been updated.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at https://doi.org/10.1007/s10265-023-01437-9.

Ayumu Kondo ayumu@meijo-u.ac.jp

Faculty of Agriculture, Meijo University, 1–501 Shiogamaguchi, Tempaku, Nagoya 468-8502, Japan

