

Extracellular matrix of *Magnaporthe oryzae* may have a role in host adhesion during fungal penetration and is digested by matrix metalloproteinases

**Kanako Inoue · Tomoko Suzuki · Kenichi Ikeda ·
Shan Jiang · Naoki Hosogi · Gang-Su Hyon ·
Sizuka Hida · Taku Yamada · Pyoyun Park**

Published online: 10 January 2008
© The Phytopathological Society of Japan and Springer 2008

**Erratum to: J Gen Plant Pathol (2007) 73:388–398
DOI 10.1007/s10327-007-0048-2**

The correct name of the sixth author should be given as
Gang-Su Hyon, not Gang-Su Hyong.

The online version of the original article can be found under
doi:[10.1007/s10327-007-0048-2](https://doi.org/10.1007/s10327-007-0048-2).

K. Inoue · T. Suzuki · K. Ikeda · S. Jiang · N. Hosogi ·
G.-S. Hyon · S. Hida · T. Yamada · P. Park (✉)
Laboratory of Stress Cytology,
Graduate School of Science and Technology,
Kobe University, Nada, Kobe 657-8501, Japan
e-mail: ppark@kobe-u.ac.jp

Present Address:
T. Suzuki
Laboratory of Plant Pathology and Genetic Engineering,
Faculty of Agriculture, Okayama University,
Okayama, Japan