



Correction to: Chlorantraniliprole application differentially affects adult emergence of *Sympetrum* dragonflies in rice paddy fields

Kosuke Nakanishi¹ · Nisikawa Usio² · Hiroyuki Yokomizo¹ · Tadao Takashima³ · Takehiko I. Hayashi⁴

Published online: 30 January 2022
© The Author(s) 2022

Correction to: Paddy and Water Environment
<https://doi.org/10.1007/s10333-021-00880-5>

The article “Chlorantraniliprole application differentially affects adult emergence of *Sympetrum* dragonflies in rice paddy fields”, written by “Kosuke Nakanishi, Nisikawa Usio, Hiroyuki Yokomizo, Tadao Takashima and Takehiko I. Hayashi”, was originally published electronically on the publisher’s Internet portal on 22 November 2021 without open access. With the author(s)’ decision to opt for Open Choice, the copyright of the article changed on 13 December 2021 to The Author(s) 2022 and the article is forthwith distributed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise

in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0>.

The original article has been corrected.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0>.

The original article can be found online at <https://doi.org/10.1007/s10333-021-00880-5>.

✉ Kosuke Nakanishi
nakanishi.kosuke@nies.go.jp

- ¹ Health and Environmental Risk Division, National Institute for Environmental Studies, Onogawa 16-2, Tsukuba, Ibaraki 305-8506, Japan
- ² Institute of Nature and Environmental Technology, Kanazawa University, Kakuma-machi, Kanazawa, Ishikawa 920-1192, Japan
- ³ Haku Agricultural Cooperatives (JA Haku), Ota-machi To 105, Haku, Ishikawa 925-8588, Japan
- ⁴ Social Systems Division, National Institute for Environmental Studies, Onogawa 16-2, Tsukuba, Ibaraki 305-8506, Japan