

Erratum to: Predicting the potential distribution of the invasive Common Waxbill *Estrilda astrild* (Passeriformes: Estrildidae)

Darius Stiels · Kathrin Schidelko · Jan O. Engler ·
Renate van den Elzen · Dennis Rödder

Published online: 19 August 2011
© Dt. Ornithologen-Gesellschaft e.V. 2011

Erratum to: J Ornithol (2011) 152:769–780 DOI 10.1007/s10336-011-0662-9

An erroneous range for the LDA overlap analyses in multivariate E-space was stated in the methods section (subsection: Comparison of climatic niches). Instead of a range of 0 (no overlap) to 1 (total overlap), the parameter ranges only from 0 (no overlap) to 0.5 (total overlap). Due to random group predictions when two groups cover

identical E-space, the maximum discriminance value cannot exceed 0.5. Thus, an overlap of 0.32 (± 0.057) (results section, subsection: Comparison of climate niches) rather indicates a “modest” than a “low” overlap. However, general conclusions as stated in the discussion section are not altered as the climate at invasive locations still nests within the substantially larger occupied E-space of the native population.

The online version of the original article can be found under
doi:[10.1007/s10336-011-0662-9](https://doi.org/10.1007/s10336-011-0662-9).

D. Stiels (✉) · K. Schidelko · J. O. Engler · R. van den Elzen ·
D. Rödder
Zoological Research Museum Alexander Koenig,
Adenauerallee 160, 53113 Bonn, Germany
e-mail: d.stiels.zfmk@uni-bonn.de

J. O. Engler · D. Rödder
Biogeography Department, Trier University,
54286 Trier, Germany