




Correction to: Determination of Commonly Used Excipients in Pharmaceutical Preparations by Microchip Electrophoresis with Conductivity Detection

Peter Troška¹ · Ewa Poboży² · Zuzana Némethová³ · Marián Masár¹ 

Published online: 21 July 2021
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: Chromatographia 82, 741–748 (2019)
<https://doi.org/10.1007/s10337-019-03691-3>

The authors regret the following mistake in the article:
the value for analyte Methylparaben in sample Paxeladine, in the column “Recovery, mean \pm SD (%)” of the Table 4, is incorrect. The value “96.6 \pm 14.2” should read “96.6 \pm 5.7”.

The original article can be found online at <https://doi.org/10.1007/s10337-019-03691-3>.

✉ Marián Masár
marian.masar@uniba.sk

- ¹ Department of Analytical Chemistry, Faculty of Natural Sciences, Comenius University in Bratislava, Mlynská dolina CH-2, Ilkovičova 6, 84215 Bratislava, Slovakia
- ² Department of Chemistry, University of Warsaw, Pasteura 1, 02093 Warsaw, Poland
- ³ Division of Natural Sciences and Forensic Analyses, Institute of Forensic Science, Sklabinská 1, 81272 Bratislava, Slovakia