

Erratum to: Leaf Removal Treatments Combined with Kaolin Particle Film Technique from Different Directions of Grapevine's Canopy Affect the Composition of Phytochemicals of cv. Muscat Hamburg (*V. Vinifera* L.)

Demir Kok¹ · Erdinc Bal¹

Published online: 18 August 2017
© Springer-Verlag GmbH Deutschland 2017

Erratum to:
Erwerbs-Obstbau 2017
DOI 10.1007/s10341-017-0337-7

The original version of this article unfortunately contained errors in Figs. 1, 3, 4, 5, 6, 7 and 8. The corrected figures are given here.

The original article has been corrected.




Fig. 1 Influences of different treatments on total soluble solids content




Fig. 3 Influences of different treatments on juice pH




Fig. 4 Influences of different treatments on *p*-value of juice

The online version of the original article can be found under doi:
[10.1007/s10341-017-0337-7](https://doi.org/10.1007/s10341-017-0337-7)

✉ Demir Kok
dkok@nku.edu.tr

¹ Agricultural Faculty, Department of Horticulture, Namık Kemal University, 59030 Tekirdağ, Turkey




Fig. 5 Influences of different treatments on total phenolic compounds content




Fig. 6 Influences of different treatments on total anthocyanin content




Fig. 7 Influences of different treatments on free volatile terpenes content




Fig. 8 Influences of different treatments on potentially volatile terpenes content