



CORRECTION



Correction to: Effectiveness of a photoscreener in identifying undiagnosed unilateral amblyopia at vision screening of 3-year-old children in Japan

Shiori Harada¹ · Yumiko Nakashima² · Masafumi Uematsu¹ · Shinpei Morimoto³ · Yasser Helmy Mohamed¹ · Takashi Kitaoka¹ · Hiroyuki Moriuchi²

Published online: 9 April 2022
© Japanese Ophthalmological Society 2022

Correction to:
Japanese Journal of Ophthalmology (2022) 66:193–198
<https://doi.org/10.1007/s10384-021-00896-8>

In the original publication, under Results section, in second paragraph, the sentence: “Of the 52 children...” should read as: Of the 52 children included in the study, the major reason for photoscreener referral was astigmatism (65.3%), followed by hyperopia (18.4%) (Table 2).

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Corresponding author: Masafumi Uematsu.

The original article can be found online at <https://doi.org/10.1007/s10384-021-00896-8>.

✉ Masafumi Uematsu
uematsu1124@outlook.jp

¹ Department of Ophthalmology and Visual Sciences, Graduate School of Biomedical Sciences, Nagasaki University, 1-7-1 Sakamoto, Nagasaki, Nagasaki 852-8501, Japan

² Department of Pediatrics, Nagasaki University Hospital, 1-7-1 Sakamoto, Nagasaki, Nagasaki 852-8501, Japan

³ Clinical Research Center, Nagasaki University Hospital, 1-7-1 Sakamoto, Nagasaki, Nagasaki 852-8501, Japan