Correction to: The early therapeutic response at 2 weeks is a crucial predictor of proton pump inhibitor-refractory gastroesophageal reflux disease

Maiko Ogawa¹ · Seiji Arihiro¹ · Nobuyuki Matsushashi² · Takashi Joh³ · Kazuhide Higuchi⁴ · Katsuhiko Iwakiri⁵ · Takeshi Kamiya⁶ · Noriaki Manabe⁷ · Kimio Isshi⁸ · Tatsuya Nakada¹ · Atsushi Hokari¹ · Masayuki Saruta⁹ · Atsushi Oshio¹⁰ · Ken Haruma¹¹ · Koji Nakada¹²

Published online: 8 April 2021
© The Japan Esophageal Society 2021

Correction to: Esophagus (2021) 18:398–406
<https://doi.org/10.1007/s10388-020-00792-z>

In the original publication of the article, Fig. 1a was published incorrectly. The number of Non-responder at 2 weeks was published as 11. The correct value is 114.

The correct Fig. 1 is provided below.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10388-020-00792-z>.

✉ Koji Nakada
nakada@jikei.ac.jp

¹ Division of Gastroenterology and Hepatology, Department of Internal Medicine, Katsushika Medical Center, The Jikei University School of Medicine, Tokyo, Japan

² Department of Gastroenterology, NTT Medical Center Tokyo, Tokyo, Japan

³ Gamagori City Hospital, Aichi, Japan

⁴ Second Department of Internal Medicine, Osaka Medical College, Osaka, Japan

⁵ Department of Gastroenterology, Nippon Medical School, Graduate School of Medicine, Tokyo, Japan

⁶ Department of Medical Innovation, Nagoya City University Graduate School Medical Sciences, Nagoya, Japan

⁷ Division of Endoscopy and Ultrasonography, Department of Laboratory Medicine, Kawasaki Medical School General Medical Center, Okayama, Japan

⁸ Isshi Gastro-Intestinal Clinic, Tokyo, Japan

⁹ Division of Gastroenterology and Hepatology, Department of Internal Medicine, The Jikei University School of Medicine, Tokyo, Japan

¹⁰ Faculty of Letters, Arts and Sciences, Waseda University, Tokyo, Japan

¹¹ Department of General Internal Medicine 2, Kawasaki Medical School Kawasaki Hospital, Okayama, Japan

¹² Department of Laboratory Medicine, The Jikei University School of Medicine, 3-25-8, Nishi-shimbashi, Minato-ku, Tokyo 105-8461, Japan

Fig. 1 Responder rates at 2 and 4 weeks of PPI treatment. **a** Responder definition by GERD-SS residual symptom rate $\leq 50\%$; **b** responder definition by patient's impression as "improved" or better (Q11); **c** responder definition by numeric rating scale ≤ 5 (Q12). *PPI* proton pump inhibitor; *GERD-SS* gastroesophageal reflux disease-subscale; *Sx* symptom