Editorial

Snezana Simic

Published online: 2 May 2010

© Springer-Verlag 2010

The current issue of the Journal of Public Health reviews complex and interlinked health determinants around the world. From New South Wales in Australia to Europe, each destination presents an intellectual challenge to understanding some intriguing public health aspects. Social, environmental, and biological determinants of health, as well as population health and health in specific groups were explored, and actions at different levels proposed. All printed articles unequivocally show that societies today face numerous threats that prompt public health professionals to look for a new way of preventing their potentially serious consequences. It is also clear that knowledge and understanding of the determinants of health are remarkably deficient, which shows the complexity of health and incorporates expertise from specialists in different disciplines.

The first group of articles explores the contribution of sociodemographic factors and their effects on ill health. Small area variation of cardio-surgery procedure rates according to sociodemographic characteristics was explored by Weerasinghe and colleagues in Australia, while Geisel-Marbaise and Stummer tried to identify gender differences in the health behavior pattern of diabetic patients' adherence to their medication regimen in Germany. Effects of social differences on cancer incidence and mortality in the population of northwest Germany were examined by Eberle and colleagues.

Some environmental and behavioral risk factors' effects on the transmission of *Helicobacter pylori* in Egypt were explored by Awdalla and colleagues; Attarchi and colleagues

appraised hearing loss brought on by a combination of smoking and continued exposure to noise in Iranian automobile industrial workers, while an assessment of the share of permanent disability pensions that can be traced back to risky health behavior among the working population in Denmark was delivered by Lund and colleagues.

Another group of articles was devoted to evaluation studies. Knowledge and practice were examined in suburban slums in India where Panigrahi assessed this aspect regarding iodized salt among retail shopkeepers, and Grunert and colleagues investigated the use and understanding of nutritional information on food labels as a precondition to healthy eating in six European countries.

Evaluation of the impact of implementing a customized training program for an integrated smoking cessation intervention in TB DOTS health workers was carried out by Awaisu and his colleagues in Malaysia, and from Serbia comes our article on teaching evaluation of the newly established Master of Health Policy and Management program at the School of Public Health.

A specific approach was developed by Porzsolt and colleagues in summarizing the deficiencies and weaknesses of the most commonly used methods for allocation of health care resources, i.e., quality-adjusted life years and efficiency frontier where groups of professionals were set up to discuss the assessment and appraisal of health services.

The starting point for assessing the effects of all listed factors on health is a benchmark that would help us determine where our greatest needs lie. Methodological approaches in those articles used all available tools, from an interview, questionnaire, and utilization of large databases, to knowledge generation and synthesis. The results provide a sound basis for activities aimed at improving social conditions, promoting healthy behavior, and protecting people from environmental hazards. Although the health

S. Simic (✉)
Institute of Social Medicine, School of Medicine,
Belgrade University,
Belgrade, Serbia
e-mail: snezanas@eunet.rs

problem patterns confronting countries differ, the role of public health is almost the same. Suggested interventions include education of people (particularly those of poor socioeconomic status) in lifestyle management, urgent curricula reforms related to the gap in public health education, and involvement of other stakeholders outside the health system because they can provide up-to-date information that represents the actual population circumstances and can serve as indicators for when to intervene. The focus on determinants of health demonstrates the need for multisectorial action and multiple approaches.

Careful reading of all these articles reveals that “public health has evolved as a multidisciplinary field that includes the use of basic and applied science, education, the social

sciences, economics, management and communication skills to promote the welfare of the individual and the community” (Tulchinsky and Varavikova, 2000). By applying this broad perspective, this issue of the Journal of Public Health should help readers appreciate the expanding knowledge and address the present and future health challenges.

References

- Tulchinsky TH, Varavikova EA (2000) *The New Public Health. An introduction for the 21st century*, 2nd edn. Academic Press, San Diego