Erratum to: Outcomes of medical management of peripheral arterial disease in general practice: follow-up results of the PACE-PAD Study

Anja Neumann · Rebecca Jahn · Curt Diehm ·
Elke Driller · Franz Hessel · Gerald Lux ·
Oliver Ommen · Holger Pfaff · Uwe Siebert ·
David Pittrow · Jürgen Wasem ·
on behalf of the Patient Care Evaluation-Peripheral
Arterial Disease (PACE-PAD) Study Investigators

Published online: 3 November 2010
© Springer-Verlag 2010

Erratum to: J Public Health
DOI 10.1007/s10389-010-0356-4

Introduction

The original version of this article unfortunately contained a mistake. The presentation of Table 1 was incorrect. The corrected Table 1 is given below.

The study group was supported by an advisory board founded in 2003 with the following members: Hans Jürgen Ahrens, Curt Diehm, Leonhard Hansen (until 2004), Klaus-Dieter Kossow.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10389-010-0356-4>.

A. Neumann (✉) · R. Jahn · F. Hessel · G. Lux · J. Wasem
Alfred Krupp von Bohlen und Halbach-Stiftungslehrstuhl für
Medizin-Management, Universität Duisburg-Essen,
Schützenbahn 70,
45127 Essen, Germany
e-mail: anja.neumann@uni-due.de

A. Neumann
e-mail: pace-pad@uni-due.de

C. Diehm
Klinikum Karlsbad-Langensteinbach,
Karlsbad, Germany

E. Driller · O. Ommen · H. Pfaff
Zentrum für Versorgungsforschung, Universität zu Köln,
Cologne, Germany

U. Siebert
Department of Public Health, Medical Decision Making and
Health Technology Assessment, University for Health Sciences,
Medical Informatics and Technology (UMIT),
Hall, Austria

D. Pittrow
Institute for Clinical Pharmacology, Medical Faculty Carl Gustav
Carus, Technical University,
Dresden, Germany

Table 1 Guideline-oriented therapy: definitions of quality indicators

Item	Definition	Specification for Fontaine stage
Anticoagulant/antiplatelet therapy	Prescription of at least 1 drug	I– IV
If present: smoking	Cessation advice, if applicable	All stages
Exercise training	Prescription or recommendation of regular training	I and II only
If present: diabetes mellitus	Dietary measure or at least 1 anti diabetic drug	All stages
If present: hypercholesterolemia	Dietary measure or at least 1 lipid-lowering drug	I– IV
If present: hypertension	Dietary measure or at least 1 antihypertensive drug	All stages