Correction to: Unmet need for mental health care within the Dutch population: exploring the role of GP

Eduard R. Wix¹ · Mark Spigt^{2,3}

Published online: 24 February 2022

© Springer-Verlag GmbH Germany, part of Springer Nature 2022

Correction to: Journal of Public Health

<https://doi.org/10.1007/s10389-021-01687-8>

The original version of this article unfortunately contained a mistake. The text under Fig. 1 is missing.

The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s10389-021-01687-8>.

✉ Eduard R. Wix
e.wix@maastrichtuniversity.nl
Mark Spigt
m.spigt@maastrichtuniversity.nl

¹ Department of Family Medicine, Maastricht University, Peter Debyeplein 1, Maastricht 6229, HA, The Netherlands

² Department of Family Medicine, CAPHRI School for Public Health and Primary Care, Maastricht University, Peter Debyeplein 1, Maastricht 6229, HA, The Netherlands

³ General Practice Research Unit, Department of Community Medicine, The Arctic University of Tromsø, Tromsø, Norway

Fig. 1 Onset of mental complaints

† 181 of 183 participants with high distress answered the questions about perceived need for care

*This represents the participants with a need for care according to our definition(21% of total study population)

**Participants with a need that did not receive treatment(44% of patients with a need for care)

***Participants with a need that did receive some treatment(56% of patients with a need for care)