



## Correction to: Rarefied gas flow through a rough channel into a vacuum


O. Sazhin<sup>1</sup>

Published online: 20 April 2020  
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

**Correction to:**  
**Microfluidics and Nanofluidics (2020) 24:27**  
<https://doi.org/10.1007/s10404-020-2330-y>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

In Table 3, there is a mistype in one cell [first row, fifth column]. Here, at  $l/h = 1$ ,  $\delta = 0$  and  $\alpha = 45^\circ$ , the mass flow rate should be  $Q^* = 0.6100$  (not 0.6010). As a result, Fig. 2a should be different, i.e. the minimum  $Q^*$  at  $\delta = 0$  and  $\alpha = 45^\circ$  disappears. Below is the correct version of Fig. 2a.



**Fig. 2 a** The relative gas flow rate  $Q/Q_{sm}$  through a modeled rough channel into a vacuum, as a function of inclination angle  $\alpha$  and rarefaction parameter  $\delta$  for the length to height ratio  $l/h = 1$

The original article can be found online at <https://doi.org/10.1007/s10404-020-2330-y>.

✉ O. Sazhin  
oleg.sazhin@urfu.ru

<sup>1</sup> Ural Federal University, Lenin av. 51, Ekaterinburg 620000, Russia