Stochastic flowering phenology in *Dactylis glomerata* populations described by Markov chain modelling

Carl A. Frisk · Beverley Adams-Groom · Carsten A. Skjøth

Received: 17 August 2020 / Accepted: 17 December 2020 / Published online: 3 February 2021
© Crown 2021

Abstract Understanding the relationship between flowering patterns and pollen dispersal is important in climate change modelling, pollen forecasting, forestry and agriculture. Enhanced understanding of this connection can be gained through detailed spatial and temporal flowering observations on a population level, combined with modelling simulating the dynamics. Species with large distribution ranges, long flowering seasons, high pollen production and naturally large populations can be used to illustrate these dynamics. Revealing and simulating species-specific demographic and stochastic elements in the flowering process will likely be important in determining when pollen release is likely to happen in flowering plants. Spatial and temporal dynamics of eight populations of *Dactylis glomerata* were collected over the course of two years to determine high-resolution demographic elements. Stochastic elements were accounted for using Markov chain approaches in order to evaluate tiller-specific contribution to overall population dynamics. Tiller-specific developmental dynamics were evaluated using three different RV matrix

correlation coefficients. We found that the demographic patterns in population development were the same for all populations with key phenological events differing only by a few days over the course of the seasons. Many tillers transitioned very quickly from non-flowering to full flowering, a process that can be replicated with Markov chain modelling. Our novel approach demonstrates the identification and quantification of stochastic elements in the flowering process of *D. glomerata*, an element likely to be found in many flowering plants. The stochastic modelling approach can be used to develop detailed pollen release models for *Dactylis*, other grass species and probably other flowering plants.

Keywords Pollen · Population statistics · Anthesis · Grass · Poaceae · UK

1 Introduction

Grass ecology is of considerable importance to human society and culture, mainly through cereal cultivation, animal fodder and forestry but also through grass pollen allergy. Grass pollen is arguably the most important outdoor aeroallergen with the highest amount of sensitization (D’Amato et al. 2007; Heinzlerling et al. 2009). Studies have shown that there is a strong relationship between airborne pollen

Supplementary Information The online version contains supplementary material available at (<https://doi.org/10.1007/s10453-020-09685-1>).

C. A. Frisk (✉) · B. Adams-Groom · C. A. Skjøth
National Pollen and Aerobiological Research Unit, School
of Science and the Environment, University of Worcester,
Worcester WR2 6AJ, UK
e-mail: c.frisk@worc.ac.uk

concentrations and the flowering phenology of grasses (Ghitarrini et al. 2017; Kmenta et al. 2016; Romero-Morte et al. 2020). The identification and modelling of demographic and stochastic processes in grass flowering should increase understanding of this relationship.

To fully understand the ecology of grasses, a population-based approach is often needed (Bolnick et al. 2003; Watkinson and Ormerod 2001). Spatial and temporal observations of flowering of multiple populations are often used to understand demographic elements in the study of grasses in agriculture (Rossignol et al. 2014; Smith 1944), ecology (Eagles 1972; Lindner and Garcia 1997) and aerobiology (Rojo et al. 2017; Romero-Morte et al. 2018). Combining this approach with a stochastic Markov chain modelling approach (Balzter 2000) and a model grass species may provide deeper multifaceted understanding with respect to underlying flowering processes. Markov chain approaches have previously been used to accurately model demographic and stochastic elements in grass populations (Nakaoka 1996) (e.g. Canales et al. 1994; Silva et al. 1991).

The species *Dactylis glomerata*, being (probably) the only species in its genus, has been investigated for more than a century regarding its growth and flowering processes (Wolfe 1925; Wolfe and Kipps 1925, 1926). This has been due to its widespread use as a common pasture grass (Eagles and Williams 1971; Mizianty 1986) and its cross-pollination capacity (Jones and Newell 1948). It occurs abundantly throughout its extensive biogeographic distribution, with a native range between the USA and Japan, and from northern Africa to northern Sweden (Clayton et al. 2002). The species can be divided into a few different sub-species based on genetic and morphological divergencies (Yan et al. 2016). However, *D. glomerata subsp. glomerata* is by far the most common sub-species, defining the standard of the species (Mizianty 1990). Cebrino et al. (2018) found that *D. glomerata* starts flowering around the same time in Spain, regardless of habitat. This holds true between years and locations (Cebrino et al. 2016). Observations in cultivated gardens have shown that earlier starting times promote longer flowering in individual flowers (Tormo-Molina et al. 2015), common in wild grass species and important for climate change scenarios (Cleland et al. 2006; Munson and Long 2017). The higher pollen production in *D.*

glomerata compared to other grasses (Aboulaich et al. 2009; Prieto-Baena et al. 2003) heightens the relevance of these processes. All the above-mentioned considerations make *D. glomerata* a model species for phenological investigation. Information about the detailed flowering dynamics in populations will enable climatic and mechanistic modelling of *D. glomerata* and other grasses with similar behaviour, a key aspect for phenological modelling of grasses and their impact on the grass pollen seasonality (García-Mozo et al. 2009). The inclusion of stochastic elements in the phenological development will be able to account for intrinsic attributes of the flowering process, which has been suggested to have implications for pollen and seed dispersal patterns (Soons et al. 2004; Tufto et al. 1997), processes-based range modelling (Evans et al. 2016) and cellular differentiation modelling (Davila-Velderrain et al. 2016). Our objective in this study was to investigate demographic and stochastic elements in grass flowering phenology by using *Dactylis glomerata* as a model species. This was done by incorporating population dynamics through detailed spatial and temporal demographic observations and by using Markov chain modelling to explain the stochastic behaviour.

2 Materials and methods

2.1 Locations

Eight locations across the surroundings of Worcester, UK (Fig. 1), were monitored over two years (15 May to 31 August 2017 and 2018) to investigate *D. glomerata* population dynamics (Table 1). The population at each location were considered as biologically distinct unit since it was spatially isolated from other specimens of the same species within the immediate surroundings. The locations are distinguished between two main populations at St Johns and Lakeside campuses belonging to the University of Worcester and six other unique locations each containing one secondary population. The populations at both campus sites were chosen as main populations due to increased accessibility and lower chances of disturbance. All locations were selected based on three criteria: 1) should contain more than 150 tillers (grass flowering stems) of *D. glomerata* within a 10 m x 10 m area, 2) should not be a road verge and 3) should

Fig. 1 Map showing the eight locations utilized in the phenological observations of *Dactylis glomerata*. The black icons represent the two main locations, while the white icons represent the six secondary locations

be easily accessible. A large minimum population size would limit the variation contributed from each tiller and produce a smaller and more stable standard error for the overall population development (Fox and

Kendall 2002). The microclimate contributed to road verges would produce uncertainty regarding the stability of population dynamics (Jantunen et al. 2007), coupled with aspects of disturbances such as

Table 1 Population type, number of individual tillers, coordinates and surrounding habitat for all locations visited during the flowering phenological observations in the populations of *Dactylis glomerata*

Location	Population Type	Tillers # 2017/2018	Latitude	Longitude	Habitats
St Johns	Main	317/355	52.196362 N	−2.241592 E	Park Landscapes, Uncut Grass Areas
Lakeside	Main	0/535	52.254039 N	−2.257216 E	Agricultural Fields, Uncut Grass Areas
Lower Lakeside	Secondary	0/210	52.226128 N	−2.255737 E	Agricultural Fields
Football Field	Secondary	0/160	52.197030 N	−2.240321 E	Park Landscapes, Uncut Grass Areas
Upper Bridge	Secondary	0/418	52.194382 N	−2.232060 E	Verges
West Field	Secondary	0/427	52.186106 N	−2.260664 E	Uncut Grass Areas
Lower Field	Secondary	0/322	52.180686 N	−2.225638 E	Uncut Grass Areas
Lower Bridge	Secondary	0/245	52.176371 N	−2.225622 E	Uncut Grass Areas, Verges

cutting or grazing (Brock et al. 1996). Easily accessible locations were due to simplicity and reproducibility of the study.

2.2 Grass flowering phenology

The flowering phenology was investigated using the ‘Biologische Bundesanstalt, Bundessortenamt und Chemische Industrie’ (BBCH) scale (Meier 2018). The scale, originally designed to monitor phenological changes in agricultural crops, functions by systematically observing a primary growth characteristic and a secondary growth characteristic, resulting in a two-number sequence. By observing changes to the sequence, detailed phenological change can be monitored (Cornelius et al. 2011, 2014; Meier et al. 2009). The primary growth characteristic focused on in the study is the flowering stage, which starts at tiller heading and ends in tiller senescence. The secondary growth characteristics are focused on the percentage of visible anthers. Each tiller is assigned one of six phases, representing an amount range of visible anthers (Table 2). The percent of visible anthers is the proxy of flowering progression due to the

increased potential of pollen release, the ultimate evolutionary and ecological goal of Poaceae-specific flowering (Khanduri 2011). The tiller is assigned phase p0 until sign of first flowering and phase p5 when the last anther has detached. The reasoning behind last anther detachment is due to the slight theoretical possibility that pollen could still be emitted from desiccated anthers. Phases p1–p4 represent fractions of flowering anthers in steps of 25%. Examples of *Dactylis glomerata* tillers in each phase can be found in the supplementary material (Figs. S1–S6). By observing each tiller within a population, it is possible to measure the phenological progression of the population. Regular observations will allow for a detailed progression profile and reveal inter- (between population) and intra- (within population) demographic variation into flowering phenology through population dynamics of the species.

2.3 Observational strategy

Each population is defined as all tillers of the species within two to four 1.5 m × 1.5 m plots with less than 3 m between each plot within each location. The

Table 2 Phases and criteria used to identify the progression of flowering phenological observations in the populations of *Dactylis glomerata*

Phase	Abbreviation	Criteria	BBCH number
0	P0	Between plant emergence and start of flowering	60
1	P1	Between 1 and 25% of tillers’ anthers are extruded	61
2	P2	Between 26 and 50% of tillers’ anthers are extruded	63
3	P3	Between 51 and 75% of tillers’ anthers are extruded	65
4	P4	Between 76 and 100% of tillers’ anthers are extruded	68
5	P5	All tillers’ anthers have detached	69

reasoning is to allow a population size large enough to capture statistically unlikely events in regard to population demographic stochasticity (Kendall and Fox 2002). Denser populations utilized the lower number of plots and sparser populations the higher number of plots. Phenological progression of all tillers in each population was observed from the first flowering tiller to the last tillers' senescence. The reasoning is to allow the full range of population dynamics within the main population locations, as adopted from previous grass population studies (e.g. Liddle et al. 1982). During 2017, the main population located in St Johns was visited once a day to investigate flowering progression. During 2018, each of the eight populations was visited. The main populations located in St Johns and Lakeside were visited every second day to investigate whether there was a difference in locations between years. All tillers at the main locations were marked with an ID tag; this allowed observations of tiller-specific progression through the season. The six secondary populations were each visited ten times over the 2018 seasons, resulting in a weekly observational frequency, widely used in grass phenology studies (e.g. García-Mozo et al. 2010; Jochner et al. 2012). Note that the population located at the location Lower Field was only visited five times, due to silage cutting, which removed all tillers. The observations of the six secondary populations stopped as the first main population reached full senescence. The distinctions between main and secondary populations are the continuity of visits, number of visits and the ID tagging of tillers. The use of frequent and continuous observations enhanced the resolution of the phenological progression within the main populations, with the St Johns 2017 population having twice the temporal resolution of the two 2018 populations.

2.4 Analytical framework

2.4.1 General flowering progression of all populations

The general flowering progression was investigated for each year and location by incorporating the average phase of all tillers at each available timestep. A mean phenological progression was then calculated per timestep and year based on all available populations. Due to the observational strategy employed,

some dates were missing flowering values. A mean value was thus estimated using the two surrounding values. For 2017, the population at St Johns was used for this mean progression. For 2018, all eight populations were incorporated into the calculation and combined with a standard error.

2.4.2 Markov chain model

The Markov chain stochastic modelling approach was utilized to analyse tiller-specific dynamics within each of the two main populations (in a similar way as Tseng et al. (2020)) for the years 2017 and 2018. The stochastic model uses the records of the temporal phase change for each tiller, by summarizing the overall phase change of all tillers into a six-by-six transitional matrix. The R package *markovchain* (Spedicato 2017) was used to create the transitional matrices. Not all phase changes are possible due to the tiller not being able to revert in phase or go directly from non-flowering to end-flowering (within the daily/bi-daily observational approach). This matrix describes the likelihood for flowering progression (or phase change) for the full season for all tillers in each of the main populations (see approach in Meyn and Tweedie (1993)). These transitional matrices can be translated into Markov chain diagrams, which display the directional change between phases and the likelihood expressed between zero and one of transitioning for all possible transitions within each population (Grinstead and Snell 1997). Note that this analysis was not possible for the secondary locations, in the same manner, due to the limited number of visits, thereby not being able to observe tiller-specific progression.

2.4.3 Correlations

The general flowering progression for each year was tested for statistical normality using the Shapiro non-normality test (Shapiro and Wilk 1965). The nonparametric Spearman rho correlation was utilized to analyse the general flowering progression due to non-normality expressed by the data. The rank-order correlation produces a more reliable estimate in cases of violated parametric assumptions (Spearman 1904). Correlation was tested without significance testing due to the strong autocorrelations found in accumulated time series. The R package *MatrixCorrelation* (Indahl et al. 2018) was utilized to analyse the similarity

between the Markov chain transitional matrices for the three main locations. Three different types of related matrix correlation coefficients were compared: RV (Robert and Escoufier 1976), RV2 (Smilde et al. 2009) and adjusted RV (Mayer et al. 2011). Three coefficients were used to avoid potential bias using only one type of correlation. All statistical analyses were employed and analysed in the statistical software R (R Core Team, 2020; R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available online at <https://www.R-project.org/>).

3 Results

3.1 Main population dynamics

The main populations at St Johns had 317 and 355 tillers in 2017 and 2018, respectively, while Lakeside had 535 (Table 1). The flowering started in late May (May 27 ± 2 days) in both years at St Johns and Lakeside (Fig. 2). During the first few days (May 29 ± 2 days), 10–20% of the tillers in the population reached the flowering phases p1, p2 and p3; hence, each of these may have up to 75% of their anther extruded. Most of these, augmented by non-flowering, tillers reached the full flowering in p4 during the first few days of June (June 1 ± 1 day). At this point, about 40% of the population was in full flowering (p4) and two days later an additional 20% of the population reached full flowering (p4) (June 3 ± 1 days). Most tillers in lower flowering phases progressed into full flowering over the coming few days (June 6 ± 2 days). At this time, most tillers were either in full flowering (p4) or had not yet started (p0). This continued until the middle of June (June 12 ± 3 days) when the first tillers reached senescence (p5), showcased by the detachment of the last anther. Around the same time (June 12 ± 1 day), the populations reached peak flowering. Peak flowering is the event where the maximum number of tillers in the season occupies the phase full flowering (p4). The phenological progression of the populations continued as an increasing number of tillers reached senescence and where the last tillers took considerable time to reach phase p5, illustrated by the long tail in Fig. 2. This progression continued until the last tiller had reached senescence,

which happened on July 17, August 9 and July 24 for each main population, respectively (Fig. 2).

3.2 Secondary population dynamics

The six secondary populations contained between 160 and 427 tillers (Table 1). The populations started flowering before the beginning of June (June 1–3–4 days) (Fig. 3). All of them had reached more than 60% full flowering (p4) in early June (June 7). By the middle of June (June 19), all populations had started senescence with at least some tillers in the senescence phase (p5). To check for spatial differentiation between the populations, phase ratios were compared. There was no discernible spatial differentiation found between the most western/eastern or northern/southern locations.

3.3 General flowering progression

The development of the 2017 population was mostly two days ahead of the 2018 population, with the full flowering phase being reached with only one day difference (Fig. 4). The main difference between the years was the amount of time needed to reach senescence: it was reached 20 days later in the 2018 population. In terms of development between the two years, the 2017 population was on average one-fifth of one phase earlier than the 2018 population. For example, if on day X the 2018 population was average phase 3.5, then the 2017 population was average phase 3.7. The standard error in the 2018 population ranged between 0.70 and 0.22 in the beginning of the season (June 1 to June 6), never reaching more than 0.15 for the rest of the season and reducing to almost zero towards the late season. See supplementary data (Fig. S7) for the contribution of each population to the general progression of 2018. A nonparametric correlation was used to compare the years due to both general populations expressing non-normality measured in the Shapiro test ($p < 0.001$ for both years). The Spearman rho correlation showed that there was a high correlation between the two years (Spearman $r_s = 0.98$).

3.4 Tiller-specific phase dynamics

The Markov chain diagrams (Fig. 5a and Table S1 for the transitional matrices) show that the 2017 St Johns

Fig. 2 Flowering phenology development of *Dactylis glomerata* populations shown per phase for the three main populations and years

Fig. 3 Flowering phenology development of *Dactylis glomerata* populations shown per phase for the secondary populations in 2018

Fig. 4 General *Dactylis glomerata* flowering progression of all observed populations for both years

population flowers first transition either to low flowering (p1) or directly to full flowering (p4), with chances of 0.41 and 0.28, respectively. The second most likely transition for all lower flowering phases (p1, p2 and p3) was to full flowering (p4), with a 0.20

to 0.25 chance for p1 and p2 but almost a 0.50 chance for p3. Tillers in p4 had only a 0.05 chance of transitioning to senescence (p5). Also note that all possible transitions were present in this population, even the less prevalent ones such as p2 to p5, with only

Fig. 5 Tiller-specific phase dynamics for the main populations and years. Arrows indicate the direction of the transition, with all arrows from each phase adding up to 1. The thickness of each arrow is directly associated with the strength of the transition

a 0.006 chance, where tillers never reached full flowering before they reached senescence. The 2018 St Johns population tillers had about a 0.73 to 0.79 chance of going into full flowering (p4) from each of the three phases p1, p2 and p3 (**Fig. 5b**). Most of the remaining chance was to stay in their respective phases. The exception was pre-flowering (p0), which had a 0.10 chance to transition into p1. Tillers in p1, p3 and p4 had chances of 0.02, 0.03 and 0.06 chance of transitioning into senescence (p5). One of the possible transitions was not observed. In the 2018 Lakeside population, tillers were most likely to transition into full flowering (p4), with chances ranging from 0.64 to 0.76 (**Fig. 5c**). The second most likely transition for the lower flowering phases was to stay in their respective phase, with chances of 0.225, 0.243 and 0.263 for p1, p2 and p3, respectively. The senescence rate for p4 into p5 was 0.074. Four of the possible transitions were not observed. The RV correlation coefficients for all the population comparisons ranged from 0.824 to 0.999 (**Table 3**). The RV coefficients ranged between 0.995 and 0.999, 0.824 and 0.932 and 0.847 and 0.950 for the spatial, temporal and spatiotemporal correlations, respectively.

4 Discussion

4.1 Demographic flowering progression

The phenological progressions of the *Dactylis glomerata* populations illustrate the same general pattern of development. The general flowering progression shows that populations in a larger area will display uniformity in average phenological progression between years. The phase distributions show small differences in day-to-day variation between all populations. Key phenological events were clearly distinguished, with start of flowering, peak flowering and

start of senescence all overlapping, in both time and space. This supports previous findings by León-Ruiz et al. (2011) that *Dactylis* flowering development is synchronistic between populations over a larger region and between years. The only main difference between populations was the senescence rates, likely attributed to local conditions responsible for drying (Atzema 1992; Hayhoe and Jackson 1974). In this study, full flowering of the first 60% of the tillers was reached within roughly 14 days, while the remaining 40% took more than a month. This demonstrates that the flowering of a population follows a non-normal distribution. We also found that tillers that start flowering later are more likely to flower for a shorter time (results not shown), as previously demonstrated by Tormo-Molina et al. (2015). The synchronal phenological progression of all populations indicates that *Dactylis* pollen release is likely to happen simultaneously over a larger area. This is supported by findings from Italy (Ghitarrini et al. 2017), Spain (Cebrino et al. 2016) and Austria (Kmenta et al. 2017). If pollen release will be proportional to the percent of extruded anthers in each tiller, then higher flowering phases will likely release more pollen (Tormo et al. 2011) and that the largest amount of pollen will be released faster than the remaining 40%. This complements a previous study by Comtois (2000), suggesting that aerobiological studies should utilize gamma distributions instead of a normal distribution when conducting a hypothesis test. Furthermore, our results illustrate that average full flowering in a population follows an uneven distribution with a mixture of some non-flowering (p0) tillers, most in full flowering (p4) and some senescent (p5). This highlights a distinct difference between peak flowering and full flowering, with peak flowering of the season being the day of maximum numbers of tillers in full flowering. Therefore, peak flowering of populations will likely be more connected with pollen release, than the average full

Table 3 RV coefficients between the three main seasonal Markov chain population matrices calculated from the flowering phenological observations in *Dactylis glomerata*

Matrix 1	Matrix 2	Correlation tested	RV coefficient	RV2 coefficient	RVadj coefficient
St Johns 2018	Lakeside 2018	Spatial	0.999	0.995	0.999
St Johns 2017	St Johns 2018	Temporal	0.849	0.932	0.824
St Johns 2017	Lakeside 2018	Temporal and spatial	0.868	0.950	0.847

flowering. If pollen has the potential to disperse evenly during the flowering progression, then peak flowering will also explain maximum pollen release potential (Frenguelli et al. 2010; Romero-Morte et al. 2018). This study therefore suggests that the daily or bi-daily resolution of *Dactylis* phenology is needed to illustrate the narrow window of maximum pollen release potential that occurs during the peak flowering of *Dactylis glomerata* populations. This information can increase the certainty of species pollen release estimates, which can be of great importance in dispersion modelling, forecasting and the usage of regional pollen calendars (e.g. Adams-Groom et al. 2020; Lo et al. 2019).

4.2 Stochastic tiller dynamics

Our findings illustrate that each population is a heterogeneous distribution of tillers occupying varying phases throughout the flowering development. The population matrix correlations are high between populations, which suggests an inherently uniform developmental dynamics shared between populations. However, the correlations also indicate that the spatial correlation is stronger than the temporal and spatiotemporal correlations in the developmental stochasticity between populations. This suggests that differences are larger between years than between separate locations, probably due to regional phenomena that vary on a temporal basis.

Although no other studies have been performed on the phase-based developmental dynamics in *Dactylis* (or any other grasses for that matter), at least two connected studies (Calder 1964a, b) propose that stage-based development is the result of physiological processes caused by photoperiod and temperature. Three stages were proposed: juvenile stage (irresponsive to photoperiod and temperature), inductive stage (responsive to photoperiod and temperature) and post-inductive stage (responsive to long days). Calder (1964a) further states the continuation of studies into *Dactylis* stage-based differentiation is essential in determining the full extent of the flowering process. Later studies confirm the differentiated physiological response to photoperiod and temperature (Broué and Nicholls 1973; Heide 1987). The likelihood of tillers to stay in lower flowering phases (as seen in 2017) is consistent with a lack of external growth stimuli to progress to full flowering (Calder 1963; Heide 1994).

On the other hand, tillers expressed high likelihoods of transitioning directly from pre-flowering (p0) to full flowering (p4) (as seen in 2018). This complements studies from Spain as it confirms *Dactylis* tillers' capacity to quickly reach full pollen release potential from one day to the next (Cebrino et al. 2016). Such rapid responses might be a contributing factor to the large daily variation in grass pollen concentrations that have previously been observed within the urban environment (Hjort et al. 2016; Simoleit et al. 2017; Werchan et al. 2017). The observed small variation in the stochastic element implies the presence of a consistent developmental progression within the species, likely to be seen in other species using the same approach. Although this study has not included a meteorological component, we suggest that small variations in meteorological factors such as solar radiation or temperature would likely explain the differential in tiller-specific flowering dynamics between 2017 and 2018. This has previously been shown using Markov chain approaches to be the case for yearly variation in Birch masting behaviour (Tseng et al. 2020). The novel aspect of including stochasticity in flowering dynamics demonstrates that grass populations have defined probabilistic developmental patterns that can be applied to populations over a larger region and potentially utilized to investigate pollen release scenarios. In contrast, the behaviour of small populations or individual plants can be difficult to predict due to the nature of probability.

4.3 Connection to grass pollen concentrations

The findings in this study increase knowledge on airborne grass pollen concentrations considerably and may have direct application in the development and improvement of numerical forecasting models for grass pollen using regional scale atmospheric transport models. In Europe, models have been developed for trees (e.g. Kurganskiy et al. 2020; Sofiev et al. 2015; Zhang et al. 2014; Zink et al. 2013) and ragweed (e.g. Prank et al. 2013; Zink et al. 2017), partly due to availability of large-scale source maps (Pauling et al. 2012; Skjøth et al. 2019) and their tendency to long distance transport (de Weger et al. 2016; Šikoparija et al. 2013; Siljamo et al. 2008; Skjøth et al. 2007). This makes regional-scale models particularly useful for simulating pollen concentrations for these species. For grasses, this may be different. Firstly, pollen

consists of a range of species all contributing to the overall pollen concentration (Ghitarrini et al. 2017). This can make current emission models developed for trees difficult to implement, depending on the local characteristics. In Australia, first attempts have been made with the CSIRO Chemical Transport Model, but with limited success (C-CTM) using mechanistic emission modelling (Emmerson et al. 2019). In the UK, the different grass species have been found to flower in a non-unified pattern throughout the country (Brennan et al. 2019), suggesting that flowering of individual species responds to local environmental variables. A recent study by Rowney et al. (2021) detecting *D. glomerata* and other grass species within the UK network found that 83% of the catch of *D. glomerata* in 2017 in Worcester was captured during the period 4–21 June. This corresponded to the period dominated by p4 and partly p5. In addition, on days before 4 June, when no or limited amounts of *D. glomerata* pollen were present in Worcester, the period was dominated by lower flowering phases (p1–p3). During this period, larger amounts of *D. glomerata* pollen were found at other sites in the UK (Rowney et al. 2021). This suggests that the first part of the flowering sequence (p1–p3) contributes only small amounts of pollen to the airborne pollen concentration. It is likely that other grass species will show a similar behaviour. It also suggests limited or no long-distance transport of grass pollen to Worcester of *D. glomerata* outside the local flowering period. This complements previous findings that grass pollen concentrations are much more localised phenomena compared to trees and ragweed with significant variations at the urban scale (e.g. Hugg et al. 2017; Skjth et al. 2013). It has been suggested that such variations are caused by variations in emission patterns between different grass species (Peel et al. 2014) and localised emission sources (e.g. Skjth et al. 2013). The Markov chain modelling approach can take this stochastic emission pattern into account, including potential rapid changes in emission potential. This concept is currently not implemented in state-of-the-art emission models designed for regional scale models (e.g. Zhang et al. 2014; Zink et al. 2013) and conceptually introduces a numerical approach to quantify the element of chance (or risk) of high pollen emission. This approach may be needed for fine-scale modelling of grass pollen concentrations at the urban scale.

5 Conclusion

This study demonstrates that the flowering progression of *Dactylis glomerata* populations follows a non-normal distribution, with a skewness towards the beginning of the season. The study shows that several populations are synchronal within an entire region and between years, with key phenological events only differing by a few days. Populations share similar phase distributions between timesteps, highlighting the uniformity of development dynamics throughout each season. Tiller-specific flowering development revealed that stochastic elements are critical in explaining inherent population demographic properties of the flowering process within grass species and that rapid development of the entire population can happen from day to day. The development may be important in forecasting. Markov chain modelling approaches utilized in the study can be further combined with local grass maps and meteorology to develop grass pollen emission models for use by numerical models.

Acknowledgements Thanks are due to Dr Geoffrey Petch for providing assistance during the fieldwork. This research was funded by the European Commission through a Marie Curie Career Integration Grant (Project ID CIG631745 and Acronym SUPREME to CAS) and by the University of Worcester.

Author contributions CAF was involved in conceptualization, methodology, validation, formal analysis, investigation, resources, data curation and writing—original draft. CAF, BAG and CAS were involved in writing—review and editing and visualization. CAS was involved in supervision, project administration and funding acquisition.

Funding This research was funded by the European Commission through a Marie Curie Career Integration Grant (Project ID CIG631745 and Acronym SUPREME to CAS) and by the University of Worcester.

Compliance with ethical standards

Conflicts of interests The authors declare that they have no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not

included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Aboulaich, N., Bouziane, H., Kadiri, M., Del Mar Trigo, M., Riadi, H., Kazzaz, M., & Merzouki, A. (2009). Pollen production in anemophilous species of the Poaceae family in Tetouan (NW Morocco). *Aerobiologia*, 25(1), 27–38. <https://doi.org/10.1007/s10453-008-9106-2>.
- Adams-Groom, B., Skjøth, C. A., Selby, K., Pashley, C., Satchwell, J., Head, K., & Ramsay, G. (2020). Regional calendars and seasonal statistics for the United Kingdom's main pollen allergens. *Allergy: European Journal of Allergy and Clinical Immunology*, 75(6), 1492–1494. <https://doi.org/10.1111/all.14168>.
- Atzema, A. J. (1992). A model for the drying of grass with realtime weather data. *Journal of Agricultural Engineering Research*, 53(C), 231–247. [https://doi.org/10.1016/0021-8634\(92\)80085-7](https://doi.org/10.1016/0021-8634(92)80085-7).
- Balzer, H. (2000). Markov chain models for vegetation dynamics. *Ecological Modelling*, 126(2–3), 139–154. [https://doi.org/10.1016/S0304-3800\(00\)00262-3](https://doi.org/10.1016/S0304-3800(00)00262-3).
- Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., & Forister, M. L. (2003). The ecology of individuals: Incidence and implications of individual specialization. *American Naturalist*, 161(1), 1–28. <https://doi.org/10.1086/343878>.
- Brennan, G. L., Potter, C., de Vere, N., Griffith, G. W., Skjøth, C. A., Osborne, N. J., et al. (2019). Temperate airborne grass pollen defined by spatio-temporal shifts in community composition. *Nature Ecology and Evolution*, 3(5), 750–754. <https://doi.org/10.1038/s41559-019-0849-7>.
- Brock, J. L., Hume, D. E., & Fletcher, R. H. (1996). Seasonal variation in the morphology of perennial ryegrass (*Lolium perenne*) and cocksfoot (*Dactylis glomerata*) plants and populations in pastures under intensive sheep grazing. *Journal of Agricultural Science*, 126(1), 37–51. <https://doi.org/10.1017/S0021859600088791>.
- Broué, P., & Nicholls, G. H. (1973). Flowering in *Dactylis Glomerata*. II. Interaction of Temperature and Photoperiod. *Australian Journal of Agricultural Research*, 24(5), 685–692. <https://doi.org/10.1093/aob/28.2.187>.
- Calder, D. M. (1963). Environmental control of flowering in *Dactylis glomerata* L. *Nature*, 197(4870), 882–883. <https://doi.org/10.1038/197882a0>.
- Calder, D. M. (1964a). Stage development and flowering in *Dactylis glomerata* L. *Annals of Botany*, 28(2), 187–206. <https://doi.org/10.1093/aob/28.2.187>.
- Calder, D. M. (1964b). Flowering Behaviour of Populations of *Dactylis Glomerata* Under Field Conditions in Britain. *Journal of Applied Ecology*, 1(2), 307–320. <https://doi.org/10.2307/2401315>.
- Canales, J., Trevisan, M. C., Silva, J. F., & Caswell, H. (1994). A demographic study of an annual grass (*Andropogon brevifolius* Schwarz) in burnt and unburnt Savanna. *Acta Oecologia*, 15(3), 261–273.
- Cebrino, J., Galán, C., & Domínguez-Vilches, E. (2016). Aerobiological and phenological study of the main Poaceae species in Córdoba City (Spain) and the surrounding hills. *Aerobiologia*, 32(4), 595–606. <https://doi.org/10.1007/s10453-016-9434-6>.
- Cebrino, J., García-Castaño, J. L., Domínguez-Vilches, E., & Galán, C. (2018). Spatio-temporal flowering patterns in Mediterranean Poaceae. A community study in SW Spain. *International Journal of Biometeorology*, 62(4), 513–523. <https://doi.org/10.1007/s00484-017-1461-7>.
- Clayton, W. D., Vorontsova, M. S., Harman, K. T., & Williamson, H. (2002). World Grass Species: Synonymy. *Kew: GrassBase*. <http://www.kew.org/data/grasses-syn.html>. Accessed 1 July 2020
- Cleland, E. E., Chiariello, N. R., Loarie, S. R., Mooney, H. A., & Field, C. B. (2006). Diverse responses of phenology to global changes in a grassland ecosystem. *Proceedings of the National Academy of Sciences*, 103(37), 13740–13744. <https://doi.org/10.1073/pnas.0600815103>.
- Comtois, P. (2000). The gamma distribution as the true aerobiological probability density function (PDF). *Aerobiologia*, 16(2), 171–176. <https://doi.org/10.1023/A:1007667531246>.
- Cornelius, C., Petermeier, H., Estrella, N., & Menzel, A. (2011). A comparison of methods to estimate seasonal phenological development from BBCH scale recording. *International Journal of Biometeorology*, 55, 867–877. <https://doi.org/10.1007/s00484-011-0421-x>.
- Cornelius, C., Petermeier, H., Estrella, N., & Menzel, A. (2014). Erratum to: A comparison of methods to estimate seasonal phenological development from BBCH scale recording. *International Journal of Biometeorology*, 58, 1707. <https://doi.org/10.1007/s00484-014-0858-9>.
- D'Amato, G., Cecchi, L., Bonini, S., Nunes, C., Annesi-Maesano, I., Behrendt, H., et al. (2007). Allergenic pollen and pollen allergy in Europe. *Allergy: European Journal of Allergy and Clinical Immunology*, 62(9), 976–990. <https://doi.org/10.1111/j.1398-9995.2007.01393.x>.
- Davila-Velderrain, J., Martinez-Garcia, J. C., & Alvarez-Buylla, E. R. (2016). Dynamic network modelling to understand flowering transition and floral patterning. *Journal of Experimental Botany*, 67(9), 2565–2572. <https://doi.org/10.1093/jxb/erw123>.
- de Weger, L. A., Pashley, C. H., Šikoparija, B., Skjøth, C. A., Kasprzyk, I., Grewling, L., et al. (2016). The long distance transport of airborne Ambrosia pollen to the UK and the Netherlands from Central and south Europe. *International Journal of Biometeorology*, 60(12), 1829–1839. <https://doi.org/10.1007/s00484-016-1170-7>.
- Eagles, C. F. (1972). Competition for Light and Nutrients Between Natural Populations of *Dactylis glomerata*. *The Journal of Applied Ecology*, 9(1), 141–151. <https://doi.org/10.2307/2402052>.
- Eagles, C. F., & Williams, D. H. (1971). Competition between natural populations of *Dactylis glomerata*. *The Journal of Agricultural Science*, 77(2), 187–193. <https://doi.org/10.1017/S0021859600024291>.

- Emmerson, K. M., Silver, J. D., Newbiggin, E., Lampugnani, E. R., Suphioglu, C., Wain, A., & Ebert, E. (2019). Development and evaluation of pollen source methodologies for the Victorian Grass Pollen Emissions Module VGPEM1.0. *Geoscientific Model Development*, 12(6), 2195–2214. <https://doi.org/10.5194/gmd-12-2195-2019>.
- Evans, M. E. K., Merow, C., Record, S., McMahon, S. M., & Enquist, B. J. (2016). Towards Process-based Range Modeling of Many Species. *Trends in Ecology and Evolution*, 31(11), 860–871. <https://doi.org/10.1016/j.tree.2016.08.005>.
- Fox, G. A., & Kendall, B. E. (2002). Demographic stochasticity and the variance reduction effect. *Ecology*, 83(7), 1928–1934. <https://doi.org/10.2307/3071775>.
- Frenguelli, G., Passalacqua, G., Bonini, S., Fiocchi, A., Incorvaia, C., Marcucci, F., et al. (2010). Bridging allergologic and botanical knowledge in seasonal allergy: A role for phenology. *Annals of Allergy, Asthma and Immunology*, 105(3), 223–227. <https://doi.org/10.1016/j.anai.2010.06.016>.
- García-Mozo, H., Galán, C., Belmonte, J., Bermejo, D., Candau, P., Díaz de la Guardia, C., et al. (2009). Predicting the start and peak dates of the Poaceae pollen season in Spain using process-based models. *Agricultural and Forest Meteorology*, 149(2), 256–262. <https://doi.org/10.1016/j.agrformet.2008.08.013>.
- García-Mozo, H., Mestre, A., & Galán, C. (2010). Phenological trends in southern Spain: A response to climate change. *Agricultural and Forest Meteorology*, 150(4), 575–580. <https://doi.org/10.1016/j.agrformet.2010.01.023>.
- Ghitarrini, S., Galán, C., Frenguelli, G., & Tedeschini, E. (2017). Phenological analysis of grasses (Poaceae) as a support for the dissection of their pollen season in Perugia (Central Italy). *Aerobiologia*, 33(3), 339–349. <https://doi.org/10.1007/s10453-017-9473-7>.
- Grinstead, C. M., & Snell, J. L. (1997). Markov chains. In *Introduction to Probability* (Second Rev., pp. 405–470). Providence, Rhode Island: American Mathematical Society.
- Hayhoe, H. N., & Jackson, L. P. (1974). Weather Effects on Hay Drying Rates. *Canadian Journal of Plant Science*, 54(3), 479–484. <https://doi.org/10.4141/cjps74-081>.
- Heide, O. M. (1987). Photoperiodic control of flowering in *Dactylis glomerata*, a true short-long-day plant. *Physiologia Plantarum*, 70(3), 523–529. <https://doi.org/10.1111/j.1399-3054.1987.tb02853.x>.
- Heide, O. M. (1994). Control of Flowering and Reproduction in Temperate Grasses. *New Phytologist*, 128(2), 347–362. <https://doi.org/10.1111/j.1469-8137.1994.tb04019.x>.
- Heinzerling, L. M., Burbach, G. J., Edenharter, G., Bachert, C., Bindslev-Jensen, C., Bonini, S., et al. (2009). GAZLEN skin test study I: GALEN harmonization of skin prick testing: Novel sensitization patterns for inhalant allergens in Europe. *Allergy: European Journal of Allergy and Clinical Immunology*, 64(10), 1498–1506. <https://doi.org/10.1111/j.1398-9995.2009.02093.x>.
- Hjort, J., Hugg, T. T., Antikainen, H., Rusanen, J., Sofiev, M., Kukkonen, J., et al. (2016). Fine-Scale exposure to allergenic pollen in the Urban environment: Evaluation of land use regression approach. *Environmental Health Perspectives*, 124(5), 619–626. <https://doi.org/10.1289/ehp.1509761>.
- Hugg, T. T., Hjort, J., Antikainen, H., Rusanen, J., Tuokila, M., Korkonen, S., et al. (2017). Urbanity as a determinant of exposure to grass pollen in Helsinki Metropolitan area. *Finland. PLoS ONE*, 12(10), 1–17. <https://doi.org/10.1371/journal.pone.0186348>.
- Indahl, U. G., Næs, T., & Liland, K. H. (2018). A similarity index for comparing coupled matrices. *Journal of Chemometrics*, 32(10), 1–18. <https://doi.org/10.1002/cem.3049>.
- Jantunen, J., Saarinen, K., Valtonen, A., & Saarnio, S. (2007). Flowering and seed production success along roads with different mowing regimes. *Applied Vegetation Science*, 10(2), 285–292. <https://doi.org/10.1111/j.1654-109X.2007.tb00528.x>.
- Jochner, S., Ziello, C., Böck, A., Estrella, N., Buters, J., Weichenmeier, I., et al. (2012). Spatio-temporal investigation of flowering dates and pollen counts in the topographically complex Zugspitze area on the German-Austrian border. *Aerobiologia*, 28(4), 541–556. <https://doi.org/10.1007/s10453-012-9255-1>.
- Jones, M. D., & Newell, L. C. (1948). Size, Variability, and Identification of Grass Pollen. *Agronomy Journal*, 40(2), 136–143. <https://doi.org/10.2134/agronj1948.00021962004000020004x>.
- Kendall, B. E., & Fox, G. A. (2002). Variation among individuals and reduced demographic stochasticity. *Conservation Biology*, 16(1), 109–116. <https://doi.org/10.1046/j.1523-1739.2002.00036.x>.
- Khanduri, V. P. (2011). Variation in Anthesis and Pollen Production in Plants. *American-Eurasian J. Agric. & Environ. Sci.*, 11(6), 834–839.
- Kmenta, M., Bastl, K., Berger, U., Kramer, M. F., Heath, M. D., Pätsi, S., et al. (2017). The grass pollen season 2015: A proof of concept multi-approach study in three different European cities. *World Allergy Organization Journal*, 10(31), 1–12. <https://doi.org/10.1186/s40413-017-0163-2>.
- Kmenta, M., Bastl, K., Kramer, M. F., Hewings, S. J., Mwange, J., Zetter, R., & Berger, U. (2016). The grass pollen season 2014 in Vienna: A pilot study combining phenology, aerobiology and symptom data. *Science of the Total Environment*, 566–567, 1614–1620. <https://doi.org/10.1016/j.scitotenv.2016.06.059>.
- Kurganskiy, A., Skjøth, C. A., Baklanov, A., Sofiev, M., Saarto, A., Severova, E., et al. (2020). Incorporation of pollen data in source maps is vital for pollen dispersion models. *Atmospheric Chemistry and Physics*, 20(4), 2099–2121. <https://doi.org/10.5194/acp-20-2099-2020>.
- León-Ruiz, E., Alcázar, P., Domínguez-Vilches, E., & Galán, C. (2011). Study of Poaceae phenology in a Mediterranean climate. Which species contribute most to airborne pollen counts? *Aerobiologia*, 27(1), 37–50. <https://doi.org/10.1007/s10453-010-9174-y>.
- Liddle, A. M. J., Budd, C. S. J., & Hutchings, M. J. (1982). Population dynamics and neighbourhood effects in establishing swards of *Festuca rubra*. *Oikos*, 38(1), 52–59. <https://doi.org/10.2307/3544567>.
- Lindner, R., & Garcia, A. (1997). Geographic distribution and genetic resources of *Dactylis* in Galicia (northwest Spain).

- Genetic Resources and Crop Evolution*, 44, 499–507. <https://doi.org/10.1023/A:1008690831828>.
- Lo, F., Bitz, C. M., Battisti, D. S., & Hess, J. J. (2019). Pollen calendars and maps of allergenic pollen in North America. *Aerobiologia*, 35(4), 613–633. <https://doi.org/10.1007/s10453-019-09601-2>.
- Mayer, C. D., Lorent, J., & Horgan, G. W. (2011). Exploratory analysis of multiple omics datasets using the adjusted RV coefficient. *Statistical Applications in Genetics and Molecular Biology*, 10(1), 1–27. <https://doi.org/10.2202/1544-6115.1540>.
- Meier, U., Bleiholder, H., Buhr, L., Feller, C., Hack, H., Heß, M., et al. (2009). The BBCH system to coding the phenological growth stages of plants-history and publications. *Journal für Kulturpflanzen*, 61(2), 41–52. <https://doi.org/10.5073/JfK.2009.02.01>.
- Meier, U. (2018). *Growth stages of mono- and dicotyledonous plants. BBCH Monograph*. Quedlinburg, Germany: Julius Kühn-Institute (JKI). <https://doi.org/10.5073/20180906-074619>.
- Meyn, S. P., & Tweedie, R. L. (1993). *Markov Chains and Stochastic Stability* (1st ed.). London: Springer-Verlag, London.
- Mizianty, M. (1986). Biosystematic studies on *Dactylis L. 1*. Review of the previous studies 1.1. Systematics, variability, ecology, biology and cultivation problems. *Acta Societatis Botanicorum Poloniae*, 55(3), 467–479. <https://doi.org/10.5586/asbp.1986.039>.
- Mizianty, M. (1990). Biosystematic studies on *Dactylis L. 1*. Review of the previous studies 1.2. Cytology, genetics, experimental studies and evolution. *Acta Societatis Botanicorum Poloniae*, 59(1), 105–118. <https://doi.org/10.5586/asbp.1990.011>.
- Munson, S. M., & Long, A. L. (2017). Climate drives shifts in grass reproductive phenology across the western USA. *New Phytologist*, 213(4), 1945–1955. <https://doi.org/10.1111/nph.14327>.
- Nakaoka, M. (1996). Dynamics of age- and size-structured populations in fluctuating environments: applications of stochastic matrix models to natural populations. *Researches on Population Ecology*, 38(2), 141–152. <https://doi.org/10.1007/BF02515722>.
- Pauling, A., Rotach, M. W., Gehrig, R., Clot, B., Jäger, S., Cerny, M., et al. (2012). A method to derive vegetation distribution maps for pollen dispersion models using birch as an example. *International Journal of Biometeorology*, 56(5), 949–958. <https://doi.org/10.1007/s00484-011-0505-7>.
- Peel, R. G., Ørby, P. V., Skjøth, C. A., Kennedy, R., Schlünssen, V., Smith, M., et al. (2014). Seasonal variation in diurnal atmospheric grass pollen concentration profiles. *Biogeosciences*, 11(3), 821–832. <https://doi.org/10.5194/bg-11-821-2014>.
- Prank, M., Chapman, D. S., Bullock, J. M., Belmonte, J., Berger, U., Dahl, A., et al. (2013). An operational model for forecasting ragweed pollen release and dispersion in Europe. *Agricultural and Forest Meteorology*, 182–183, 43–53. <https://doi.org/10.1016/j.agrformet.2013.08.003>.
- Prieto-Baena, J. C., Hidalgo, P. J., Domínguez, E., & Galán, C. (2003). Pollen production in the Poaceae family. *Grana*, 42(3), 153–160. <https://doi.org/10.1080/00173130310011810>.
- Robert, P., & Escoufier, Y. (1976). A Unifying Tool for Linear Multivariate Statistical Methods: The RV-Coefficient. *Applied Statistics*, 25(3), 257–265. <https://doi.org/10.2307/2347233>.
- Rojo, J., Rivero, R., Romero-Morte, J., Fernández-González, F., & Pérez-Badia, R. (2017). Modeling pollen time series using seasonal-trend decomposition procedure based on LOESS smoothing. *International Journal of Biometeorology*, 61(2), 335–348. <https://doi.org/10.1007/s00484-016-1215-y>.
- Romero-Morte, J., Rojo, J., & Pérez-Badia, R. (2020). Meteorological factors driving airborne grass pollen concentration in central Iberian Peninsula. *Aerobiologia*, 36, 527–540. <https://doi.org/10.1007/s10453-020-09647-7>.
- Romero-Morte, J., Rojo, J., Rivero, R., Fernández-González, F., & Pérez-Badia, R. (2018). Standardised index for measuring atmospheric grass-pollen emission. *Science of the Total Environment*, 612, 180–191. <https://doi.org/10.1016/j.scitotenv.2017.08.139>.
- Rossignol, N., Andueza, D., Carrère, P., Cruz, P., Duru, M., Fiorelli, J. L., et al. (2014). Assessing population maturity of three perennial grass species: Influence of phenology and tiller demography along latitudinal and altitudinal gradients. *Grass and Forage Science*, 69(3), 534–548. <https://doi.org/10.1111/gfs.12067>.
- Rowney, F. M., Brennan, G. L., Skjøth, C. A., Griffith, G. W., McInnes, R. N., Clewlow, et al. (2021). Environmental DNA reveals links between abundance and composition of airborne grass pollen and respiratory health. [In press].
- Shapiro, S. S., & Wilk, M. B. (1965). An Analysis of Variance Test for Normality (Complete Samples). *Biometrika*, 52(3), 591–611. <https://doi.org/10.2307/2333709>.
- Šikoparija, B., Skjøth, C. A., Alm Kübler, K., Dahl, A., Sommer, J., Grewling, L., et al. (2013). A mechanism for long distance transport of Ambrosia pollen from the Pannonian Plain. *Agricultural and Forest Meteorology*, 180, 112–117. <https://doi.org/10.1016/j.agrformet.2013.05.014>.
- Siljamo, P., Sofiev, M., Ranta, H., Linkosalo, T., Kubin, E., Ahas, R., et al. (2008). Representativeness of point-wise phenological Betula data collected in different parts of Europe. *Global Ecology and Biogeography*, 17(4), 489–502. <https://doi.org/10.1111/j.1466-8238.2008.00383.x>.
- Silva, J. F., Raventos, J., Caswell, H., & Trevisan, M. C. (1991). Population Responses to Fire in a Tropical Savanna Grass, *Andropogon semiberbis*: A Matrix Model Approach. *The Journal of Ecology*, 79(2), 345. <https://doi.org/10.2307/2260717>.
- Simoleit, A., Werchan, M., Werchan, B., Mücke, H.-G., Gauger, U., Zuberbier, T., & Bergmann, K.-C. (2017). Birch, grass, and mugwort pollen concentrations and intradiurnal patterns at two different urban sites in Berlin. *Germany. Allergo Journal International*, 26(5), 155–164. <https://doi.org/10.1007/s40629-017-0012-4>.
- Skjøth, C. A., Ørby, P. V., Becker, T., Geels, C., Schlünssen, V., Sigsgaard, T., et al. (2013). Identifying urban sources as cause of elevated grass pollen concentrations using GIS and remote sensing. *Biogeosciences*, 10(1), 541–554. <https://doi.org/10.5194/bg-10-541-2013>.

- Skjøth, C. A., Sommer, J., Stach, A., Smith, M., & Brandt, J. (2007). The long-range transport of birch (*Betula*) pollen from Poland and Germany causes significant pre-season concentrations in Denmark. *Clinical and Experimental Allergy*, 37(8), 1204–1212. <https://doi.org/10.1111/j.1365-2222.2007.02771.x>.
- Skjøth, C. A., Sun, Y., Karrer, G., Sikoparija, B., Smith, M., Schaffner, U., & Müller-Schärer, H. (2019). Predicting abundances of invasive ragweed across Europe using a “top-down” approach. *Science of the Total Environment*, 686, 212–222. <https://doi.org/10.1016/j.scitotenv.2019.05.215>.
- Smilde, A. K., Kiers, H. A. L., Bijlsma, S., Rubingh, C. M., & Van Erk, M. J. (2009). Matrix correlations for high-dimensional data: The modified RV-coefficient. *Bioinformatics*, 25(3), 401–405. <https://doi.org/10.1093/bioinformatics/btn634>.
- Smith, D. C. (1944). Pollination and seeds formation in grasses. *Journal of Agricultural Research*, 68(2), 79–95.
- Sofiev, M., Berger, U., Prank, M., Vira, J., Arteta, J., Belmonte, J., et al. (2015). MACC regional multi-model ensemble simulations of birch pollen dispersion in Europe. *Atmospheric Chemistry and Physics*, 15(14), 8115–8130. <https://doi.org/10.5194/acp-15-8115-2015>.
- Soons, M. B., Heil, G. W., Nathan, R., & Katul, G. G. (2004). Determinants of long-distance seed dispersal by wind in grasslands. *Ecology*, 85(11), 3056–3068. <https://doi.org/10.1890/03-0522>.
- Spearman, C. (1904). The Proof and Measurement of Association Between Two Things. *The American Journal of Psychology* 15(1), 72–101. <https://archive.org/details/jstor-4576614>
- Spedicato, G. A. (2017). Discrete time Markov chains with R. *The R Journal*, 9(2), 84–104. <https://doi.org/10.32614/RJ-2017-036>.
- Tormo-Molina, R., Maya-Manzano, J. M., Silva-Palacios, I., Fernández-Rodríguez, S., & Gonzalo-Garijo, Á. (2015). Flower production and phenology in *Dactylis glomerata*. *Aerobiologia*, 31(4), 469–479. <https://doi.org/10.1007/s10453-015-9381-7>.
- Tormo, R., Silva, I., Gonzalo, Á., Moreno, A., Pérez, R., & Fernández, S. (2011). Phenological records as a complement to aerobiological data. *International Journal of Biometeorology*, 55(1), 51–65. <https://doi.org/10.1007/s00484-010-0308-2>.
- Tseng, Y. T., Kawashima, S., Kobayashi, S., Takeuchi, S., & Nakamura, K. (2020). Forecasting the seasonal pollen index by using a hidden Markov model combining meteorological and biological factors. *Science of the Total Environment*, 698, 1–10. <https://doi.org/10.1016/j.scitotenv.2019.134246>.
- Tufto, J., Engen, S., & Hindar, K. (1997). Stochastic dispersal processes in plant populations. *Theoretical Population Biology*, 52(1), 16–26. <https://doi.org/10.1006/tpbi.1997.1306>.
- Watkinson, A. R., & Ormerod, S. J. (2001). Grasslands, grazing and biodiversity: Editors’ introduction. *Journal of Applied Ecology*, 38(2), 233–237. <https://doi.org/10.1046/j.1365-2664.2001.00621.x>.
- Werchan, B., Werchan, M., Mücke, H. G., Gauger, U., Simoleit, A., Zuberbier, T., & Bergmann, K. C. (2017). Spatial distribution of allergenic pollen through a large metropolitan area. *Environmental Monitoring and Assessment*, 189, 1–19. <https://doi.org/10.1007/s10661-017-5876-8>.
- Wolfe, T. K. (1925). Observations on the blooming of orchard grass flowers. *Agronomy Journal*, 17(10), 605–618. <https://doi.org/10.2134/agronj1925.00021962001700100004x>.
- Wolfe, T. K., & Kipps, M. S. (1925). Pollination studies with orchard grass. *Agronomy Journal*, 17(11), 748–752. <https://doi.org/10.2134/agronj1925.00021962001700110014x>.
- Wolfe, T. K., & Kipps, M. S. (1926). Further studies of the pollination of orchard grass. *Agronomy Journal*, 18(12), 1121–1127. <https://doi.org/10.2134/agronj1926.00021962001800120012x>.
- Yan, D., Zhao, X., Cheng, Y., Ma, X., Huang, L., & Zhang, X. (2016). Phylogenetic and diversity analysis of *Dactylis glomerata* subspecies using SSR and IT-ISJ Markers. *Molecules*, 21(12), 1–13. <https://doi.org/10.3390/molecules21111459>.
- Zhang, R., Duhl, T., Salam, M. T., House, J. M., Flagan, R. C., Avol, E. L., et al. (2014). Development of a regional-scale pollen emission and transport modeling framework for investigating the impact of climate change on allergic airway disease. *Biogeosciences*, 11(6), 1461–1478. <https://doi.org/10.5194/bg-11-1461-2014>.
- Zink, K., Pauling, A., Rotach, M. W., Vogel, H., Kaufmann, P., & Clot, B. (2013). EMPOL 1.0: a new parameterization of pollen emission in numerical weather prediction models. *Geoscientific Model Development Discussions*, 6(2), 3137–3178. <https://doi.org/10.5194/gmdd-6-3137-2013>.
- Zink, K., Kaufmann, P., Petitpierre, B., Broennimann, O., Guisan, A., Gentilini, E., & Rotach, M. W. (2017). Numerical ragweed pollen forecasts using different source maps: a comparison for France. *International Journal of Biometeorology*, 61(1), 23–33. <https://doi.org/10.1007/s00484-016-1188-x>.