

# Endothelium-derived fibronectin regulates neonatal vascular morphogenesis in an autocrine fashion

Christopher J. Turner<sup>1,2</sup> · Kwabena Badu-Nkansah<sup>1,3</sup> · Richard O. Hynes<sup>1</sup>

Received: 15 February 2017 / Accepted: 21 June 2017 / Published online: 30 June 2017  
© The Author(s) 2017. This article is an open access publication

**Abstract** Fibronectin containing alternatively spliced EIIIA and EIIIB domains is largely absent from mature quiescent vessels in adults, but is highly expressed around blood vessels during developmental and pathological angiogenesis. The precise functions of fibronectin and its splice variants during developmental angiogenesis however remain unclear due to the presence of cardiac, somitic, mesodermal and neural defects in existing global fibronectin KO mouse models. Using a rare family of surviving EIIIA EIIIB double KO mice, as well as inducible endothelial-specific fibronectin-deficient mutant mice, we show that vascular development in the neonatal retina is regulated in an autocrine manner by endothelium-derived fibronectin, and requires both EIIIA and EIIIB domains and the RGD-binding  $\alpha 5$  and  $\alpha v$  integrins for its function. Exogenous sources of fibronectin do not fully substitute for the autocrine function of endothelial fibronectin, demonstrating that fibronectins from different sources contribute differentially to specific aspects of angiogenesis.

**Keywords** Fibronectin · Angiogenesis · Integrins · EIIIA · EIIIB · Autocrine

## Introduction

In addition to providing structural strength and elasticity to blood vessels, the extracellular matrix (ECM) provides instructional signals that control the development, patterning, and stability of the vasculature [1]. The ECM achieves this, in part, by binding and regulating the distribution and activity of growth factors such as the vascular endothelial growth factors (VEGFs), platelet-derived growth factor (PDGFs), fibroblast growth factor (FGFs) and transforming growth factor- $\beta$  (TGF- $\beta$ ) [2]. The ECM also regulates vascular development by directly binding and conveying both biochemical and biomechanical signals through integrin receptors [3]. Integrins comprise a family of heterodimeric adhesion receptors that contains 16 $\alpha$  and 8 $\beta$  subunits that associate to form 24 different receptors that bind to the ECM with distinct yet often overlapping specificities. Previous studies have shown that the interaction of integrins with the ECM is essential for endothelial cell (EC) adhesion and, as a consequence, regulates EC proliferation, migration, and the sprouting of new vessels [4]. Indeed, concentration gradients of immobilised ECM proteins have been shown to control both the direction and speed of EC migration in the absence of chemokines *in vitro* [5, 6]. Furthermore, ECM-integrin interactions have been shown to regulate EC junction formation, vessel stability and integrity [7], cell polarity and vessel lumen formation [8].

One of the most extensively studied ECM proteins involved in vascular development is the glycoprotein fibronectin (Fn) [9]. Fibronectin is a modular protein

**Electronic supplementary material** The online version of this article (doi:10.1007/s10456-017-9563-8) contains supplementary material, which is available to authorized users.

✉ Richard O. Hynes  
rohynes@mit.edu

<sup>1</sup> Howard Hughes Medical Institute, Koch Institute for Integrative Cancer Research, Massachusetts Institute of Technology, 77 Massachusetts Ave, 76-361, Cambridge, MA 02139, USA

<sup>2</sup> Present Address: University of Suffolk, James Hehir Building, University Avenue, Ipswich, Suffolk IP3 0FS, UK

<sup>3</sup> Present Address: Duke University Medical Center, 307 Research Drive, Durham, NC 27710, USA

consisting of type I, II, and III repeating units and is alternatively spliced to exclude extra EIIIA and EIIIB domains, and portions of the variable (V or IIICS) domain. Fibronectin is found only in vertebrates with an endothelium-lined vasculature [10] and is expressed around early embryonic vessels before the presence of other basement membrane or structural ECM proteins [11, 12]. Interestingly, Fn containing EIIIA and EIIIB domains is highly expressed around angiogenic vessels [13, 14], but is largely absent from mature quiescent vessels in adults until subjected to injury or low or disturbed flow [15, 16], suggesting that both domains are required for the development and remodelling of the vasculature.

The functional importance of fibronectin can be seen from several genetic studies [9]. Global deletion of *Fn* leads to early embryonic lethality due to severe neural, mesodermal, cardiac and vascular defects in mice [12, 17, 18]. Individually EIIIA-null and EIIIB-null mice are viable, fertile, and lack reported defects in either developmental or tumour angiogenesis [19, 20], however mice lacking both EIIIA and EIIIB domains die around E10.5 with multiple developmental and cardiovascular defects [21]. Interestingly, just as observed in *Fn*-null mice and zebrafish (*natter1* mutants), the severity of the defects in EIIIA EIIIB double KO mice varies with genetic background, suggesting the presence of a genetic modifier(s) [21–23]. The RGD motif within Fn, recognised by  $\alpha 5\beta 1$  and  $\alpha v$  integrins (i.e.  $\alpha v\beta 1$ ,  $\alpha v\beta 3$ ,  $\alpha v\beta 5$ ,  $\alpha v\beta 6$  and  $\alpha v\beta 8$ ), is also essential for embryonic development. Replacement of the RGD by an inactive RGE motif leads to neural, somitic and cardiovascular defects [17, 24], while mice lacking endothelial expression of  $\alpha 5$  and  $\alpha v$  integrins, the major RGD-binding receptors expressed on ECs, die around E14.5 with defects in the development of the heart, great vessels and lymphatic vasculature [25, 26].

More recently, Fn secreted and assembled ahead of the developing vasculature by astrocytes has been shown to guide EC migration by binding and presenting VEGF to EC tip cells [27], while Fn, including contributions from plasma, has been shown to promote tumour angiogenesis through increased retention of VEGF in the tumour environment [28]. The precise role of endothelial-derived Fn in developmental angiogenesis however remains unclear. Fn-depleted ECs display defects in angiogenic assays in vitro [29], however postnatal deletion of endothelial *Fn* fails to inhibit tumour angiogenesis in mice [30].

To investigate further the role of fibronectin and its alternatively spliced EIIIA and EIIIB domains in developmental angiogenesis in the absence of heart, great vessel, somite, or neural crest defects, we have analysed in detail the postnatal growth and patterning of blood vessels within the retinas of inducible global *Fn* KO mice, EIIIA and EIIIB double KO mice, and inducible endothelial-specific

*Fn* mutants. Our results show that vessel outgrowth, branching, sprouting and stability are regulated to a significant degree by EC-derived EIIIA<sup>+</sup>/EIIIB<sup>+</sup>-containing fibronectin and require endothelial expression of either  $\alpha 5\beta 1$  or the  $\alpha v$  integrins.

## Methods

### Mouse lines

All mice were housed and handled in accordance with approved Massachusetts Institute of Technology Division of Comparative Medicine protocols (IACUC approval 0412-033-15). *Fn* floxed [31], *Itga5* floxed [26], *Itgav* floxed [32], Rosa26-CreER<sup>T2</sup> [33], Cdh5(PAC)-CreER<sup>T2</sup> [34], and mTmG [35], mouse lines have all been described previously. Live EIIIA/EIIIB KO mice were established by extensively backcrossing EIIIA/EIIIB heterozygous mice [21] to C57BL/6 J wild-type mice and inter-crossing the resulting mice at  $n = 10$ . Cre activity and gene deletion were induced in *Fn*<sup>iKO</sup> and *Fn*<sup>iEC KO</sup> mice through consecutive intraperitoneal injections of 50 mg tamoxifen (Sigma) at P1, P2 and P3. Genotyping was performed on DNA isolated from tail snips in-house or by Transnetyx.

### Immunofluorescence staining

Whole-mount immunofluorescence staining of retinas was achieved following methods previously described in Pitulescu et al. [36]. Briefly, eyes were freshly isolated and fixed in 4% paraformaldehyde (PFA) in PBS at 4 °C overnight, blocked in PBS containing 0.5% Tween (PBS-T) and 2% goat, donkey or fibronectin-depleted goat serum and incubated overnight at 4 °C with primary antibodies either in 0.25% Tween/1% serum in PBS (staining buffer) or, for isolectin-B4 staining, in Pblec buffer (1% Triton X-100, 1 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub> and 1 mM MnCl<sub>2</sub> in PBS, pH 6.8). After washes in PBS, retinas were incubated either at RT for 2 h, or overnight at 4 °C, with fluorophore-conjugated secondary antibodies diluted in staining buffer. Retinas were then washed in PBS, cut into three segments and flat-mounted onto coverslips in Fluoromount (Southern Biotech).

### Antibodies

Primary antibodies: Rat anti-mouse PECAM-1 MEC13.3 (1:100, BD Pharmingen), rabbit anti-Fibronectin (1:200, Hynes lab) [16], mouse anti-EIIIA fibronectin antibody (1:100, Abcam, IST-9), rabbit anti-Collagen IV (1:400, Abcam, ab19808), rabbit anti-Desmin (1:200, Abcam, ab15200), biotinylated Isolectin-B4 (1:50, Vector Labs,

B-1205), mouse anti-EIIIB fibronectin (1:100, Amy McMahan, Hynes lab). Secondary antibodies were Alexa488, Alexa594, and Alexa647 conjugated antibodies (1:500, Invitrogen), and Alexa-Fluor–streptavidin-conjugated antibodies (1:200, Molecular Probes).

### Image acquisition and processing

All images were acquired using Zeiss LSM 510 or Nikon A1R scanning laser confocal microscopes and processed using Volocity (Perkin Elmer) or Nikon Elements software and Adobe Photoshop.

### Quantitative analyses of the retinal vasculature

All quantifications were completed using Volocity (Perkin Elmer) software using maximal intensity projection images.

Radial outgrowth/vessel migration was measured in a straight line from the optic nerve to the angiogenic front of the retinal plexus in 6 different mice from each genotype group ( $n = 6$ ). Branch points were calculated from  $250 \mu\text{m} \times 500 \mu\text{m}$  fields of view (FOV) adjacent to retinal veins behind the angiogenic front using 6 different retina samples from each genotype group ( $n = 6$ ).

Endothelial coverage, tip cell sprouts, filopodial numbers and lengths were all measured using only isolectin-B4 stained retinas. Endothelial percentage coverage is defined as the area of isolectin-B4-positive immunofluorescence divided by the total area, and was calculated from  $340 \text{mm}^2$  fields within the capillary plexus behind the angiogenic front ( $n = 6$  mice per genotype). Endothelial tip cell numbers were quantified by counting endothelial sprouts at the angiogenic front of the entire vascular plexus ( $n = 3$  retinas per genotype). Endothelial tip cell filopodial numbers and lengths were calculated from high-resolution confocal images (60X objective, thin z-sections of sample) of 12 randomly selected tip cells at the leading edge of the vascular plexus from a minimum of 3 retinas per genotype ( $n = 12$ ). Filopodia lengths were calculated from a minimum of 50 filopodia ( $n = 50$ ).

Fibronectin deposition around the vasculature was quantified by measuring the mean fibronectin immunofluorescent pixel intensity in 10 randomly selected PECAM1-positive  $500 \mu\text{m}^2$  fields within  $200 \mu\text{m}$  of the leading edge of the capillary plexus, using a minimum of 2 retinas per genotype ( $n = 10$ ). Vessel regression events were measured by counting PECAM1<sup>-</sup>/collagen IV<sup>+</sup> structures within 6 FOV (sized  $300 \mu\text{m} \times 300 \mu\text{m}$ ) adjacent to the retinal veins behind the angiogenic front using 3 retinas per genotype ( $n = 6$ ). Endothelial cell proliferation was calculated by measuring the number of Ki-67<sup>+</sup> PECAM1<sup>+</sup> cells within 2 retinas and normalising to  $100 \mu\text{m}$  vessel

length ( $n = 2$ ). Pericyte numbers were calculated from 6 fields (sized  $230 \mu\text{m} \times 230 \mu\text{m}$ ) within the capillary plexus using a minimum of 3 retinas per genotype ( $n = 6$ ).


Results are plotted as mean  $\pm$  s.e.m and were analysed using Student's *t* test and considered significant when \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$  or \*\*\*\* $P < 0.0001$ .

## Results

### Fibronectin controls the patterning of the retinal vasculature

Fibronectin is highly expressed throughout the developing retinal vasculature (Fig. 1a–f) and is assembled ahead of the vascular plexus by retinal astrocytes (Fig. 1a–c) forming a scaffold for ECs migration [27, 37]. In contrast with the fibronectin assembled by astrocytes, the majority of the fibronectin surrounding the vasculature contained the alternatively spliced EIIIA and/or EIIIB domain(s) (Fig. 1b–f), with expression of both EIIIA and EIIIB especially pronounced around the edges of transcapillary pillars (holes) in vessels undergoing intussusceptive angiogenesis (Fig. 1d–f).

To investigate the role of fibronectin in regulating the development of the retinal vasculature we first deleted *Fn* from all postnatal tissue (*Fn*<sup>iKO</sup>) by administering tamoxifen to newborn mice carrying the inducible (ubiquitously expressed) ROSA26-CreER<sup>T2</sup> transgene and a loxP-flanked *Fn* gene (Fig. S1a). Surprisingly, despite previous studies showing that fibronectin is essential for early developmental angiogenesis [12, 18, 38, 39], whole-mount immunofluorescence staining of retinas revealed that both *Fn*<sup>flox/flox</sup> mice (control) and Rosa26-CreER<sup>T2+</sup> *Fn*<sup>flox/flox</sup> mutants (*Fn*<sup>iKO</sup>) develop a hierarchical vascular tree of arteries, veins and capillaries by postnatal day 6 (P6) (Fig. 2a–d). In contrast to control mice however, radial growth and vessel branching were significantly compromised in *Fn*<sup>iKO</sup> mice (Fig. 2a–f) and, as a consequence, endothelial coverage within the retinal tissue was significantly reduced (Fig. 2a–i). Since the EIIIA and EIIIB domains of fibronectin have been shown to regulate vascular morphogenesis [21], we next analysed the retinal vasculature in a rare family of surviving C57BL/6 EIIIA<sup>-</sup> EIIIB<sup>-</sup> double-knockout mice (*Fn*<sup>AB KO</sup>). These mice express fibronectin around their vasculature at equivalent levels to control mice, however the Fn expressed no longer contains the EIIIA and EIIIB domains (Fig. S1b). Interestingly, phenocopying the *Fn*<sup>iKO</sup> mutants, *Fn*<sup>AB KO</sup> mice also displayed reduced vessel migration, branching and coverage within their retinas at P6 (Fig. 2e–h). Because EIIIA/EIIIB-containing fibronectin was predominantly localised around the developing vasculature in control




**Fig. 1** Fibronectin localisation within the retinal vasculature. **a–f** Whole-mount immunofluorescence staining showing the localisation of fibronectin and its alternatively spliced EIIIA<sup>+</sup> and EIIIB<sup>+</sup>-containing variants within the retinal vasculature of a P6 mouse. Fibronectin is localised throughout the vasculature (**a**) and is assembled just ahead of the vascular plexus by astrocytes (*arrowheads* **a–c**). EIIIA<sup>+</sup> (**b**) and EIIIB<sup>+</sup> (**c**) domains are present at low

levels in the fibronectin network assembled by astrocytes (*arrowheads* **b, c**), but are found at higher levels in the fibronectin surrounding blood vessels. Note that expression of EIIIA (**d**, enlarged in **e**) and EIIIB (**f**) are especially pronounced around the edges of transcapillary pillars (*holes*) within vessels (*arrows*). Scale bars: 50 μm (**a–d**); 10 μm (**e, f**)

retinas (Fig. 1b–f), and branching defects have not been reported in mice lacking astrocyte-derived fibronectin [27], we next examined whether the defects in both the  $Fn^{iKO}$

and  $Fn^{AB KO}$  could be due to the loss of just EC-derived EIIIA<sup>+</sup>/EIIIB<sup>+</sup> fibronectin. To avoid embryonic developmental defects, we once again used an inducible loss-of-



**Fig. 2** Abnormal patterning of the retinal vasculature in *Fn* mutants. **a–f** Confocal micrographs of the retinal vasculature in control, *Fn<sup>iKO</sup>*, *Fn<sup>AB KO</sup>* and *Fn<sup>iEC KO</sup>* mice at P6. **a, b** Representative low magnification images showing that, in the absence of fibronectin, *Fn<sup>iKO</sup>* mice develop a vascular plexus containing arteries (A), veins (V) and capillaries, but display reduced vessel outgrowth (*dashed lines* illustrate measurements used to quantify distance of vessel

migration), vascular branching and vascular coverage in their retinas (**b**, higher magnification **c, d**). These defects are phenocopied in *Fn<sup>AB KO</sup>* (**e**) and *Fn<sup>iEC KO</sup>* (**f**) mice. Quantification of vessel outgrowth (**g**), vessel branching (**h**) and endothelial coverage (**i**) (*see Methods*) reveals defects in the *Fn* mutants ( $n = 6$ , mice per genotype). *Scale bars*: 500 μm (**a, b**), 100 μm (**c–f**)

function approach and deleted *Fn* expression specifically in ECs from P1 onwards through the administration of tamoxifen to mice carrying a *Cdh5*(PAC)-CreER<sup>T2</sup> transgene [34] and loxP-flanked *Fn* gene (*Fn<sup>iEC KO</sup>*). Just as observed in *Fn<sup>iKO</sup>* and *Fn<sup>AB KO</sup>* mice, radial expansion, branching and density of retinal blood vessels were all significantly reduced in *Fn<sup>iEC KO</sup>* mutants (Fig. 2f, g–i).

However, in contrast to both *Fn<sup>iKO</sup>* and *Fn<sup>AB KO</sup>* mutants (Fig. S1), and despite efficient Cre-mediated excision from endothelial cells (Fig. S2), blood vessels in *Fn<sup>iEC KO</sup>* mice remained covered with EIIIA<sup>+</sup>/EIIIB<sup>+</sup> fibronectin at similar levels as observed in controls (Fig. 3a–c). This supports previous data showing that the majority of the fibronectin deposited around the vasculature in the retina is

derived from astrocytes [27, 37], but suggests that astrocyte fibronectin cannot compensate fully for the loss of endothelial-derived fibronectin. Vessel patterning in the retina therefore is regulated, at least in part, in an autocrine manner by endothelial EIIIA<sup>+</sup>/EIIIB<sup>+</sup> fibronectin.


### Endothelium-derived fibronectin affects tip cell numbers in the retina

Examination of the leading edge of the vascular plexus revealed that the reduced vessel density in all three *Fn* mutants is, in part, due to a reduced number of tip cells at the angiogenic front of the retinal vasculature (Fig. 4a–e). In addition, global loss of fibronectin also led to increased numbers of thick, long, abnormally shaped angiogenic sprouts at the front of the plexus (Fig. 4a–d). Loss of fibronectin did not appear however to affect either the number or length of filopodia extending from individual tip cells in *Fn* mutants (Fig. 4f, g). In addition, filopodia extending from endothelial cells in *Fn*<sup>iEC KO</sup> mutants aligned with the fibronectin network assembled ahead of the plexus by the astrocytes (Fig. 3a).

**Fig. 4** *Fn* mutants display reduced numbers of tip cells. Confocal images of isolectin-B4-stained tip cells at the angiogenic front of (a) control, (b) *Fn*<sup>iKO</sup>, (c) *Fn*<sup>AB KO</sup> and (d) *Fn*<sup>iEC KO</sup> retinas at P6. *Fn* mutants have reduced numbers of extending tip cell vessel sprouts ( $n = 3$ , mice per genotype) (e), but have similar numbers of filopodial extensions ( $n = 12$ , tip cells) (f) and filopodial lengths ( $n = 50$ , filopodia) (g) per tip cell as control mice. Higher magnification images of individual tip cells and filopodial extensions in right panels. Note the thicker more irregular tip cell morphology in *Fn*<sup>iKO</sup> mutants (a–d). NS not significant. Scale bars: 50  $\mu$ m





### Endothelium-derived fibronectin regulates vessel stability


Further analysis revealed that EC-derived fibronectin also has a critical role in controlling vessel stability. At P6, *Fn*<sup>iEC KO</sup> mutants had increased numbers of thin empty PECAM1-negative/collagen IV-positive basement membrane sleeves, which are left behind by regressing endothelial cells and serve as a historical record of pre-existing vessels [40, 41], indicating increased pruning/regression of vessels (Fig. 5a, b). Proliferation of endothelial cells in *Fn*<sup>iEC KO</sup> mutants however appeared largely unaffected by the loss of fibronectin (Fig. 5c, d). Previous studies have shown that vessel stability is dependent on the



**Fig. 3** Deposition of fibronectin splice variants around the vessel wall of *Fn*<sup>iEC KO</sup> mice. Whole-mount immunofluorescence staining showing that *Fn*<sup>iEC KO</sup> mice develop vascular defects despite the presence of exogenous EIIIA<sup>+</sup> (a) and EIIIB<sup>+</sup> (b) fibronectin surrounding their vessels. c Quantification of fibronectin

immunofluorescence around the retinal vasculature of *Fn* mutants showing that equivalent levels of fibronectin are deposited around the vessels of control, *Fn*<sup>iEC KO</sup> and *Fn*<sup>AB KO</sup> mice at P6 ( $n = 10$ , FOV). NS not significant. Scale bars: 50  $\mu$ m





**Fig. 5** Loss of EC-derived fibronectin leads to ectopic vessel regression. **a**  $Fn^{IEC\ KO}$  mice display increased numbers of thin, empty (PECAM1-negative) sleeves of collagen IV matrix (arrows), indicating increased levels of vessel regression within their retinas at P6. **b** Quantification of PECAM1<sup>-</sup> collagen IV<sup>+</sup> vessel regression segments ( $n = 6$ , FOV). **c** Immunofluorescence staining for the proliferation marker Ki-67, and **(d)** quantification of proliferating

endothelial cells, revealed no significant differences in endothelial cell (EC) proliferation within the retinas of  $Fn^{IEC\ KO}$  mice at P6 ( $n = 2$ , mice per genotype). **e** Desmin-positive pericytes remain in close association with the capillary endothelium in  $Fn^{IEC\ KO}$  mice, and are found in similar numbers to those seen in control mice **(f)** ( $n = 6$ , FOV). Scale bars: 50  $\mu\text{m}$  (**a**, **e**); 100  $\mu\text{m}$  (**c**)

recruitment of mural cells [42], namely pericytes and vascular smooth muscle cells. We therefore analysed whether loss of EC-derived fibronectin inhibited the recruitment and incorporation of mural cells around retinal vessels. Analysis of P6 retinas stained with anti-desmin antibodies however revealed no obvious defects in mural cell recruitment or attachment to the capillaries of  $Fn^{IEC\ KO}$  mutants (Fig. 5e, f).


### Mice lacking endothelial $\alpha 5$ and $\alpha v$ integrins phenocopy $Fn^{IEC\ KO}$ mutants

Since  $Fn^{IEC\ KO}$  mice displayed vascular defects, despite displaying apparently normal levels of cellular fibronectin around their vessels (Fig. 3), we next investigated whether vascular patterning in the retina is dependent on expression of the major endothelial fibronectin-binding receptors,



integrin  $\alpha 5$  and  $\alpha v$ . Just like fibronectin,  $\alpha 5$  and  $\alpha v$  integrins are poorly expressed on quiescent endothelium but are highly expressed around blood vessels during developmental or tumour angiogenesis. Previous studies have shown that mice lacking endothelial expression of both  $\alpha 5$  and  $\alpha v$  die at E14.5 with heart, great vessel and lymphatic defects [26, 43] but, in contrast to numerous in vitro studies, lack angiogenic defects [26]. To examine the role of endothelial  $\alpha 5$  and  $\alpha v$  integrins in postnatal developmental angiogenesis, we crossed female double-homozygous *Itga5/Itgav*-floxed mice to *Cdh5(PAC)-CreER<sup>T2</sup>* mice (to generate *Itga5/av<sup>iEC KO</sup>* mice) and deleted both genes

from the endothelium through administration of tamoxifen from P1 to bypass embryonic lethality. Despite the requirement for both  $\alpha 5$  and  $\alpha v$  integrins for fibronectin fibrillogenesis in vitro, consistent with previous in vivo studies [25, 26], *Itga5/av<sup>iEC KO</sup>* mice displayed no obvious defects in the assembly of Fn around their vasculature (Fig. 6a). However, just as observed in *Fn<sup>iEC KO</sup>* mutants, doubly deficient *Itga5/av<sup>iEC KO</sup>* (but not singly deficient *Itga5<sup>iEC KO</sup>* or *Itgav<sup>iEC KO</sup>* mice, data not shown) still displayed reduced radial growth, vessel branching, endothelial coverage, and tip cell numbers within their retinas at P6 (Fig. 6b–g). Surprisingly, in contrast to mice



**Fig. 6** Loss of both endothelial  $\alpha 5$  and  $\alpha v$  integrins phenocopies *Fn* mutants. **a** Immunofluorescence staining showing deposition of fibronectin around the retinal vasculature of an *Itga5/av<sup>iEC KO</sup>* mouse at P6, despite the lack of the major fibronectin-binding integrins. **b** Isolectin-B4-labeled retinas showing the decreased vessel branching and density in *Itga5/av<sup>iEC KO</sup>* retinas. Quantification of **(c)** radial

outgrowth, **(d)** branchpoints and **(e)** endothelial coverage in *Itga5/av<sup>iEC KO</sup>* mutants ( $n = 6$ , mice per genotype). **f** Representative image of the tip-cell sprouts at the angiogenic front of the developing vasculature in *Itga5/av<sup>iEC KO</sup>* mice. **g** Quantification of *Itga5/av<sup>iEC KO</sup>* tip-cell deficit ( $n = 3$ , mice per genotype). Scale bars: 50  $\mu\text{m}$  (**a**, **b**, **f**)

in which *Itga5* had been deleted from the endothelium using Tie2-Cre [27], endothelial tip-cell filopodia appeared unaffected by the loss of both  $\alpha 5$  and  $\alpha v$  integrins and aligned to the fibronectin network assembled by astrocytes (Fig. 6a). The possibility exists that other fibronectin-binding integrins could also be involved in retinal angiogenesis or that they might partially compensate for the absence of  $\alpha 5$  and  $\alpha v$ . Nonetheless the results do show the involvement of these two integrin subunits.

## Discussion

In this study, we have shown that vessel patterning in the retina is regulated in an autocrine manner by EC-derived EIIIA<sup>+</sup>/EIIIB<sup>+</sup> fibronectin and that these functions are not fully compensated by exogenous sources of fibronectin such as astrocytes or plasma. We have also shown that postnatal developmental angiogenesis requires endothelial expression of at least one of the RGD-binding integrin receptors and that interaction of  $\alpha 5$  and  $\alpha v$  integrins with EC-derived fibronectin regulates vessel sprouting and vessel stability during vascular development.

### Fibronectin regulates blood vessel development

It has long been suggested that fibronectin plays a key role in the development of the vascular system. Fibronectin is found only in vertebrates with an endothelial-lined vasculature [10] and forms a central node within the “angiome” [44]. The exact functions of Fn and its splice variants during vascular development however have remained unclear due to mesodermal, neural tube and cardiac defects hampering the interpretation of the vascular defects observed in existing *Fn* models [12, 17, 18, 21]. Our detailed analyses of retinal vascular development show that, in the absence of embryonic defects, mice temporally lacking *Fn* expression throughout the tissue develop severe vascular patterning defects (Fig. 2). *Fn*<sup>iKO</sup> mice are able to develop vessels, but radial expansion, vessel branching and density of blood vessels are all reduced in the absence of *Fn* (Fig. 2). While the reduced radial outgrowth of blood vessels can in part be attributed to the loss of astrocyte-derived Fn within the retina and reduced VEGFR2 and PI3 K/Akt signalling [27], our results indicate that it is the loss of EC-derived *Fn* that is the predominant cause of the vascular defects observed. *Fn*<sup>iEC KO</sup> mutants display decreased vessel outgrowth, branching, and vessel coverage, phenocopying the defects observed in global *Fn* KO mice (Fig. 2). These defects are, at least in part, due to decreased levels of vessel sprouting at the angiogenic front (Fig. 4) and increased regression and pruning within the vascular plexus (Fig. 5). Since Fn expression is especially

pronounced around the edges of transcapillary pillars, we cannot however rule out the possibility that it may also regulate intussusceptive angiogenesis, which has also been shown to increase vessel branching and vascular expansion in numerous tissues including the retina [45].

### Cell autonomous roles for fibronectin during vascular development

A remarkable finding in our study is that *Fn*<sup>iEC KO</sup> mice develop vascular defects despite displaying apparently normal levels of Fn around their vessels (Fig. 3), suggesting that EC-derived Fn plays a distinct role in regulating vascular development. During the development of the retina, *Fn* is expressed by ECs, astrocytes [27, 37] and, to some extent, pericytes [46], although the functional importance of the latter appears to be minimal since no defects have been observed in mice lacking pericyte expression of *Fn* (data not shown). Astrocytes deposit a Fn scaffold ahead of the vascular plexus to support EC migration into the avascular areas of the retina (Fig. 1), but upon contact with the advancing vasculature astrocytes downregulate fibronectin expression [27, 37]. Interestingly, in contrast to *Fn*<sup>iEC KO</sup> mutants, astrocyte-specific deletion of *Fn* leads to increased numbers of tip-cell filopodia, increased branching and an increase in vessel density [27]. Taken together, this suggests that Fn regulates vascular development through distinct paracrine and autocrine mechanisms, with astrocyte Fn inhibiting and EC-derived Fn promoting sprouting and branching of vasculature. Autocrine fibronectin has previously been suggested to have a distinct role in controlling EC behaviour in vitro [29], and has been shown to play an important role in regulating cardiovascular development in vivo [47]. The exact mechanisms by which autocrine Fn elicits its differential response however remain unclear.

### Roles of EIIIA/EIIIB-containing fibronectin

One possible hypothesis for the differential response to paracrine and autocrine Fn is that ECs may produce a different form of fibronectin. Fn containing EIIIA/EIIIB domains is highly expressed around the developing vasculature, but these domains are almost undetectable in the Fn scaffold deposited by astrocytes in the retina (Fig. 1). Furthermore, loss of both EIIIA and EIIIB domains, but of neither one alone (data not shown), replicates the vascular defects observed in the *Fn*<sup>iEC KO</sup> mutants (Figs. 2, 4). Arguing against this hypothesis however is the observation that *Fn*<sup>iEC KO</sup> mutants still display EIIIA- and EIIIB-containing fibronectin around their vessels (Fig. S1b). A caveat of this analysis however is that it is almost impossible to distinguish whether the Fn surrounding the vessels is

EIIIA<sup>+</sup> EIIBB<sup>+</sup> or just fibrils of Fn containing a mixture of EIIIA<sup>+</sup> EIIBB<sup>-</sup> and EIIIA<sup>-</sup> EIIBB<sup>+</sup> fibronectins. Since neither EIIIA KO nor EIIBB single KO mice display any vascular defects [19], it is possible that only EIIIA<sup>+</sup> EIIBB<sup>+</sup> Fn is expressed by ECs, and it is this specific form of Fn that is essential for regulating EC function. So, how might the EIIIA and EIIBB domains regulate EC function? Previous studies have shown that Fn can bind and regulate the activity of numerous growth factors [48], it is possible therefore that the addition of EIIIA and EIIBB domains may modulate growth factor signalling within the vascular endothelium. Indeed, EIIIA and EIIBB KO cells have reduced growth and proliferation in vitro [20, 49]. It is unlikely that this is the cause of the vascular phenotype seen in *Fn*<sup>iEC KO</sup> mice however, since no obvious proliferation defects were observed in our mutant mice (Fig. 5). A second possibility is that the addition of EIIIA and EIIBB domains may alter the physiological properties of Fn. Fibronectin is one of the most extendable biological fibres [50, 51] and upon extension becomes more rigid [52]. It is conceivable therefore that addition of EIIIA and EIIBB may alter the structural and mechano-transductive properties of the protein. Fn fibrillogenesis [53], assembly of collagen I [54], and vasodilation of vessels [55] have all been shown to be mechano-regulated by stretch-induced conformational changes in Fn. Finally, insertion of additional EIIIA and EIIBB domains may increase adhesiveness of Fn to its integrin receptors. The EIIIA domain contains additional binding sites for  $\alpha 4\beta 1$  and  $\alpha 9\beta 1$  integrins [56], while inclusion of EIIBB has been shown to induce a conformational change that unmask a cryptic binding site [57] and affects the exposure of the RGD loop [58] recognised by both  $\alpha 5$  and  $\alpha v$  integrins within Fn.

### Role of $\alpha 5$ and $\alpha v$ integrins during vascular development

In contrast to studies in the embryo [26] and during tumorigenesis [30], we found that endothelial expression of both  $\alpha 5$  and  $\alpha v$  integrins is essential for proper angiogenesis in the retina (Fig. 6). Just as observed in both *Fn*<sup>iEC KO</sup> and *Fn*<sup>AB KO</sup> mutants, loss of both  $\alpha 5$  and  $\alpha v$  integrins leads to defects in vessel growth, branching, and vascular sprouting (Fig. 6). Interestingly, similar defects have been reported in mice lacking EC expression of integrin  $\beta 1$  [7], suggesting that interaction of EC-derived EIIIA<sup>+</sup> EIIBB<sup>+</sup> Fn with  $\alpha 5\beta 1$  and  $\alpha v\beta 1$  regulates vessel patterning and stability during retinal angiogenesis. We have previously shown that  $\alpha 5$  and  $\alpha v$  integrins cooperate to regulate vascular smooth muscle cell function in vivo [25], it is therefore increasingly clear that both receptors play a key role(s) in controlling the development of the vascular

system. These results do not rule out the participation of other integrins recognising fibronectin or, indeed, other ECM proteins in retinal angiogenesis.

### Conclusion

Very few prior studies have given much attention to the cellular sources or specific splice variants of Fn within their experiments. Our results have shown that EC-derived Fn provides distinct signals from those derived from exogenous sources of Fn (such as astrocytes, pericytes, and the plasma) and is indispensable for proper vascular development in the retina. Furthermore, we have shown that EC-derived fibronectin requires both EIIIA and EIIBB domains for its function and that it signals through  $\alpha 5$  and  $\alpha v$  integrins to regulate vessel patterning. It is quite possible that detailed analyses of other angiogenic processes may reveal analogous distinctions among the contributions of different cell types and isoforms to specific aspects of angiogenesis. The exact mechanisms by which EIIIA and EIIBB domains within Fn regulate angiogenesis and the precise roles of the individual Fn integrins on vascular cells however remain unclear. Future experiments will need to examine the biomechanical and biochemical signalling changes caused by the addition of EIIIA and EIIBB domains within fibronectin and use multiple inducible cell-specific integrin and Fn mutants. It will also be important to identify the genetic modifiers that strongly influence the phenotypes of both integrin and fibronectin mutants.

**Acknowledgements** We thank members of the Hynes laboratory, especially Patrick Murphy, for discussions and advice. This work was supported by grants from the National Institutes of Health (PO1-HL66105, PI, Monty Krieger), the NIGMS Cell Migration Consortium, (GC11451.126452, PI, A.F. Horwitz), by the Koch Institute Support (core) Grant P30-CA14051 from the National Cancer Institute and by the Howard Hughes Medical Institute. CJT was a post-doctoral associate and ROH is an Investigator of the Howard Hughes Medical Institute.

**Author contributions** Experiments were conceived, designed and interpreted by CJT, KB-N, and ROH. Experiments were performed by CJT and KB-N. The manuscript was written by CJT and ROH.

### Compliance with ethical standards

**Conflict of interest** The authors declared that they have no conflict of interest.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Senger DR, Davis GE (2011) Angiogenesis. *Cold Spring Harb Perspect Biol* 3(8):a005090. doi:[10.1101/cshperspect.a005090](https://doi.org/10.1101/cshperspect.a005090)
- Hynes RO (2009) The extracellular matrix: not just pretty fibrils. *Science* 326(5957):1216–1219. doi:[10.1126/science.1176009](https://doi.org/10.1126/science.1176009)
- Hynes RO (2007) Cell-matrix adhesion in vascular development. *J Thromb Haemost* 5(Suppl 1):32–40. doi:[10.1111/j.1538-7836.2007.02569.x](https://doi.org/10.1111/j.1538-7836.2007.02569.x)
- Avraamides CJ, Garmy-Susini B, Varner JA (2008) Integrins in angiogenesis and lymphangiogenesis. *Nat Rev Cancer* 8(8):604–617. doi:[10.1038/nrc2353](https://doi.org/10.1038/nrc2353)
- Senger DR, Perruzzi CA, Streit M, Kotliansky VE, de Fougères AR, Detmar M (2002) The alpha(1)beta(1) and alpha(2)-beta(1) integrins provide critical support for vascular endothelial growth factor signaling, endothelial cell migration, and tumor angiogenesis. *Am J Pathol* 160(1):195–204
- Smith JT, Elkin JT, Reichert WM (2006) Directed cell migration on fibronectin gradients: effect of gradient slope. *Exp Cell Res* 312(13):2424–2432. doi:[10.1016/j.yexcr.2006.04.005](https://doi.org/10.1016/j.yexcr.2006.04.005)
- Yamamoto H, Ehling M, Kato K, Kanai K, van Lessen M, Frye M, Zeuschner D, Nakayama M, Vestweber D, Adams RH (2015) Integrin beta1 controls VE-cadherin localization and blood vessel stability. *Nat Commun* 6:6429. doi:[10.1038/ncomms7429](https://doi.org/10.1038/ncomms7429)
- Iruela-Arispe ML, Davis GE (2009) Cellular and molecular mechanisms of vascular lumen formation. *Dev Cell* 16(2):222–231. doi:[10.1016/j.devcel.2009.01.013](https://doi.org/10.1016/j.devcel.2009.01.013)
- Astrof S, Hynes RO (2009) Fibronectins in vascular morphogenesis. *Angiogenesis* 12(2):165–175. doi:[10.1007/s10456-009-9136-6](https://doi.org/10.1007/s10456-009-9136-6)
- Whittaker CA, Bergeron KF, Whittle J, Brandhorst BP, Burke RD, Hynes RO (2006) The echinoderm adhesome. *Dev Biol* 300(1):252–266. doi:[10.1016/j.ydbio.2006.07.044](https://doi.org/10.1016/j.ydbio.2006.07.044)
- Risau W, Lemmon V (1988) Changes in the vascular extracellular matrix during embryonic vasculogenesis and angiogenesis. *Dev Biol* 125(2):441–450
- George EL, Georges-Labouesse EN, Patel-King RS, Rayburn H, Hynes RO (1993) Defects in mesoderm, neural tube and vascular development in mouse embryos lacking fibronectin. *Development* 119(4):1079–1091
- Peters JH, Hynes RO (1996) Fibronectin isoform distribution in the mouse. I. The alternatively spliced EIIIB, EIIIA, and V segments show widespread codistribution in the developing mouse embryo. *Cell Adhes Commun* 4(2):103–125
- Ffrench-Constant C, Hynes RO (1989) Alternative splicing of fibronectin is temporally and spatially regulated in the chicken embryo. *Development* 106(2):375–388
- Dubin D, Peters JH, Brown LF, Logan B, Kent KC, Berse B, Berven S, Cercek B, Sharifi BG, Pratt RE et al (1995) Balloon catheterization induced arterial expression of embryonic fibronectins. *Arterioscler Thromb Vasc Biol* 15(11):1958–1967
- Murphy PA, Hynes RO (2014) Alternative splicing of endothelial fibronectin is induced by disturbed hemodynamics and protects against hemorrhage of the vessel wall. *Arterioscler Thromb Vasc Biol* 34(9):2042–2050. doi:[10.1161/ATVBAHA.114.303879](https://doi.org/10.1161/ATVBAHA.114.303879)
- Takahashi S, Leiss M, Moser M, Ohashi T, Kitao T, Heckmann D, Pfeifer A, Kessler H, Takagi J, Erickson HP, Fassler R (2007) The RGD motif in fibronectin is essential for development but dispensable for fibril assembly. *J Cell Biol* 178(1):167–178. doi:[10.1083/jcb.200703021](https://doi.org/10.1083/jcb.200703021)
- George EL, Baldwin HS, Hynes RO (1997) Fibronectins are essential for heart and blood vessel morphogenesis but are dispensable for initial specification of precursor cells. *Blood* 90(8):3073–3081
- Astrof S, Crowley D, George EL, Fukuda T, Sekiguchi K, Hanahan D, Hynes RO (2004) Direct test of potential roles of EIIIA and EIIIB alternatively spliced segments of fibronectin in physiological and tumor angiogenesis. *Mol Cell Biol* 24(19):8662–8670. doi:[10.1128/MCB.24.19.8662-8670.2004](https://doi.org/10.1128/MCB.24.19.8662-8670.2004)
- Fukuda T, Yoshida N, Kataoka Y, Manabe R, Mizuno-Horikawa Y, Sato M, Kuriyama K, Yasui N, Sekiguchi K (2002) Mice lacking the EDB segment of fibronectin develop normally but exhibit reduced cell growth and fibronectin matrix assembly in vitro. *Cancer Res* 62(19):5603–5610
- Astrof S, Crowley D, Hynes RO (2007) Multiple cardiovascular defects caused by the absence of alternatively spliced segments of fibronectin. *Dev Biol* 311(1):11–24. doi:[10.1016/j.ydbio.2007.07.005](https://doi.org/10.1016/j.ydbio.2007.07.005)
- Astrof S, Kirby A, Lindblad-Toh K, Daly M, Hynes RO (2007) Heart development in fibronectin-null mice is governed by a genetic modifier on chromosome four. *Mech Dev* 124(7–8):551–558. doi:[10.1016/j.mod.2007.05.004](https://doi.org/10.1016/j.mod.2007.05.004)
- Trinh LA, Stainier DY (2004) Fibronectin regulates epithelial organization during myocardial migration in zebrafish. *Dev Cell* 6(3):371–382
- Giros A, Grgur K, Gossler A, Costell M (2011) alpha5beta1 integrin-mediated adhesion to fibronectin is required for axis elongation and somitogenesis in mice. *PLoS ONE* 6(7):e22002. doi:[10.1371/journal.pone.0022002](https://doi.org/10.1371/journal.pone.0022002)
- Turner CJ, Badu-Nkansah K, Crowley D, van der Flier A, Hynes RO (2015) alpha5 and alphaV integrins cooperate to regulate vascular smooth muscle and neural crest functions in vivo. *Development* 142(4):797–808. doi:[10.1242/dev.117572](https://doi.org/10.1242/dev.117572)
- van der Flier A, Badu-Nkansah K, Whittaker CA, Crowley D, Bronson RT, Lacy-Hulbert A, Hynes RO (2010) Endothelial alpha5 and alphaV integrins cooperate in remodeling of the vasculature during development. *Development* 137(14):2439–2449. doi:[10.1242/dev.049551](https://doi.org/10.1242/dev.049551)
- Stenzel D, Lundkvist A, Sauvaget D, Busse M, Graupera M, van der Flier A, Wijelath ES, Murray J, Sobel M, Costell M, Takahashi S, Fassler R, Yamaguchi Y, Gutmann DH, Hynes RO, Gerhardt H (2011) Integrin-dependent and -independent functions of astrocytic fibronectin in retinal angiogenesis. *Development* 138(20):4451–4463. doi:[10.1242/dev.071381](https://doi.org/10.1242/dev.071381)
- von Au A, Vasel M, Kraft S, Sens C, Hackl N, Marx A, Stroebel P, Hennenlotter J, Todenhofer T, Stenzl A, Schott S, Sinn HP, Wetterwald A, Bermejo JL, Cecchini MG, Nakchbandi IA (2013) Circulating fibronectin controls tumor growth. *Neoplasia* 15(8):925–938
- Cseh B, Fernandez-Sauze S, Grall D, Schaub S, Doma E, Van Obberghen-Schilling E (2010) Autocrine fibronectin directs matrix assembly and crosstalk between cell–matrix and cell–cell adhesion in vascular endothelial cells. *J Cell Sci* 123(Pt 22):3989–3999. doi:[10.1242/jcs.073346](https://doi.org/10.1242/jcs.073346)
- Murphy PA, Begum S, Hynes RO (2015) Tumor angiogenesis in the absence of fibronectin or its cognate integrin receptors. *PLoS ONE* 10(3):e0120872. doi:[10.1371/journal.pone.0120872](https://doi.org/10.1371/journal.pone.0120872)
- Sakai T, Johnson KJ, Murozono M, Sakai K, Magnuson MA, Wieloch T, Cronberg T, Isshiki A, Erickson HP, Fassler R (2001) Plasma fibronectin supports neuronal survival and reduces brain injury following transient focal cerebral ischemia but is not essential for skin-wound healing and hemostasis. *Nat Med* 7(3):324–330. doi:[10.1038/85471](https://doi.org/10.1038/85471)
- Lacy-Hulbert A, Smith AM, Tissire H, Barry M, Crowley D, Bronson RT, Roes JT, Savill JS, Hynes RO (2007) Ulcerative colitis and autoimmunity induced by loss of myeloid alphaV integrins. *Proc Natl Acad Sci U S A* 104(40):15823–15828. doi:[10.1073/pnas.0707421104](https://doi.org/10.1073/pnas.0707421104)
- Ventura A, Kirsch DG, McLaughlin ME, Tuveson DA, Grimm J, Lintault L, Newman J, Reczek EE, Weissleder R, Jacks T (2007) Restoration of p53 function leads to tumour regression in vivo. *Nature* 445(7128):661–665. doi:[10.1038/nature05541](https://doi.org/10.1038/nature05541)

34. Wang Y, Nakayama M, Pitulescu ME, Schmidt TS, Bochenek ML, Sakakibara A, Adams S, Davy A, Deutsch U, Luthi U, Barberis A, Benjamin LE, Makinen T, Nobes CD, Adams RH (2010) Ephrin-B2 controls VEGF-induced angiogenesis and lymphangiogenesis. *Nature* 465(7297):483–486. doi:[10.1038/nature09002](https://doi.org/10.1038/nature09002)
35. Muzumdar MD, Tasic B, Miyamichi K, Li L, Luo L (2007) A global double-fluorescent Cre reporter mouse. *Genesis* 45(9):593–605. doi:[10.1002/dvg.20335](https://doi.org/10.1002/dvg.20335)
36. Pitulescu ME, Schmidt I, Benedetto R, Adams RH (2010) Inducible gene targeting in the neonatal vasculature and analysis of retinal angiogenesis in mice. *Nat Protoc* 5(9):1518–1534. doi:[10.1038/nprot.2010.113](https://doi.org/10.1038/nprot.2010.113)
37. Uemura A, Kusuhara S, Wiegand SJ, Yu RT, Nishikawa S (2006) Tlx acts as a proangiogenic switch by regulating extracellular assembly of fibronectin matrices in retinal astrocytes. *J Clin Invest* 116(2):369–377. doi:[10.1172/JCI25964](https://doi.org/10.1172/JCI25964)
38. Zhou X, Rowe RG, Hiraoka N, George JP, Wirtz D, Mosher DF, Virtanen I, Chernousov MA, Weiss SJ (2008) Fibronectin fibrillogenesis regulates three-dimensional neovessel formation. *Genes Dev* 22(9):1231–1243. doi:[10.1101/gad.1643308](https://doi.org/10.1101/gad.1643308)
39. Francis SE, Goh KL, Hodivala-Dilke K, Bader BL, Stark M, Davidson D, Hynes RO (2002) Central roles of alpha5beta1 integrin and fibronectin in vascular development in mouse embryos and embryoid bodies. *Arterioscler Thromb Vasc Biol* 22(6):927–933
40. Baffert F, Le T, Sennino B, Thurston G, Kuo CJ, Hu-Lowe D, McDonald DM (2006) Cellular changes in normal blood capillaries undergoing regression after inhibition of VEGF signaling. *Am J Physiol Heart Circ Physiol* 290(2):H547–H559. doi:[10.1152/ajpheart.00616.2005](https://doi.org/10.1152/ajpheart.00616.2005)
41. Baluk P, Morikawa S, Haskell A, Mancuso M, McDonald DM (2003) Abnormalities of basement membrane on blood vessels and endothelial sprouts in tumors. *Am J Pathol* 163(5):1801–1815. doi:[10.1016/S0002-9440\(10\)63540-7](https://doi.org/10.1016/S0002-9440(10)63540-7)
42. Gaengel K, Genove G, Armulik A, Betsholtz C (2009) Endothelial-mural cell signaling in vascular development and angiogenesis. *Arterioscler Thromb Vasc Biol* 29(5):630–638. doi:[10.1161/ATVBAHA.107.161521](https://doi.org/10.1161/ATVBAHA.107.161521)
43. Turner CJ, Badu-Nkansah K, Crowley D, van der Flier A, Hynes RO (2014) Integrin-alpha5beta1 is not required for mural cell functions during development of blood vessels but is required for lymphatic-blood vessel separation and lymphovenous valve formation. *Dev Biol* 392(2):381–392. doi:[10.1016/j.ydbio.2014.05.006](https://doi.org/10.1016/j.ydbio.2014.05.006)
44. Chu LH, Rivera CG, Popel AS, Bader JS (2012) Constructing the angiome: a global angiogenesis protein interaction network. *Physiol Genomics* 44(19):915–924. doi:[10.1152/physiolgenomics.00181.2011](https://doi.org/10.1152/physiolgenomics.00181.2011)
45. Burri PH, Hlushchuk R, Djonov V (2004) Intussusceptive angiogenesis: its emergence, its characteristics, and its significance. *Dev Dyn* 231(3):474–488. doi:[10.1002/dvdy.20184](https://doi.org/10.1002/dvdy.20184)
46. Stratman AN, Malotte KM, Mahan RD, Davis MJ, Davis GE (2009) Pericyte recruitment during vasculogenic tube assembly stimulates endothelial basement membrane matrix formation. *Blood* 114(24):5091–5101. doi:[10.1182/blood-2009-05-222364](https://doi.org/10.1182/blood-2009-05-222364)
47. Wang X, Astrof S (2016) Neural crest cell-autonomous roles of fibronectin in cardiovascular development. *Development* 143(1):88–100. doi:[10.1242/dev.125286](https://doi.org/10.1242/dev.125286)
48. Miyamoto S, Teramoto H, Gutkind JS, Yamada KM (1996) Integrins can collaborate with growth factors for phosphorylation of receptor tyrosine kinases and MAP kinase activation: roles of integrin aggregation and occupancy of receptors. *J Cell Biol* 135(6 Pt 1):1633–1642
49. White ES, Baralle FE, Muro AF (2008) New insights into form and function of fibronectin splice variants. *J Pathol* 216(1):1–14. doi:[10.1002/path.2388](https://doi.org/10.1002/path.2388)
50. Ohashi T, Kiehart DP, Erickson HP (1999) Dynamics and elasticity of the fibronectin matrix in living cell culture visualized by fibronectin-green fluorescent protein. *Proc Natl Acad Sci U S A* 96(5):2153–2158
51. Little WC, Smith ML, Ebnetter U, Vogel V (2008) Assay to mechanically tune and optically probe fibrillar fibronectin conformations from fully relaxed to breakage. *Matrix Biol* 27(5):451–461. doi:[10.1016/j.matbio.2008.02.003](https://doi.org/10.1016/j.matbio.2008.02.003)
52. Klotzsch E, Smith ML, Kubow KE, Muntwyler S, Little WC, Beyeler F, Gourdon D, Nelson BJ, Vogel V (2009) Fibronectin forms the most extensible biological fibers displaying switchable force-exposed cryptic binding sites. *Proc Natl Acad Sci U S A* 106(43):18267–18272. doi:[10.1073/pnas.0907518106](https://doi.org/10.1073/pnas.0907518106)
53. Zhong C, Chrzanowska-Wodnicka M, Brown J, Shaub A, Belkin AM, Burrige K (1998) Rho-mediated contractility exposes a cryptic site in fibronectin and induces fibronectin matrix assembly. *J Cell Biol* 141(2):539–551
54. Kubow KE, Vukmirovic R, Zhe L, Klotzsch E, Smith ML, Gourdon D, Luna S, Vogel V (2015) Mechanical forces regulate the interactions of fibronectin and collagen I in extracellular matrix. *Nat Commun* 6:8026. doi:[10.1038/ncomms9026](https://doi.org/10.1038/ncomms9026)
55. Hocking DC, Titus PA, Sumagin R, Sarelius IH (2008) Extracellular matrix fibronectin mechanically couples skeletal muscle contraction with local vasodilation. *Circ Res* 102(3):372–379. doi:[10.1161/CIRCRESAHA.107.158501](https://doi.org/10.1161/CIRCRESAHA.107.158501)
56. Liao YF, Gotwals PJ, Kotliansky VE, Sheppard D, Van De Water L (2002) The EIIIA segment of fibronectin is a ligand for integrins alpha 9beta 1 and alpha 4beta 1 providing a novel mechanism for regulating cell adhesion by alternative splicing. *J Biol Chem* 277(17):14467–14474. doi:[10.1074/jbc.M201100200](https://doi.org/10.1074/jbc.M201100200)
57. Balza E, Sassi F, Ventura E, Parodi A, Fossati S, Blalock W, Carnemolla B, Castellani P, Zardi L, Borsi L (2009) A novel human fibronectin cryptic sequence unmasked by the insertion of the angiogenesis-associated extra type III domain B. *Int J Cancer* 125(4):751–758. doi:[10.1002/ijc.24473](https://doi.org/10.1002/ijc.24473)
58. Carnemolla B, Leprini A, Allemanni G, Saginati M, Zardi L (1992) The inclusion of the type III repeat ED-B in the fibronectin molecule generates conformational modifications that unmask a cryptic sequence. *J Biol Chem* 267(34):24689–24692