CORRECTION

Correction to: Climate change impacts on potential distribution of multipurpose agro-forestry species: *Argania spinosa* (L.) Skeels as case study

Said Moukrim · Said Lahssini · Mouhssine Rhazi · Hicham Mharzi Alaoui · Abdelkader Benabou · Imane Wahby · Mohammed El Madihi · Moustapha Arahou · Laila Rhazi

Published online: 17 May 2018
© Springer Science+Business Media B.V., part of Springer Nature 2018

Correction to: Agroforest Syst

<https://doi.org/10.1007/s10457-018-0232-8>

Unfortunately, in the original publication of the article, the second line arrangement of Table 1 has

been listed incorrectly. The correct version of Table 1 is provided. The original article has also been corrected.

The original article can be found online at
<https://doi.org/10.1007/s10457-018-0232-8>.

S. Moukrim (✉) · A. Benabou · I. Wahby · M. El Madihi · M. Arahou · L. Rhazi
Research center of plant and microbial biotechnology, biodiversity and environment, Faculty of Sciences, Mohammed V University in Rabat, Avenue Ibn-Battouta, B.P. 1014 RP, Rabat, Morocco
e-mail: maildemoukrim@gmail.com

S. Moukrim · H. M. Alaoui · A. Benabou
Water and Forests Department, 3 rue Harroun Errachid–Agdal, Rabat, Morocco

S. Lahssini
National School of Forest Engineer,
Bd My Youssef, BP. 511. Tabriquet, 11000 Salé, Morocco

M. Rhazi
Department of Biology, Faculty of Sciences and Techniques, Moulay Ismail University, BP509, Boutalamine, Errachidia, Morocco

Table 1 Predicted change in suitability areas for argan tree distribution (in square kilometers) by short and middle-term climate

	Ranges of suitable habitat				Potential production	
	Total	Stable	Decrease	Increase	Forage	Argan oil
Actual occurrence of <i>A. spinosa</i>	8712				175,000	4008
Current climate prediction ($R > \frac{1}{2}T$)	39,070				781,000	17,900
Future climate prediction (Averaged GCMs)						
2050						
RCP2.6	28,818 (− 26)	26,726 (68)	12,344 (− 32)	2092 (+ 5)	(− 32)	
RCP4.5	30,944 (− 21)	25,585 (65)	13,486 (− 34)	5359 (+ 14)	(− 34)	
RCP6.0	34,057 (− 31)	26,455 (67)	12,615 (− 32)	7602 (+ 19)	(− 32)	
RCP8.5	38,927 (− 0.4)	23,536 (60)	15,534 (− 40)	15,391 (+ 39)	(− 40)	
2070						
RCP2.6	27,858 (− 29)	25,680 (66)	13,390 (− 34)	2178 (+ 6)	(− 34)	
RCP4.5	40,389 (+ 3)	24,489 (63)	14,582 (− 37)	15,900 (+ 41)	(− 37)	
RCP6.0	26,507 (− 32)	22,046 (56)	17,024 (− 44)	4461 (+ 11)	(− 44)	
RCP8.5	35,798 (− 8)	18,272 (47)	20,798 (− 53)	17,526 (+ 45)	(− 53)	

For the habitat suitability, the percent changes from current climate predictions are provided in parenthesis. For the potential production of argan oil and forage is respectively (in $\times 10^3$ L and $\times 10^3$ FU), and parenthesis provides the percent loss of the potential production in the future. *R* relative habitat suitability; *T* predefined threshold; *GCMs* general circulation models