PREFACE

Yaozhong HU (胡耀忠)

*Department of Mathematical and Statistical Sciences, University of Alberta,
Edmonton, T6G 2G1, Canada*

There are large amount of research papers on stochastic partial differential equations (SPDEs). This volume attempts to collect some recent progresses on some very special topics in this broad field. Our concentration will be the stochastic heat (wave) equations driven by Gaussian noises:

$$\partial_t^\beta u(t, x) = \frac{1}{2} \Delta u(t, x) + \sigma(u) \dot{W}, \quad t \geq 0, \quad x \in \mathbb{R}^d,$$

where ∂_t^β denotes the partial derivative with respect to t ($\beta = 1$ first order, $\beta = 2$ second order), $\Delta = \sum_{k=1}^d \frac{\partial^2}{\partial x_k^2}$ denotes the Laplacian in $\mathbb{R}^d$, and $\dot{W} = \frac{\partial^{d+1}}{\partial t \partial x_1 \cdots \partial x_d} W$ is the Gaussian noise.

In the above, $\{W(t, x) = W(t, x, \omega), t \geq 0, x \in \mathbb{R}^d, \omega \in \Omega\}$ is a Gaussian random field on some probability space $(\Omega, \mathcal{F}, P)$ of mean zero and with some given specific covariance structure:

$$\mathbb{E}(W(s, x)) = 0$$

$$\mathbb{E}[W(s, x)W(t, y)] = q(s, t; x, y), \quad s, t \geq 0, \quad x, y \in \mathbb{R}^d.$$

Some special covariance structures are

$$q_1(s, t; x, y) = \frac{1}{2^d} (s \wedge t) \left(\prod_{k=1}^d [|x_k|^{2H} + |y_k|^{2H} - |x_k - y_k|^{2H}] \right);$$

$$\frac{\partial^{2d+2}}{\partial t \partial t \partial x \partial y} q_2(s, t; x, y) = c |t - s|^{-\beta} \int_{\mathbb{R}^d} e^{\iota \xi(x-y)} \mu(d\xi);$$

for some constant c and some measure (or generalized function) μ , where $\iota = \sqrt{-1}$ is the imaginary number, and $\partial x = \partial x_1 \cdots \partial x_d$. For example, the most studied covariance structure is when $H = 1/2$ in q_1 . In this case, the noise $\dot{W}$ is called space time white noise.

Usually and in the most interesting situation, the derivative $\frac{\partial^{d+1}}{\partial t \partial x_1 \cdots \partial x_d} W$ does not exist. So one can not no longer use the classical PDE method to study this equation for any fixed sample path ω . The probabilistic method is needed.

When $\sigma(u) = u$, the above mentioned equation is one of the simplest nontrivial stochastic partial differential equations, for which one may obtain some sharp properties of the solution. For example, one can obtain the exact moment bound and exact asymptotic behavior of the solution (see the references of the papers collected in this volume); one can also express the exact probability distribution of the solution $u(t, x)$ by using the Tracy-Widom law when the noise is space time white and when the initial condition is the Dirac delta function (in one space dimension). This expression can be used to obtain the sharp asymptotic behavior of the probability density of solution $u(t, x)$ ([1, 2]). See also [4] for the pretty sharp density asymptotics of general Gaussian noises. On the other hand, this equation is relevant to the parabolic Anderson localization, in particular in its discrete form ([5]). It is also related to the KPZ (Mehran Kardar, Giorgio Parisi, and Yi-Cheng Zhang) equation (see [3]), which is the field theory of many surface growth models, such as the Eden model, ballistic deposition, and the SOS model.

In addition to the above equation, we also collected some papers in related fields such as those dealing with the SPDEs arising from the branching processes, stochastic Hamilton-Jacobi equation, McKean-Vlasov equation, and complex Wiener-Itô chaos expansion.

References

- [1] Amir G, Corwin I, Quastel J. Probability distribution of the free energy of the continuum directed random polymer in $1 + 1$ dimensions. *Comm Pure Appl Math*, 2011, **64**(4): 466–537
- [2] Corwin I, Ghosal P. Lower tail of the KPZ equation. *ArXiv* 1802.03273v1, 2018
- [3] Hairer M. Solving the KPZ equation. *Ann of Math*, 2013, **178**(2): 559–664
- [4] Hu Y, Lê K. Asymptotics of the density of parabolic Anderson random fields. *arXiv*:1801.03386v1, 2018
- [5] König W. The parabolic Anderson model. *Random walk in random potential. Pathways in Mathematics*. Birkhäuser/Springer, 2016