

Correction to: Population genetic structure of lumpfish along the Norwegian coast: aquaculture implications

Ólöf Dóra Bartels Jónsdóttir¹ · Julia Schregel² ·
Snorre B. Hagen² · Camilla Tobiassen² ·
Siv Grethe Aarnes² · Albert K. D. Imsland^{1,3}

Published online: 26 October 2017
© Springer International Publishing AG 2017

Correction to: Aquacult Int
<https://doi.org/10.1007/s10499-017-0194-2>

The original version of this article unfortunately contained an error where the figure caption for Figures 3 and 4 was mixed.

The correct caption for those figures should read as follows:

Fig. 3 Relationships between geographical (km shortest waterway) and genetic distance [estimated as $F_{ST} (1-F_{ST})^{-1}$] among pairs of lumpfish samples along the Norwegian coastline. The dotted line shows the results of non-significant ($y = 1E-06x + 0.0152$, $R^2 = 0.0043$, $P > 0.9$) regression between geographic and genetic distance.

Fig. 4 Bayesian cluster analyses performed in STRUCTURE (Pritchard et al. 2000) on lumpfish samples from six sample localities. Each vertical bar represents an individual, whilst colours denote inferred clusters. The y-axis indicates the probability of an individual belonging to a particular cluster.

The online version of the original article can be found at <https://doi.org/10.1007/s10499-017-0194-2>

✉ Albert K. D. Imsland
albert.imsland@akvaplan.niva.no

¹ Akvaplan-niva, Iceland Office, Akralind 4, 201 Kópavogur, Iceland

² Norwegian Institute of Bioeconomy Research, Svanhovd, 9925 Svanvik, Norway

³ Department of Biology, University of Bergen, High Technology Centre, 5020 Bergen, Norway