Correction to: Decline of rare and specialist species across multiple taxonomic groups after grassland intensification and abandonment

Andreas Hilpold¹ · Julia Seeber^{1,2} · Veronika Fontana¹ · Georg Niedrist¹ · Alexander Rief² · Michael Steinwandter¹ · Erich Tasser¹ · Ulrike Tappeiner^{1,2}

Published online: 29 May 2019
© Springer Nature B.V. 2019

Correction to: Biodiversity and Conservation (2018) 27:3729–3744
<https://doi.org/10.1007/s10531-018-1623-x>

In the original publication of the article, Fig. 2 was published without the labels.
The corrected figure and figure legend are provided in this Correction.

Fig. 2 Percentage of rare (left) and specialist species (middle) and total species number (right) in intensively used hay meadows (H), dry pastures (P) and larch forests (L); letters indicate significant differences between habitats based on χ^2 -tests at significance level $p < 0.05$

Andreas Hilpold and Julia Seeber have contributed equally as first authors.

The original article can be found online at <https://doi.org/10.1007/s10531-018-1623-x>.

✉ Andreas Hilpold
Andreas.Hilpold@eurac.edu

¹ Institute for Alpine Environment, Eurac Research, Drususallee 1, 39100 Bolzano, Italy

² Department of Ecology, University of Innsbruck, Technikerstraße 25, 6020 Innsbruck, Austria

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.