



Correction to: Beaver creates early successional hotspots for water beetles

Petri Nummi¹  · Wenfei Liao²  · Juliette van der Schoor^{3,4} · John Loehr⁴ 

Published online: 3 July 2021
© Springer Nature B.V. 2021

Correction to: Biodiversity and Conservation

<https://doi.org/10.1007/s10531-021-02213-8>

In the original publication of the article, the captions of figures 1 and 2 were interchanged. The correct figure captions and figures are given in Figs. 1 and 2.

The original article can be found online at <https://doi.org/10.1007/s10531-021-02213-8>.

✉ Wenfei Liao
wenfei.liao@helsinki.fi

¹ Department of Forest Sciences, University of Helsinki, P.O. Box 27, 00014 Helsinki, Finland

² Faculty of Biological and Environmental Sciences, University of Helsinki, P.O. Box 65, 00014 Helsinki, Finland

³ Department of Applied Biology, HAS University of Applied Sciences, Onderwijsboulevard 221, 5223 DE 's-Hertogenbosch, The Netherlands

⁴ Lammi Biological Station, University of Helsinki, Pääjärventie 320, 16900 Lammi, Finland




Fig. 1 Water beetle species presence in study ponds. A dark square indicates presence. Note that only ponds with beetles shown




Fig. 2 Raw data of water beetle species richness and abundance per trap in each pond type. The box represents the data from Quantile 1 to Quantile 3; the black bar in each box represents the median; the open circles represent outliers. The different letters in the figure represent significant difference. For example, new beaver ponds, labelled with **a**, have significantly more species than old beaver ponds, labelled with **b**

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.