



Correction to: Elevational changes in canopy Collembola community composition are primarily driven by species turnover on Changbai Mountain, northeastern China

Yunga Wu^{1,2} · Zhijing Xie¹ · Zhuoma Wan¹ · Qiao-Qiao Ji³ · Jingjing Yang^{1,4} · Ting-Wen Chen^{2,5} · Donghui Wu^{1,3,6,7} · Stefan Scheu^{2,8}

Published online: 9 December 2023
© The Author(s) 2023

Correction to: Biodiversity and Conservation
<https://doi.org/10.1007/s10531-023-02734-4>

In the original article, the Funding information section was missed out and should have read as below,

This work was supported by the National Natural Science Foundation of China (31861133006), the National Science and Technology Fundamental Resources Investigation Program of China (2018FY100300), Young Scientists Fund of the National Natural Science Foundation of China (42,301,072 to ZX), the China Scholarship Council (CSC No. 202106620038 to YW), the Program of Introducing Talents of Discipline to Universities (No. B16011), and the German

The original article can be found online at <https://doi.org/10.1007/s10531-023-02734-4>.

✉ Ting-Wen Chen
tchen2@gwdg.de

✉ Donghui Wu
wudonghui@iga.ac.cn

¹ Key Laboratory of Vegetation Ecology, Ministry of Education, Northeast Normal University, Changchun 130024, China

² J.F. Blumenbach Institute of Zoology and Anthropology, University of Göttingen, Göttingen, Germany

³ Key Laboratory of Wetland Ecology and Environment, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun 130102, China

⁴ School of Biotechnology, Key Laboratory of Applied Chemistry and Technology in Oilfield, Daqing Normal University, DaqingHeilongjiang 163712, China

⁵ Biology Centre of the Czech Academy of Sciences, Institute of Soil Biology and Biogeochemistry, České Budějovice, Czech Republic

⁶ State Environmental Protection Key Laboratory of Wetland Ecology and Vegetation Restoration, School of Environment, Northeast Normal University, Changchun 130117, China

⁷ Jilin Provincial Key Laboratory of Animal Resource Conservation and Utilization, Northeast Normal University, Changchun 130117, China

⁸ Centre of Biodiversity and Sustainable Land Use, University of Göttingen, Göttingen, Germany

Research Foundation (DFG; SCHE 376/42-1). Open Access funding enabled and organized by Projekt DEAL.

Also, the Tables 1 and 2 have missed grey background in specific cells. The correct version of tables is given in this erratum.

Table 1 Pairwise distance matrix for the abundance-based partitioning of pairwise Bray-Curtis dissimilarity, with spatial turnover measured as balanced variation in abundance between elevations (β_{bal}) (bold) and nestedness measured as the unidirectional abundance gradients between elevations (β_{gra}) (unbold) between Collembola communities of different elevations on Changbai Mountain. The percentage contribution to total beta diversity is given in parentheses

	800 m	1100 m	1400 m	1700 m
800 m		0.095 (13.0%)	0.049 (5.0%)	0.018 (1.8%)
1100 m	0.637 (87.0%)		0.281 (37.8%)	0.280 (41.8%)
1400 m	0.930 (95.0%)	0.463 (62.2%)		0.034 (5.4%)
1700 m	0.972 (98.2%)	0.390 (58.2%)	0.599 (94.6%)	

Table 2 Analysis of similarity (SIMPER) of Collembola families and the three most abundant species contributing to the dissimilarity of Collembola communities between elevations. Av_a and Av_b represent the average abundance of Collembola at the respective two elevations compared. Only families contributing > 10% to the community difference are shown

	Taxa	Contribution (%)	Av_a (ind./sample)	Av_b (ind./sample)	P value
Family					
800 m vs. 1100 m	Entomobryidae	17.75	179.0	252.0	0.033
800 m vs. 1400 m	Tomoceridae	23.67	7.0	245.5	0.001
800 m vs. 1700 m	Sminthuridae	22.29	0.8	271.0	0.001
1100 m vs. 1400 m	Tomoceridae	20.79	1.8	245.5	0.004
1100 m vs. 1700 m	Sminthuridae	17.83	18.8	271.0	0.002
Species					
800 m vs. 1100 m	<i>Entomobrya</i> sp.4	23.31	0	122.0	0.004
	<i>Entomobrya</i> sp.8	17.56	98.8	0	0.001
	<i>Entomobrya</i> sp.1	8.02	51.0	91.0	0.028
800 m vs. 1400 m	Tomoceridae sp.1	24.21	0	242.8	0.001
	Hypogastruridae sp.7	15.95	0.5	404.8	0.001
	Xenylla sp.1	3.84	0	75.0	0.001
800 m vs. 1700 m	<i>Entomobrya</i> sp.4	22.91	0	278.5	0.002
	Sminthuridae sp.1	22.03	0.3	266.8	0.001
	Isotomidae sp.4	5.18	0	69.3	0.001
1100 m vs. 1400 m	Tomoceridae sp.1	20.78	0	242.8	0.003
	Hypogastruridae sp.7	15.06	0.5	404.8	0.031
	Entomobrya sp.1	7.68	91.0	8.0	0.038
1100 m vs. 1700 m	Sminthuridae sp.1	17.55	18.8	266.8	0.002
	<i>Entomobrya</i> sp.1	8.28	91.0	2.5	0.015
	Isotomidae sp.4	4.58	0	69.3	0.005

The original online version of this article was revised.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.