


Soil phosphorus supply controls P nutrition strategies of beech forest ecosystems in Central Europe

F. Lang  · J. Krüger · W. Amelung · S. Willbold · E. Frossard ·
E. K. Bünemann · J. Bauhus · R. Nitschke · E. Kandeler · S. Marhan ·
S. Schulz · F. Bergkemper · M. Schloter · J. Luster · F. Guggisberg ·
K. Kaiser · R. Mikutta · G. Guggenberger · A. Polle · R. Pena ·
J. Prietzel · A. Rodionov · U. Talkner · H. Meesenburg · K. von Wilpert ·
A. Hölscher · H. P. Dietrich · I. Chmara

Received: 6 April 2017 / Accepted: 2 September 2017 / Published online: 9 September 2017
© The Author(s) 2017. This article is an open access publication

Abstract Phosphorus availability may shape plant–microorganism–soil interactions in forest ecosystems. Our aim was to quantify the interactions between soil P availability and P nutrition strategies of European beech (*Fagus sylvatica*) forests. We assumed that plants and microorganisms of P-rich forests carry over mineral-bound P into the biogeochemical P cycle (acquiring strategy). In contrast, P-poor ecosystems establish tight P cycles to sustain their P demand (recycling strategy). We tested if this conceptual

model on supply-controlled P nutrition strategies was consistent with data from five European beech forest ecosystems with different parent materials (geosequence), covering a wide range of total soil P stocks (160–900 g P m⁻²; <1 m depth). We analyzed numerous soil chemical and biological properties. Especially P-rich beech ecosystems accumulated P in topsoil horizons in moderately labile forms. Forest floor turnover rates decreased with decreasing total P stocks (from 1/5 to 1/40 per year) while ratios between organic carbon and organic phosphorus (C:P_{org}) increased from 110 to 984 (A horizons). High proportions of fine-root biomass in forest floors seemed to favor tight P recycling. Phosphorus in fine-root biomass increased relative to microbial P

Responsible Editor: Edith Bai.

Electronic supplementary material The online version of this article (doi:[10.1007/s10533-017-0375-0](https://doi.org/10.1007/s10533-017-0375-0)) contains supplementary material, which is available to authorized users.

F. Lang (✉) · J. Krüger
University of Freiburg, Chair of Soil Ecology,
Bertholdstraße 17, 79085 Freiburg (I Br.), Germany
e-mail: fritzi.lang@bodenkunde.uni-freiburg.de

W. Amelung · A. Rodionov
University of Bonn, Institute of Crop Science and
Resource Conservation, Division Soil Science, Nußallee
13, 53115 Bonn, Germany

S. Willbold
Research Centre Juelich, ZEA-3, Wilhelm-Johnen-Str.,
52428 Jülich, Germany

E. Frossard · E. K. Bünemann
ETH Zurich, Institute of Agricultural Sciences, Eschikon
33, 8315 Lindau, Switzerland

J. Bauhus · R. Nitschke
University of Freiburg, Chair of Silviculture,
Tennenbacherstr. 4, 79085 Freiburg (I. Br.), Germany

E. Kandeler · S. Marhan
University of Hohenheim, Soil Biology, Emil-Wolff-Str.
27, 70593 Stuttgart, Germany

S. Schulz · F. Bergkemper · M. Schloter
Helmholtz Zentrum München, Research Unit for
Comparative Microbiome Analyses, Ingolstädter
Landstraße 1, 85764 Neuherberg, Germany

J. Luster · F. Guggisberg
Biogeochemistry Group, Swiss Federal Institute for
Forest, Snow and Landscape Research WSL,
Zürcherstrasse 111, 8903 Birmensdorf, Switzerland

with decreasing P stocks. Concomitantly, phosphodiesterase activity decreased, which might explain increasing proportions of diester-P remaining in the soil organic matter. With decreasing P supply indicator values for P acquisition decreased and those for recycling increased, implying adjustment of plant–microorganism–soil feedbacks to soil P availability. Intense recycling improves the P use efficiency of beech forests.

Keywords Forest ecosystem nutrition · P geosequence · P acquiring · P-recycling

Introduction

The functioning of terrestrial ecosystems depends on P availability (Achat et al. 2016) but quantitative knowledge regarding its relationship to soil P stocks is missing. Results obtained from studies along chronosequences showed that aboveground and belowground P pools of ecosystems reflect the current state of interactions between P availability of soils and P uptake and usage by plants and microorganisms occurring at these sites (Pearson and Vitousek 2002; Selmants and Hart 2010; Turner and Condron 2013; Galván-Tejada et al. 2014; Wu et al. 2014). These studies suggest that the P contents of the parent material affect the development of ecosystems, especially in the soil compartment, based on different P-related plant–microorganism–soil interactions.

According to Odum’s hypothesis on the nutrition strategies of vegetation, P cycles tend to “tighten” during succession (Odum 1969): young ecosystems are characterized by open P cycles; mature ecosystems establish closed P cycles. The rate of changes in P pools with succession likely depends on the parent material of the soils (Laliberté et al. 2013). Recently, the role of the parent material in the P nutrition of terrestrial ecosystems has thus gained more and more attention (Augusto et al. 2017). Based on their analyses of the P contents of different types of parent material, Porder and Ramachandran (2013) outlined the need for quantifying links between the P content of the parent material and ecosystem P dynamics. Owing to our limited understanding of the underlying processes, it is currently not possible to predict, how P nutrition strategies of late successional forest ecosystems are controlled by the P supply from parent material.

The recent decline in P nutrition of European beech (*Fagus sylvatica*) documented in several publications (e.g., Talkner et al. 2015) has intensified the scientific debate about possible underlying mechanisms (Jonard et al. 2015). So far no consensus has been reached and many open questions remain regarding the P nutrition of beech forests. Natural *F. sylvatica* forests cover a wide range of soils, which mainly formed since the last ice age and developed from various parent materials (Peters 1997). The prevalence of *F. sylvatica* ecosystems in Central Europe enables exploring the P nutrition strategies of plant and microbial communities with different soil P

K. Kaiser · R. Mikutta
Soil Sciences, Martin Luther University Halle-
Wittenberg, Von-Seckendorff-Platz 3,
06120 Halle (Saale), Germany

G. Guggenberger
Institute of Soil Science, Leibniz Universität Hannover,
Herrenhäuser Str. 2, 30419 Hannover, Germany

A. Polle · R. Pena
Department of Forest Botany and Tree Physiology,
Georg-August-Universität Göttingen, Büsgenweg 2,
37077 Göttingen, Germany

J. Prietzel
Technical University of Munich, Research Department
Ecology and Ecosystem Management, Chair of Soil
Science, Emil-Ramann-Straße 2, 85354 Freising,
Germany

U. Talkner · H. Meesenburg
Nordwestdeutsche Forstliche Versuchsanstalt, Grätzelstr.
2, 37079 Göttingen, Germany

K. von Wilpert · A. Hölscher
Forstliche Versuchs- und Versuchsanstalt Baden-
Württemberg, Abteilung Boden und Umwelt,
Wonnhaldestraße 4, 79100 Freiburg, Germany

H. P. Dietrich
Bayerische Landesanstalt für Wald und Forstwirtschaft,
Abteilung Boden und Klima, Hans-Carl-von-Carlowitz-
Platz 1, 85354 Freising, Germany

I. Chmara
Thüringen Forst, Forstlichen Forschungs- und
Kompetenzzentrum Gotha, Jägerstraße 1, 99867 Gotha,
Germany

contents but similar state of forest development. This provides an opportunity to test the conceptual model, that P stocks in parent material control P nutrition of mature forest ecosystems (Lang et al. 2016). According to this model, plant and microbial communities at P-rich sites transfer P from soil minerals into the biogeochemical P cycle. The set of mechanisms involved in this transfer is what we term *P acquiring strategy*. In contrast, tight P cycling is expected at sites poor in P. That means plants and microbes use P from organic sources and minimize P losses from the biogeochemical cycle. The set of mechanisms contributing to this tightening of the P cycle is what we term *P recycling strategy*. The aim of our study was to test if this conceptual model of supply-controlled P nutrition is consistent with analytical data obtained from beech forest ecosystems. Additionally, we addressed the question of *how* P acquisition and recycling changes with changing P supply. First, we analyzed a variety of P-related ecosystem properties of beech forest ecosystems at sites with different parent materials, covering a wide range of different P stocks. Second, we quantified indicator values for the proposed P nutrition strategies and related them to the P soil stocks of the five study sites.

To the best of our knowledge, the study design as well as the diversity of analyses applied is unprecedented: we studied five forest ecosystems dominated by the same tree species but differing in P stocks of soils by a factor of six. In contrast to the concept of chronosequences, the sites represent a geosequence with different P contents in the parent material and are spatially independent. The chemical analyses covered a wide range of P forms, included soil stocks at high depth resolution down to 1 m, as well as P mobilization kinetics. These analyses were combined with analyses of microbiological and root characteristics as well as P concentrations in beech leaves and permitted the solid analysis of soil–plant–microorganism interactions in beech forest ecosystems along the P geosequence. Based on these data and on a novel indicator approach we provide evidence that P nutrition strategies of beech forests adapt continuously to changing P stocks.

Materials and methods

Study sites

We selected five study sites differing in their parent material and supporting 120–140 year old beech forests (Table 1). Methods used for analyzing stand characteristics are summarized in supplementary S1. The parent material ranged from P-rich basaltic rock (Porder and Ramachandran 2013) to P-poor sandy till (Table 1). All sites were in periglacial zones during the last glaciation in central Europe (Fiebig et al. 2011; Geyer and Gwinner 1986; Ergenzinger 1967). Thus, the development of the present soil profiles started at the end of the last ice-age, 10–12,000 years ago (Eberle and Allgaier 2010). All study sites are Level II intensive monitoring plots of the Pan-European International Co-operative Program on assessment and monitoring of air pollution effects on forests (ICP Forests) under UNECE (Lorenz 1995; Vries et al. 2003). Soil and tree properties have been monitored for the last 15–25 years. Four sites (Bad Brückenau, BBR; Mitterfels, MIT; Vessertal, VES; Conventwald, CON) are located at intermediate elevation in the German central and southern uplands; the P-poorest site Lüss (LUE) is located in the North German lowlands.

Sampling design

The characterization of forest stands, vegetation, and litterfall was performed according to the ICP Forest manual at a monitoring plot with an area of 0.25 ha representing the forest stand. The analyses of microbial biomass were performed with samples from soil cores derived from five sampling points distributed within the buffer zone around the monitoring plot (see below). For the other soil analyses, we used soil samples derived from volume based sampling of a complete soil profile performed over an area of 0.25–0.56 m², down to 90–100 cm below the mineral soil surface. Due to the high stone contents of the soils, we decided to use the “quantitative soil pit” (QP) approach developed by Hamburg (1984) and adjusted recently by Vadeboncoeur et al. (2012) to quantify carbon (C) stocks in stony forest soils. The method provides volume-based samples of fine earth material and other constituents, such as roots and stones, of a soil pit. By analyzing a soil volume with a large cross

Table 1 Site and stand characteristics of the study sites Bad Brückenau (BBR), Mitterfels (MIT), Vessertal (VES), Conventwald (CON), Lüss (LUE)

Study site	BBR	MIT	VES	CON	LUE
Soil type (WRB 2015)	Dystric skeletal cambisol (Hyperhumic. Loamic)	Hyperdystric chromic folic cambisol (Humic. Loamic. Nechic)	Hyperdystric skeletal chromic cambisol (Hyperhumic. Loamic)	Hyperdystric skeletal folic cambisol (Hyperhumic. Loamic)	Hyperdystric folic cambisol (Arenic. Loamic. Nechic. Protospodic)
Parent material	Basalt	Paragneiss	Trachyandesite	Paragneiss	Sandy till (Geschiebedecksand)
Humus form (Ad hoc-AG Boden 2005)	Mull-like Moder	Moder	Moder	Mor-like Moder	Mor-like Moder
Texture (topsoil) (WRB 2015)	Silty clay loam	Loam	Loam	Loam	Loamy sand
Texture (subsoil) (WRB 2015)	Loam	Sandy loam	Sandy loam	Sandy loam	Sand
Altitude (m a.s.l.)	809	1023	810	840	115
Mean annual precipitation (mm)	1031	1299	1200	1749	779
Tree species composition (%)	<i>Fagus sylvatica</i> (99) <i>Acer pseudoplatanus</i> (1)	<i>Fagus sylvatica</i> (96) <i>Picea abies</i> (2) <i>Abies alba</i> (2)	<i>Fagus sylvatica</i> (100)	<i>Fagus sylvatica</i> (69) <i>Abies alba</i> (31)	<i>Fagus sylvatica</i> (91) <i>Quercus petraea</i> (9)
Age <i>beeches</i> (a)	137	131	123	132	132
Height <i>beech</i> (mean basal area tree) (m)	26.8	20.8	29.3	27.6	27.3
Diameter at breast height <i>beeches</i> (cm)	36.8	37.6	40.1	39.9	27.5
Number of trees (ha ⁻¹)	335	252	506	312	480
Basal area (m ² ha ⁻¹)	35.6	28.1	31.4	44.0	36.7
Standing volume (m ³ ha ⁻¹)	495	274	550	685	529
Vegetation (pot. nat.)	Hordelymo-Fagetum	Dryopteris-Fagetum	Luzulo-Fagetum	Galio-Fagetum	Luzulo-Fagetum

section representing a large portion of the rooting space of an adult tree, QP sampling allows to obtain a more coherent picture of the system than by analyzing several small soil volumes.

The QPs were established in the buffer zone of the monitoring plots within the Level II sites. Only one QP per study site could be established. The exact position was determined randomly but had to meet the

following criteria: (1) thickness of forest floor and distribution of mineral soil horizons representative for the forest stand (identified based on auger screening of the monitoring site), (2) a minimum distance of at least 3 m to the next trees (diameter at breast height (DBH) >10 cm), and (3) not covered by understory or downed deadwood.

Soil sampling and fractionation

For quantitative pit establishment, a square wooden frame with an interior lateral length of 50 cm (75 cm at site BBR) was prepared and registered optical targets for photogrammetric analysis were fixed to the upper side of the frame. The frame was fixed as reference plane to the soil surface using steel pins. The entire organic layer was cut off alongside the inner edge of the frame with a knife. The individual humus layers (Oi, Oe, Oa) were sampled separately by hand. Roots crossing different layers were cut off at the layer boundary and removed to prevent mixing of material from different layers. The mineral soil sampling followed layers representing single diagnostic horizons. If the thickness of a diagnostic horizon exceeded 5 cm for A or E horizons and 10 cm for B horizons, a new sample layer was started. The procedure was repeated until a maximum sampling depth of 1 m was reached. Following this procedure, we obtained 10 (BBR) to 15 (MIT and VES) different depth layers per site. Detailed information on soil sampling depths at the different sites is given in the supplementary material (S2). All soil material (including rocks and roots) was placed in containers, transported to the laboratory, and air-dried (40 °C). Then, the soil samples were manually separated into the following fractions: fine earth (<2 mm), gravel (2–20 mm), stones (>20 mm), coarse roots (>2 mm), fine roots (<2 mm), and other soil constituents (e.g., wood or seedlings). All fractions were weighed and stored dry, cold, and in the dark for further analysis. Dried fine earth material was used for soil analyses. The volume quantification of the different soil layers was carried out based on photogrammetry (Haas et al. 2016).

For microbiological analyses, five contiguous soil cores (circular distance 2–3 m) were taken and the Oe and Oa horizons removed before samples were pooled together and sieved to 2 mm. This procedure was conducted five times giving five pooled soil samples

per site. Soil samples were stored at 4 °C prior to analysis.

The analyses of the bulk samples from the quantitative pits were conducted in duplicate (soil chemical analyses) or triplicate (analyses of enzyme activity) to account for analytical variability. The mean coefficients of variation of the analyzed soil samples were below 10% and indicate good reproducibility of applied methods. We do not have exact information regarding the spatial heterogeneity of all the analyzed properties across study sites. However, there are clear indications that the results obtained from QP sampling represent the study site properties: (1) QP location was determined based on a soil survey of the study area (see description above). (2) In frame of a geostatistical analysis, we determined citrate extractable P concentrations, carbon to nitrogen (C:N) ratios, and pH values of the forest floor and three soil depth increments at 48 sampling points within 50 × 50 m areas at four of the study sites (LUE, CON, MIT, BBR). The results for the pit samples were mostly within the 68% confidence range of the grid data for the analyzed soil properties and soil horizons (results not shown). (3) We analyzed 10–15 depth intervals at the different sites and outliers would have been identified based on extraordinary discontinuities of analyzed soil properties along the depth gradient. (4) Total P concentrations are assumed to be controlled by the P content of the parent soil material (Turner and Engelbrecht 2011), which is homogeneous within the study sites. Furthermore, published information on the heterogeneity of P concentrations in soil within an area of uniform morphology and geology pointed to only small variation with coefficients of variation less than 10% (e.g., Turner et al. 2012; Chen et al. 2015).

Basic soil chemical characterization

Total contents of soil C and N were measured on ground samples dried at 105 °C using an elemental analyzer (Vario EL cube, Elementar, Germany). Soil pH of air-dried samples (40 °C) was determined in deionized water and in 1 M KCl at a soil-to-solution ratio of 1:2.5 (w/v).

Determination of the cation exchange capacity (CEC) and exchangeable cations was carried out using ammonium acetate at pH 7 and KCl (Hendershot et al. 2008). Concentrations of extracted Ca, Mg, K, and Na were determined by inductively coupled plasma-

optical emission spectroscopy (ICP-OES, Ultima 2, Horiba Jobin–Yvon S.A.S., Longjumeau, France); ammonium in KCl extracts was determined using an automated photometer (SANplus, Skalar Analytical B.V., Breda, The Netherlands). The difference between the CEC and the sum of Ca, Mg, K, and Na is an estimate of H^+ and Al^{3+} occupation of the CEC.

The hot dithionite–citrate–bicarbonate extraction of Fe (Fe_{DCB}), as outlined by Mehra and Jackson (1960), was used to estimate total pedogenic Fe oxide phases. Extraction with NH_4 oxalate at pH 3.0 and 2 h shaking in the dark (Schwertmann 1964) was carried out to estimate Al and Fe in short range-ordered forms and organic complexes (Al_{ox} and Fe_{ox}). The concentrations of extracted Al and Fe were determined by ICP-OES.

Soil phosphorus analyses

Total P, citrate extractable P, and organic P

Contents of total soil phosphorus were determined on ground samples dried at 105 °C after microwave-digestion with 42% HF and H_2O_2 (both: Suprapur®, Merck Millipore, Germany) using ICP-OES (CIROS CDD, Side-On plasma, Spectro, Germany). Organic acid extraction for the quantification of plant-available P in soils has been recommended to simulate organic acid secretion by plant roots (Gerke and Hermann 1992). We analyzed plant-available P by extraction of sieved subsamples with 1% citric acid (50 mM citrate). Similar as described by Hayes et al. (2000), we used a soil-to-solution ratio of 1:10 (w/v), and determined orthophosphate-P concentrations using the ascorbic acid method of Murphy and Riley (1962) as modified by John (1970). Total extractable P was quantified by ICP-OES analyses. Organic P (*P_{org}*) in soil samples was analyzed using the ignition method of Saunders and Williams (1955). Each sample was extracted with or without preceding ignition at 550 °C with 0.5 M H_2SO_4 and the fraction of *P_{org}* was quantified as difference of extracted orthophosphate quantified using the malachite green colorimetric method (Ohno and Zibilske 1991). We used *P_{org}* obtained by the method of Saunders and Williams for calculating the C:*P_{org}* ratios of soil organic matter. Uncertainties related to this approach are discussed in supplementary material S3.

Hedley fractionation

Soil samples were analyzed by sequential extraction according to Hedley and Stewart (1982) as modified by Tiessen and Moir (2008). We used 24 samples in one batch consisting of 23 individual samples including six replicate samples (as random quality check) and one in-house soil standard for quality testing. Each 0.5 g soil was extracted with solutions of increasing extraction strength, starting with distilled water containing an anion exchange resin (Dowex 18, 20–50 mesh, Sigma-Aldrich, Taufkirchen, Germany), followed by 0.5 M $NaHCO_3$; 0.1 M NaOH; 1 M HCl; HCl (37%w/w) and a final acid digestion with 65% HNO_3 and 30% H_2O_2 . Orthophosphate-P concentrations of the different extracts were determined photometrically (Murphy and Riley 1962). We combined the different fractions according their mobility and speciation (organic vs. inorganic P) to describe the following P pools:

Sorbed inorganic P: inorganic P extractable by resin, $NaHCO_3$, and NaOH and *sorbed organic P*: organic P extractable by $NaHCO_3$ and NaOH, *Ca-phosphates*: the P fraction mobilized by 1 M HCl; *stable P*: sum of the P fractions dissolved by HCl (conc) or acid digestion.

Solution-state ^{31}P nuclear magnetic resonance (NMR) spectroscopy

For solution-state ^{31}P -NMR analyses, three soil horizons (Ah horizons, as well as two B horizons from about 30 and 90 cm soil depth) were extracted using 0.25 M NaOH plus 0.05 M Na_2EDTA (1:1/v/v), as described by Cade-Menun (2005). Samples were thereafter centrifuged ($1500\times g$, 20 min), and the remaining supernatant was then split into two halves. One half was lyophilized directly (Thermo Freeze Dryer, Heto PowerDry PL6000). The second half of the supernatant was dialyzed (molecular weight cut off, MWCO, was 14,000; thickness 0.041 mm; Visking, Cellulose, Roth, (Sumann et al. 1998; Amelung et al. 2001). To prepare the samples for NMR spectroscopy, the freeze-dried extracts were resolved by 1 ml *aqua dest.*, 0.5 ml of D_2O , and 10 M NaOH to increase and to standardize the pH for optimal peak separation (Crouse et al. 2000). Samples were centrifuged ($1500\times g$, 20 min) und decanted into NMR tubes.

Spectra were recorded on an NMR spectrometer (Inova 400, Varian, USA) with power-gated proton decoupling at a temperature of 295 K. An acquisition time of 0.7 s, a 30° pulse, and 0.5 s of relaxation delay were used. Chemical shifts of signals were measured in parts per million (ppm) relative to 85% orthophosphoric acid. Approximately 24,576 scans were acquired for each sample. Spectra for the soil samples were recorded with a line broadening of 3.0 Hz. Terminology and interpretations of the spectra followed Cade-Menun (2005, 2015), Bol et al. (2006), Vincent et al. (2013). The chemical shift of the spectra was analyzed as described by Turner (2004). Further details and a discussion of uncertainties are provided in the supplementary material (S4).

Isotopic exchange kinetic

Briefly, a given amount of $\text{H}_3^{33}\text{PO}_4$ was added to a soil water suspension pre-equilibrated (steady-state conditions for P) and the decrease of radioactivity remaining in the solution was measured over time. At the end of the experiment (after 80 min for all samples except for LUE, AE, for which the experiments lasted 420 min because of the very low rate of isotopic exchange), the concentration of orthophosphate in the solution (C_p) was measured after centrifugation for 10 min at 10 000 g and filtration at 0.2 μm using the malachite green colorimetric method (Ohno and Zibilske 1991). Isotopic exchange kinetic (IEK) experiments were detailed by Fardeau (1993, 1996) and Randriamanantsoa et al. (2013). The decrease with time of the ^{33}P added in solution can be described by the following equation (Fardeau 1993):

$$\frac{r_t}{R} = m \times \left(t + m^{\frac{1}{n}}\right)^{-n} + \frac{r_\infty}{R} \quad (1)$$

where r_t and r_∞ (MBq) are the radioactivity remaining in solution after t min of exchange and after an infinite time of exchange, respectively; R (MBq) is the initially added radioactivity; t is the time (in min) elapsed after the radioactivity addition and m and n are soil specific parameters calculated from a non-linear regression between r_t/R and t . The r_∞/R value is estimated as the ratio of the water soluble P expressed in mg P kg^{-1} to the total inorganic P. The amount of

water soluble P is calculated as $10 \times C_p$, with C_p expressed in mg P l^{-1} and 10 being the solution to soil ratio (100 ml:10 g).

As described by Fardeau (1993) the amount of soil isotopically exchangeable P (E_t , expressed in mg P kg^{-1} soil) is calculated based on Eq. 2:

$$E_t = 10 \times C_p \times \left(\frac{R}{r_t}\right) \quad (2)$$

The following variables were calculated for selected samples (Fardeau 1993): m , n , C_p , and the amounts of P isotopically exchangeable within 1 min ($E_{1 \text{ min}}$, mg P kg^{-1} soil), between 1 min and 1 day ($E_{1 \text{ min}-1 \text{ day}}$), between 1 day and 3 months ($E_{1 \text{ day}-3 \text{ months}}$), and the amount of P that cannot be exchanged within 3 months ($E_{>3 \text{ months}}$) by taking the difference between total inorganic P and $E_{3 \text{ months}}$. The samples were analyzed in duplicate.

Fine roots

Fine-root biomass was quantified as described above. Fine roots (<2 mm diameter) were weighed and used for mycorrhizal analyses. All root tips of each weighed sample were inspected under a dissecting microscope (M205 FA; Leica, Wetzlar, Germany) and classified as vital non-mycorrhizal, vital mycorrhizal or dead. The percentage of mycorrhizal colonization was calculated as (n vital mycorrhizal root tips/n vital root tips) \times 100. Vital root tips were distinguished from dead and dry root tips according to the method used by Winkler et al. (2010). Root tip vitality (%) was calculated (n vital root tips/n total root tips) \times 100. For the determination of P concentration in fine roots, fine root material was washed, ground and digested by HNO_3 (65% w/w). Phosphorus concentrations were analyzed by ICP-OES (CIROS CDD, Side-On plasma, Spectro, Germany).

Microorganisms

Quantification of microbial biomass

Soil samples (A horizon) were extracted according to Brankatschk et al. (2011) for estimation of microbial biomass C (C_{mic}), N (N_{mic}), and P (P_{mic}). Measurements of C_{mic} and N_{mic} were performed using the chloroform fumigation-extraction (CFE) method

according to Vance et al. (1987), Joergensen (1996) (k_{EC} 0.45) and Joergensen and Mueller (1996) (k_{EN} 0.54). For determination of microbial biomass P the CFE-method modified from Brookes et al. (1982) (k_{EP} 0.4) was applied. Since C_{mic} , N_{mic} and P_{mic} were intended to be compared from the same extract, 0.01 M calcium chloride was used instead of 0.5 M sodium hydrogen carbonate for inorganic-P extraction. Orthophosphate was quantified as molybdenum blue using commercial tube tests "NANOCOLOR ortho- and total-Phosphate 1" (Macherey–Nagel, Germany).

Determination of acid phosphomonoesterase and phosphodiesterase activities

Acid phosphomonoesterase (EC 3.1.3.2) activity was determined using a modified disodium phenylphosphate method. Briefly, each soil sample (field-moist samples, stored at -20°C and sieved) was split into three subsamples and two controls of 1 g each. Soil suspensions were prepared with 10 ml acetate buffer (pH 5) and 5 ml of 20 mM disodium phenylphosphate (EC 3279-54-7) as substrate solution; in controls, substrate solution was replaced by deionized water. All soil suspensions were incubated at 37°C and continuous shaking (100 rpm) for 3 h. The release of phenol was determined colorimetrically at 614 nm (ELx808, Absorbance Microplate Reader, BioTek Instruments Inc., Winooski, VT, USA), using 2,6-dibromchinone-chlorimide (EC 202-937-2) as coloring reagent (Hoffmann 1968, modified by Öhlinger 1996).

The activity of phosphodiesterase (EC 3.1.4.1) was measured using bis(p-nitrophenyl) phosphate (EC 223-739-2) as substrate and ris(hydroxymethyl)aminomethane as the p-nitrophenol color reagent, according to a modified procedure of Margesin (1996). Each fresh soil sample was split into three subsamples and two controls of each 1 g. Soil suspensions were prepared with 4 ml of 0.05 M Tris(THAM) buffer (pH 8.0) and 1 ml of 5 mM substrate solution; in controls, substrate solution is replaced by deionized water. Soil suspensions were incubated at 37°C for 1 h at continuous shaking (100 rpm). After incubation, 1 ml 0.5 M NaCl solution and 4 ml 0.1 M Tris(THAM) buffer (pH 12.0) were added to each subsample, whereas the controls received additionally 1 ml of the substrate solution.

Soil suspensions were filtered and pipetted into 96-well microplate (PS F transparent 96 well; Greiner Bio-one GmbH, Frickenhausen, Germany). The enzyme activity was measured photometrically at 405 nm on a microplate reader (ELx808, Absorbance Microplate Reader, BioTek Instruments Inc., Winooski, VT, USA).

Leaves and litterfall

P concentrations of beech leaves

Current year beech leaves were sampled in July/August from five trees (9 at LUE). South-facing branches from the upper third of the crown were selected (Rautio et al. 2010). Leaf material was washed, ground and digested by HNO_3 (65%w/w). Phosphorus concentrations were analyzed by ICP-OES. Mean values for leaves from 3–5 years of sampling (between 1995 and 2013, integrating mast years and non-mast years) are presented.

Litterfall

Litterfall at the study sites except VES, where no litterfall sampling took place, was determined according to Pitman et al. (2010) using 10 (12 at LUE) litter collectors distributed among the monitoring plot. Each litter collector had an area of 0.25 m^2 . Litterfall was collected monthly and in periods of strong litterfall bi-weekly. Values presented here represent the sampling period 2013. For litter weight documentation the total mass of litter was determined. P concentrations were only determined for the leaf component of litter fall.

Indicators for P acquisition and P recycling

The most direct approach for testing the hypothesis that soil P stocks control the P nutrition strategy of beech forest ecosystems would be analyzing the P fluxes within the soil and from soil to plants and microorganisms. Some of the relevant P fluxes are hard to quantify (Bol et al. 2016) and not yet available at the study sites. However, the different P nutrition strategies can modify the studied P characteristics of the ecosystems as well as other ecosystem properties. Based on current knowledge and theoretical assumptions we used indicators to assess P nutrition strategies. We defined three indicators for P acquisition and

four indicators for P recycling and analyzed how indicator values were related to the soil P stocks at the study sites. Regarding P recycling we selected properties indicating tight cycling of P between plants and the organic soil P pool. We did not consider plant internal P cycling here although it had been shown to be important for the nutrition of European beech at P poor sites (Netzer et al. 2017). The indicators used are listed and their justifications given in Table 2. The table also informs on the way the indicator values were calculated. In general, indicators representative of acquiring systems address P mobilization from the mineral soil and indicators representative of recycling systems refer to P associated with soil organic matter. To enable the comparison of the different indicators along the geosequence, we normalized all the different indicator values according to Eq. 3:

$$N_{ai} = \frac{I_{ai}}{I_{am}} \quad (3)$$

where N represents the normalized indicator value, the index a represents the indicator addressed, the index i represents the study site, I represents the indicator value, and the index m represents the P-richest site (= BBR) for acquisition indicators and the P-poorest site (= LUE) for recycling indicators.

In addition, we calculated the P uptake efficiency of beech trees according to Marschner et al. (2015) based on Eq. (4):

$$\text{Uptake efficiency} = \frac{\text{Nutrient}_{\text{leaves}}}{\text{Nutrient}_{\text{soil}}} \quad (4)$$

where $\text{Nutrient}_{\text{leaves}}$ is the P stock in beech leaves and $\text{Nutrient}_{\text{soil}}$ the P stock in entire soil profile. The P stock of beech leaves was estimated based on the P concentration of the leaves and the annual litterfall.

Results

Analyses of the soils' solid phase

Variation of basic soil properties along the geosequence

Total P stocks down to 1 m depth of the mineral soil of the geosequence differed by a factor of 5.6 between the poorest site LUE and the richest site BBR (Table 3, supplementary material S5). The P

concentrations in topsoil horizons even differed by a factor of 27 among sites (supplementary material, S5). Nearly all extraction methods confirmed the P gradient for the first meter of the soil profile: $\text{BBR} > \text{MIT} \geq \text{VES} > \text{CON} > \text{LUE}$. The soils of all study sites were acidic (Table 3, supplementary S2) and showed intermediate to high contents of organic C (C_{org}). Soil textures were mostly loam, apart from the P-poorest site LUE with a loamy sand texture. The plant availability of the major nutrients, as indicated by C:N ratio or exchangeable cations, differed strongly among the sites. Although total N contents of the sites were significantly related to P contents ($r^2 = 0.74$; $p < 0.05$), there was still a considerable variation in N/P ratios. The difference in P-related soil properties was larger than that of all the other properties (Table 3).

The depth distribution of P in the soils

Soil P concentrations in the forest floor horizons covered a much smaller range than in the mineral soil horizons (Fig. 1; e.g., Oe horizons: 847–1570 mg kg^{-1} ; A horizons: 196–3265 mg kg^{-1}). Consequently, the portion of P located in the forest floor increased with decreasing P in the mineral soil but was generally rather small (0.4 to 9.2% of total P stock; supplementary material S5). In addition, the mass of forest floor material tended to increase along the P geosequence while its turnover rate decreased continuously (Table 4). In the mineral soil, we observed decreasing concentrations of total P with increasing soil depth at all sites; the decrease was more pronounced at P-rich than at P-poor sites (Fig. 1). While the soil profiles at MIT and LUE indicate vertical translocation of Al and Fe, no such re-location of P could be observed (S6).

Phosphorus fractionation and speciation

Phosphorus concentrations of all Hedley fractions analyzed decreased along the P geosequence with decreasing total soil P stocks (supplementary material S7). Most pronounced with decreasing soil P stocks was the increase in P in the $\text{HCl}_{(\text{conc})}$ and residual P fractions (supplementary material S7). These two fractions contributed up to 20% of the total P at the P-richer sites BBR, MIT and VES. In contrast, they accounted for more than 35% in topsoil horizons and

Table 2 Indicators for P-acquisition and P-recycling applied in this study, method of calculation of indicator values and justification of indicator

Acquiring indicators		
Variable	Calculation of indicator values	Assumed underlying process
P-enrichment in topsoil	P stock in the upper 50% of the soil fine earth mass divided by the total soil P stock (up to 1 m)	Spatial redistribution induced by the P pumping of trees in the long term: root uptake of P in the subsoil, P deposition with litter at the topsoil and adsorption after mineralization. This mechanism was described by Guckland et al. (2009) for Ca. A general discussion on nutrient pumping was provided by Jobbágy and Jackson (2004)
Proportion of non-stable P in the profile	Stock (up to 1 m) of non-stable P. (i.e., sum of Hedley P minus $P_{\text{HCl conc}}$ and P_{residual}) relative to total Hedley P	Chemical redistribution due to biological mobilization of P from primary minerals (Prietz et al. 2016). Nutrient demand had been discussed as the reason for root induced weathering (Kelly et al. 1998)
Phosphate mobilization within 1 min and 1 day	Concentration of isotopically exchangeable P within 1 min and 1 day of topsoil horizons as described in the methods chapter	P mobilization based on physicochemical processes; indicator for P availability (Achat et al. 2016; Bünemann et al. 2016; Oehl et al. 2001)
Recycling indicators		
Variable	Calculation of indicator values	Assumed underlying process
Accumulation of P in the forest floor	P stock in the forest floor related to total P stock (up to 1 m mineral soil depth)	Forest floor pathways as short cut for plant P uptake without passing of P through the fixing mineral soil. P turnover in the forest floor as an emergent property of beech forest ecosystems had been discussed by Talkner et al. (2009)
Concentration of fine-root biomass in the forest floor	Total fine root biomass in the forest floor and in the upper 0–5 cm mineral soil in relation to total fine root biomass (up to 1 m mineral soil depth)	Peak concentrations of fine roots in the forest floor have been assumed to favor tight P cycling in acid temperate forest ecosystems (Wood et al. 1984). Results obtained were much clearer when we added the 0–5 cm increment of the mineral soil. This is in agreement with a smooth transition between Oa and Ah horizons observed at most of the sites
Enrichment of diester-P	Diester-P/monoester P ratio in the topsoil horizon as calculated from NMR spectra	Increased proportions of diester P have been observed in acid soils and were explained by changes in enzyme activity (Turner and Haygarth 2005), and decreased accessibility of diester P for microbial decay due to accumulation within large organic molecules (Turner et al. 2007)
Mean residence time of the forest floor	Mass of the forest floor related to the mass of annual litter fall	Limited decay of soil organic matter enhances tight P recycling by providing forest floor P-pathways for tree nutrition

up to 60% of total P in subsoil horizons at the P-poor sites. The stocks of the P fractions at the different sites also highlighted the increasing proportion of the stable P fractions with decreasing total P stock (Fig. 2). In the subsoil horizons, organic Hedley fractions were greater at sites BBR and MIT than at the P-poor sites CON and LUE (supplementary material S7). The sum of both organic Hedley

fractions corresponded well with the results obtained by the method of Saunders and Williams (1955, see below).

The percentage of organic P of the mineral soils decreased with increasing soil depth and decreasing C_{org} concentration. Regarding the share of P_{org} of the total P stock of the soil profiles, we observed slightly lower contributions of P_{org} at the two sites lower in P

Table 3 Basic characteristics of the soils at the study sites and their variation as indicated by the coefficient of variation among the different sites

Study sites	BBR	MIT	VES	CON	LUE	Coefficient of variation (%)
Element stocks up to 1 m soil depth and forest floor						
P_{tot} (g m^{-2})	904	678	464	231	164	63
N_{tot} (kg m^{-2})	1.3	1.4	1.1	0.8	0.7	27
C_{tot} (kg m^{-2})	18	26	19	18	16	20
Chemical characteristics of A horizons (0–5 cm)						
P_{tot} (mg kg^{-1})	2966	1375	1017	929	195	79
P_{resin} (mg kg^{-1})	116	70	40	24	11	80
C (mg g^{-1})	175	174	126	149	95	24
C:N	16	18	18	22	26	20
C:P	59	127	124	160	493	89
N:P	4	7	7	7	19	67
pH (H_2O)	3.8	3.6	3.4	4.0	3.5	7
pH (KCl)	3.3	3.0	2.8	3.2	2.5	11
CEC meq (kg^{-1})	371	408	197	293	108	45
$(\text{Al} + \text{H})_{\text{ex}}$ meq (kg^{-1})	297	388	184	286	100	44
Texture of the A horizon (0–5 cm)						
Clay (%)	37	24	24	27	6	47
Silt (%)	55	32	46	33	19	37
Sand (%)	8	44	30	40	75	62
Stone content						
Content of stony fragments (profile average) % w/w	78	25	63	69	43	38

P related results are presented in bold letters




Fig. 1 Total P concentrations and depth distributions of P along the analyzed geosequence

(CON and LUE; P_{org} contributes 26% to total P) than at the three sites at the upper end of the P gradient (P_{org} contributed 38–44–38% to total P stock). While there was no clear trend in the percentage of P_{org} in the different horizons along the P geosequence, the C: P_{org}

ratio of the mineral soil increased significantly with decreasing P status, which was most pronounced for the A horizons (Table 5). For the sites BBR, MIT, VES and CON, the C: P_{org} ratio declined with increasing soil depth; at the site LUE the C: P_{org} ratio increased from the Oe to the AE horizon and decreased only beneath.

NMR data showed that the most prominent organic P species were diesters and monoesters (Table 6). The dialyzed EDTA/NaOH extracts contained between 35.6 and 85.2% of monoester-P and between 6.4 and 33.6% of diester-P. Despite the preceding dialysis, the NMR spectra, however, also revealed that the samples contained up to 15% orthophosphate-P. In addition, the samples contained traces of polyphosphate-P (1%, detected in surface soils only) and pyrophosphate-P (6%). Unusually large portions of phosphonate-P were detected, which reached 22% of total signal intensity in the topsoil material at LUE (Table 6). In addition, the spectra of topsoil samples showed increasing proportions of diester-P with decreasing P stock. The

Table 4 Litterfall and P concentrations in leaves, leaf litter and fine roots as well as the turnover rate of the forest floor calculated based on the mass of forest floor and annual litterfall

Study site	BBR	MIT	VES	CON	LUE
P in leaf litter ($\text{g m}^{-2} \text{ a}^{-1}$)	0.229	0.213	n.d.	n.d.	0.156
P leaves ($\text{mg g}^{-1} \text{ d.w.}$)	1.41 (0.21)	1.66 (0.16)	1.64 (0.14)	1.22 (0.14)	1.21 (0.08)
N leaves ($\text{mg g}^{-1} \text{ d.w.}$)	22.9 (2.27)	26.5 (2.27)	24.0 (1.63)	25.2 (0.49)	22.7 (1.39)
P fine roots forest floor (mg g^{-1})	0.96	0.96	0.93	0.80	0.76
P fine roots mineral soil (mean) (mg g^{-1})	0.88 (0.27)	0.82 (0.14)	0.77 (0.13)	0.49 (0.17)	0.54 (0.27)
Mass of forest floor (kg m^{-2})	1.83	5.05	13.98	11.74	12.84
Turnover rate of the forest floor (years^{-1})	1/5	1/14	n.d.	1/36	1/39

Values in parentheses represent standard deviation

n.d. not determined




Fig. 2 Stocks of stable P ($\text{P HCl}_{\text{con}}$, P residual (acid digestion)), P Ca (P 1 M HCl), sorbed inorganic P (P resin, $\text{P}_i \text{ NaHCO}_3$, $\text{P}_i \text{ NaOH}$), and organically bound P ($\text{P}_o \text{ NaHCO}_3$, $\text{P}_o \text{ NaOH}$, P_{tot} (HF) from organic layer) along the analyzed geosequence

increase in diester-P went along with an increase in phosphonate-P, while the proportions of monoester-P declined correspondingly. The ratio of diester-P/mo-
noester-P increased in the same direction: $\text{BBR} < \text{MIT} = \text{VES} < \text{CON} < \text{LUE}$ (Table 6).

Phosphorus mobilization kinetics by isotopic exchange

The concentrations of water-extractable inorganic P (C_P) ranged between 1.3 and 0.01 mg P L^{-1} in the LUE AE and the CON BA horizons, respectively (Table 7). In all soils, C_P values decreased with increasing depth. The loss rate of ^{33}P radioactivity from solution in these laboratory experiments was described using two fitting parameters (Eq. 1). The fitting parameter m followed the same trends as C_P .

The fitting parameter n was between 0.4 and 0.8 in all samples except those from LUE where values were much lower. The amount of P isotopically exchangeable between 1 min and 3 months ($E_{1 \text{ min}-3 \text{ months}}$) in the topsoil horizon decreased in the order $\text{BBR} > \text{MIT} > \text{CON} > \text{VES} > \text{LUE}$, for LUE it was very little. In contrast, the amount of P isotopically exchangeable within 1 min in the AE horizon of LUE was nearly as high as that in the Ah1 horizons of VES and MIT.

Trees and soil microorganisms

Leaves and litter fall

Despite the strong differences in P stocks and availability, above-ground woody biomass was similar among the sites. The P concentrations of beech leaves decreased only slightly with decreasing P stocks in soils (Table 4). Leaf litter P was slightly higher at BBR and MIT than at CON and LUE (Table 4). The decrease in leaf litter P concentrations was much steeper than the decrease in the P concentrations of living leaves (Table 4).

Fine roots

The total fine-root biomass per area slightly increased along the geosequence (BBR 815 mg m^{-2} , MIT 1160 mg m^{-2} , VES 1098 mg m^{-2} , CON 923 mg m^{-2} , LUE 1668 mg m^{-2}). Concentrations of P in fine roots were lower than in leaves (Table 4). Especially at the P-poor sites CON and LUE, root P

Table 5 C and P_{org} concentration and C:P_{org} mass ratio of the forest floor and the mineral soil horizons

Horizon	C _{org} mg g ⁻¹ mg g ⁻¹					Proportion P _{org} % of total P (HF digestion)					C/P _{org} g g ⁻¹					
	BBR	MIT	VES	CON	LUE	BBR	MIT	VES	CON	LUE	BBR	MIT	VES	CON	LUE	
Oi	506	517	510	521	502	56	n.d.	55	n.d.	78	931	n.d.	685	n.d.	792	
Oe	482	503	484	500	387	48	72	67	45	85	634	565	530	1180	536	
Oa	n.p.	349	313	319	463	n.p.	67	83	77	71	n.p.	308	285	330	972	
A	175	174	126	149	96	54	69	66	71	50	110	183	188	226	984	
	97	73	68		18	49	75	66		35	60	89	105		447	
BA transition	78	50	46	80	14	50	66	59	60	37	47	75	85	200	323	
	50				10	51				35	36				225	
B	34	45	46	56	13	50	64	53	61	54	29	70	85	156	132	
		33	40	35			64	51	43			61	78	161		
		34	35	22	10		65	40	40	58		59	81	119	97	
	22	31	29			44	45	35			24	73	74			
		30	23	18	4		46	31	34	54		73	69	119	39	
		25	17				53	24				58	70			
		26	33	12	8	2	41	47	20	34	38	31	73	66	57	27
		27	8				46	1				64	74			
CB transition	11	16		3	2	16	41		28	35	37	47		28	26	
CB transition	b.	b.	6	b.	b.	b.	b.	14	b.	b.	b.	b.	45	b.	b.	

Please consider that sampling depths vary depending on study site. Sampling depths are documented in the supplementary material (S2)

n.p. horizon not present, *n.d.* not determined, *b.* below max. sampling depth

concentrations in mineral soil horizons were much lower than those of fine roots in the forest floor (Table 4). In addition, root biomass was increasingly concentrated in the forest floor and topsoil horizons with decreasing soil P stocks. Only 20% of the total fine-root biomass within the top meter of the soil profile was located in the forest floor and the A horizon at the P-rich site BBR; the respective share increased to nearly 70% at the P-poor site LUE (Fig. 3). The degree of mycorrhization of the vital root tips was 100% across all sites and soil horizons. Root vitality was similar along all study sites, but showed decreasing vitality with increasing soil depth, starting at 60–70% in the forest floor and topsoil horizons and decreasing to about 30–40% in the subsoil horizons.

Microbial biomass

Highest microbial biomass P (P_{mic}) in the topsoil was detected at site MIT (111 µg g⁻¹), followed by CON (104 µg g⁻¹), BBR (100 µg g⁻¹), and VES (79 µg g⁻¹); LUE showed the lowest values (10 µg g⁻¹). Regarding C_{mic} and N_{mic}, a different

order was observed. While LUE again had consistently low values; C_{mic} and N_{mic} reached maxima at the sites CON and BBR (Table 8). The mass ratios of microbial C and P (C_{mic}:P_{mic}) decreased from 20 at LUE to seven at MIT (Table 8). The latter soil likewise showed the lowest ratios of N_{mic}:P_{mic} (0.7) and C_{mic}:N_{mic} (10); highest values were detected at CON (1.3) and LUE (16). In summary, N_{mic}:P_{mic} and C_{mic}:P_{mic} ratios did not strictly follow the P contents of the soils, with the P-richest site of the geosequence (BBR) having larger N_{mic}:P_{mic} and C_{mic}:P_{mic} ratios in the Ah horizon than the P-poorer sites MIT and VES. At all sites, the C:P ratios of microorganisms were much smaller than the C:P ratio of soil organic matter. The imbalances between soil organic matter and microbial ratios increased with decreasing P contents of the soil (Table 8).

Enzyme activities involved in P mineralization

Phosphomonoesterase activity (supplementary S8) was much higher in the forest floor material than in all mineral soil samples. This was also true for

Table 6 Species composition of soil phosphorus forms in EDTA/NaOH extracts after dialysis

Site/ horizon	Organic P species according to chemical shift range (calculated as percent of total signal intensity)						Diester-P/ Monoester- P ratio	NMR Recovery in dialyzed samples	
	Ortho- P (%)	Monoesters (%)	Diesters (%)	Pyrophosphate (%)	Phosphonate (%)	Polyphosphate (%)		Organic species relative to P _{org} (%)	Inorganic species relative to P _i (%)
BBR									
Ah1	16.7	68.8	8.8	1.7	3.3	0.7	0.13	37	10
BA2	9.9	81.4	6.4	0.5	1.8	n.d.	0.08	29	4
BC	4.9	85.2	9.0	0.1	0.8	n.d.	0.11	23	0
MIT									
Ah1	12.4	57.3	12.6	4.9	10.1	2.7	0.22	40	14
Bw3	4.1	85.2	8.6	0.0	2.0	n.d.	0.10	68	3
BC	5.7	85.6	6.8	n.d.	1.9	n.d.	0.08	30	0
VES									
Ah1	8.2	57.0	12.5	3.0	15.2	4.2	0.22	28	19
Bw3	5.9	81.5	10.8	n.d.	1.7	n.d.	0.13	26	3
BC	4.4	83.2	11.6	n.d.	0.8	n.d.	0.14	8	0
CON									
Ah1	13.7	48.5	20.3	4.7	9.7	3.2	0.42	39	22
Bw1	11.9	50.7	33.6	2.0	1.8	n.d.	0.66	35	3
BC	9.0	59.5	30.8	0.7	n.d.	n.d.	0.52	50	2
LUE									
AE	11.3	35.6	23.6	3.8	22.1	3.6	0.66	52	12
Bsw	6.0	69.7	21.3	0.6	2.4	n.d.	0.31	8	1
BC	10.7	69.8	13.4	6.1	n.d.	n.d.	0.19	3	0

For recovery calculations P_{org} as determined according to Saunders and Williams (1955) was used, P_i was calculated as the difference between the concentrations of HF P and P_{org}

n.d. not determined

phosphodiesterase activity (S8). Phosphomonoesterase activities decreased more strongly from forest floor to the mineral soil than phosphodiesterase activities, resulting in smaller monoesterase:diesterase ratios in the mineral soil than in the forest floor, except for LUE (Fig. 4). Phosphomonoesterase activities of soil samples from different soil horizons were closely related to fine-root biomass, with site-specific slopes (data not shown). In contrast, phosphodiesterase activities were related to microbial biomass ($r^2 = 0.81$; $p < 0.05$).

Discussion

Solid phase properties

The range of soil properties covered by the P geosequence

The study sites differed clearly in total P stocks of the mineral soils. The range in P stocks and concentrations in the soils (fine-earth fractions) was as large as the range covered by published chronosequence studies (Table 3, supplementary material S5). The P supply by the soil fine-earth material roughly mirrored the P level of the parent material as described by Porder and Ramachandran 2013. The P contents of the parent

Table 7 Results of isotopic exchange analyses for the Ah horizons: water extractable P (CP), and total inorganic P (Pi)

Site	Horizon	C_P (mg P L ⁻¹)	P_i (mg P kg ⁻¹)	m	n	$E_{1 \text{ min}}$ mg P kg ⁻¹ soil	$E_{1 \text{ min}-1 \text{ day}}$	$E_{1 \text{ day}-3 \text{ months}}$	$E_{>3 \text{ months}}$
BBR	Ah1	0.21	1375.0	0.18	0.59	13.3	533.8	637.9	190.0
	Ah2	0.07	1660.0	0.06	0.57	11.9	504.9	901.6	241.6
	BA1	0.04	1626.0	0.03	0.55	13.6	494.7	862.1	255.6
	BA2	0.07	1383.0	0.03	0.53	21.6	568.7	640.2	152.5
MIT	Ah1	0.27	422.0	0.34	0.76	9.2	339.0	70.9	3.0
	Ah2	0.05	271.0	0.11	0.56	4.7	130.8	114.1	21.5
	BA	0.02	347.0	0.06	0.60	3.0	138.3	158.3	47.4
VES	Ah1	0.70	348.0	0.68	0.40	11.4	108.8	142.9	84.9
	Ah2	0.10	329.0	0.21	0.60	5.2	172.6	132.7	18.6
	BA	0.04	384.0	0.06	0.51	6.8	157.6	174.4	45.1
CON	Ah	0.04	269.0	0.09	0.57	4.0	126.3	118.3	20.4
	BA	0.01	263.0	0.03	0.50	5.0	104.1	119.3	34.7
LUE	AE	1.31	98.0	0.86	0.03	13.2	2.8	2.0	80.0
	E	0.10	78.0	0.97	0.22	1.0	3.9	7.0	66.1

Fitting parameters describing the decrease of radioactivity in the solution with time (m and n), amount of P isotopically exchangeable within 1 min ($E_{1 \text{ min}}$), between 1 min and 1 day ($E_{1 \text{ min}-1 \text{ day}}$), between 1 day and 3 months ($E_{1 \text{ day}-3 \text{ months}}$), and amount of P that cannot be exchanged within 3 months ($E_{>3 \text{ months}}$)




Fig. 3 Three of the soil properties applied as indicators for the recycling strategy against the total P stock of the soil at the five study sites

materials as well as pedogenic processes seem to control the total P stocks of the soils at the different study sites.

Phosphorus associated to Al and Fe oxy(hydr)oxides

Iron and Al oxy(hydr)oxides are important sorbents for inorganic and organic P components. This role was confirmed for the sites BBR, MIT, and CON by P

K-edge XANES analyses (Prietz et al. 2016). At all sites, the amount of oxalate-extractable Al and/or Fe increased—at least slightly—with increasing soil depth, indicating beginning podsolization (Lundström et al. 2000). Podsolization was most obvious at the site LUE. Nevertheless, P was not leached together with Fe and Al but it was incorporated into organic P fractions (supplementary material S7). In consequence, the degree of P saturation of the soil sorbents decreased with increasing soil depth (S6). In contrast, several studies showed translocation of P by podsolization (Turner et al. 2012; Wu et al. 2014; Celi et al. 2013; Wood et al. 1984), especially in cold temperate climate (Väänänen et al. 2008). Also Backnäs et al. (2012) observed P translocation from topsoil to subsoil horizons during podsolization. At their study site in Southern Finnish Lapland, tree growth was much less intense than at our most strongly podsolized site LUE. While the volume of the tree stand at Lapland was 70 m³ ha⁻¹, for instance, it amounted to 529 m³ ha⁻¹ at site LUE. Consequently, the lack of translocation of P at site LUE could be due to stronger biological uptake. Quantifying the true magnitude and thus significance of P fluxes within the

Table 8 Microbial biomass and elemental ratios of microbial biomass as well as P of fine root biomass (P_{fr}) of mineral topsoil horizons (0–20 cm) and stoichiometric imbalances

Site	C_{mic} $\mu\text{g g}^{-1}$	N_{mic}	P_{mic}	P_{fr} g g^{-1}	$C_{mic}:P_{mic}$	$N_{mic}:P_{mic}$	$C_{mic}:N_{mic}$	$C_{Org}:P_{Org}$	$N_{Org}:P_{Org}$	$C_{Org}:N_{Org}$	C:P imbalance
BBR	1223	87	100	5.5	12.2	0.9	14	103	6	15.3	9
MIT	795	82	111	3.9	7.2	0.7	10	198	11	17.5	28
VES	810	64	79	3.3	10.3	0.8	13	226	13	16.8	23
CON	1392	130	104	6.0	13.4	1.3	11	393	18	22.5	30
LUE	192	12	10	1.6	19.8	1.2	16	905	34	25.4	45

between resources (i.e., soil organic matter) and microbes calculated as the mass ratio of $C_{resource}:P_{resource}$ over $C_{mic}:P_{mic}$ according to Mooshammer et al. (2014)




Fig. 4 Ratio of monoesterase activity:diesterase activity at different soil depth and study sites

soil profiles warrants further attention (see also Bol et al., 2016).

Organic P

The percentage of P_{org} in soils has been assumed to be an emergent property representing complex interactions within the ecosystem (Turner and Engelbrecht 2011). The mean values for the proportion of P_{org} found in our study (Table 5) were very close to results observed in other studies, for example for birch forests and tundra soils (mean value 62%; Turner et al. 2004) and for temperate rain forests in New Zealand (mean value 52%; Turner et al. 2007). The proportions of P_{org} in the upper soil profile extended the range of values published for temperate European beech forest ecosystems on loess (44–55%; Talkner et al. 2009; our study: 35–83%), with the largest proportions of P_{org} found at the upper end of the P gradient. Also Ratios of $C:P_{org}$ of the soils continuously increased with decreasing P

stock (Table 5). Both observations suggest more intense usage of P_{org} at the P-poor sites.

Considering the classification of $C:P_{org}$ ratios in Oa and A horizons used in Germany for forest site assessment (Arbeitsgemeinschaft Forsteinrichtung 1996), the site LUE is more P limited than all other sites, which had ratios in the medium (Oa) and moderately small (uppermost A horizon) range. Remarkably, $C:P_{org}$ ratios of soil organic matter of the different study sites converged in subsoil horizons (Table 5). This might be due to specific $C:P$ ratios of dissolved organic matter, which is the most important source of P_{org} in subsoil horizons. This is supported by a recent meta-analysis of soil organic matter stoichiometry in a large number of soils worldwide by Tipping et al. (2016). Their data analysis suggests that relatively P and N-rich organic matter is selectively sorbed by mineral surfaces.

Any intense use of soil P_{org} structures should alter its chemical composition. Indeed, we observed distinct changes in the quality of P_{org} by NMR analyses. Proportions of diester-P and phosphonate-P in organic P forms increased along the decreasing P gradient (Table 6). It has been assumed that diester-P is more readily available to microbial breakdown than monoester-P (Hinedi et al. 1988, Guggenberger et al. 1996). Large proportions of diester-P have also been observed in acid soils of other studies. The controls on the proportion of diester-P are still under debate. Diester-P enrichment has been explained by inclusion of diester-P into large organic molecules (Turner et al. 2007) or by low activity of degrading enzymes (see below).

At the P-poor sites, the proportions of phosphonate-P exceeded by far the levels reported for more fertile soils (e.g., Sumann et al. 1998). Accumulation of

phosphonate-P has been observed for very acidic or water-logged conditions (Condrón et al. 2005). In contrast to our observation, phosphonate-P concentrations showed extremely low concentrations at the late and P-poor states of the Franz-Josef chronosequence (Turner et al. 2007). A relative increase in phosphonate-P along the geosequence (towards LUE) could be favoured by elevated chemical stability of these compounds; particularly biogenic phosphonates were shown to resist chemical degradation (Ternan et al. 1998). Enzymes involved in the degradation of P_{org} can be both, regulated by the concentration of inorganic P (as part of the Pho regulon) or substrate-induced (LysR regulated). While Pho-regulated phosphonases are primarily expressed to meet the organisms' demand for P during phosphate starvation, the latter ones also provide C and N to cells (McGrath et al. 2013). Once easily degradable phosphoester-bonds are hydrolyzed, the more stable organophosphonates remain in the soil. In line with the NMR findings, metagenomic datasets confirmed a significantly elevated microbial potential for one type of phosphonate degrading enzyme (phosphonoacetaldehyde hydrolase) at site LUE (Bergkemper et al. 2016). Probably, the LUE microbial community has adapted to the elevated levels of organophosphonates, thus, showing a higher abundance of the respective genes.

Phosphorus mobilization

The results above mainly address the amount of P in the solid phase. Analysis of the isotopic exchange kinetics provides information on the exchangeability of P between the soil solid phase and the soil solution (Fardeau 1996) and, therefore, on the amount of inorganic P that may diffuse to the absorbing surface of roots or hyphae. The parameter n describes the rate of disappearance of the tracer from the solution after 1 min, thus, accounts for slower physical–chemical reactions (Randriamanantsoa et al. 2013). The parameter m informs on the capacity of the soil to sorb inorganic P (Randriamanantsoa et al. 2013). Small m values indicate a high sorption potential. For the set of soil samples studied, m was significantly correlated to Fe_{CBD} (which is related to the total number of sorption sites) and C_P ($m = 0.47 + 0.39 * C_P - 0.016 * Fe_{CBD}$; $r^2 = 0.67$, $p < 0.01$). The values obtained for m confirm the low sorption capacity for inorganic P of LUE, and the high P sorption capacity of the soils at the other

sites. Low sorption capacity of the soils from site LUE might be explained by the low concentration of iron and aluminum oxides at this site (S6). Interestingly, water soluble P was much higher for the LUE AE horizon than for the other samples. This high value can be explained by the release of microbial P following drying before analysis and subsequent rewetting during analysis (Bünemann et al. 2013). This indicates the high relevance of the microbial pool for the storage of P at the P-poor site LUE, which is in agreement with the metagenomics and NMR data discussed above.

E_{1min} values were in most cases equal or larger than 5 mg P kg^{-1} , the level above which the yield of crop plants does not increase following P application (Gallet et al. 2003). This threshold was, however, found to be valid in soils that had sufficient P exchangeable on the medium term (i.e., between 1 min and 3 months) and were able to sustain significant P desorption flux into the soil solution. This was not the case at LUE. Samples from the LUE AE horizon were characterized by very little P exchangeable between 1 min and 3 months. In agreement with the results from a subsequent soil sampling at sites BBR and LUE (Bünemann et al. 2016), the results obtained here suggest that diffusion of inorganic P into the solution can sustain P availability at most sites but is probably not sufficient to cover the plant demands at LUE. Thus, beech forests at LUE strongly depend on P recycling from soil organic matter.

Response of soil organisms and trees to P stocks of soils

Microbial biomass

The $C_{mic}:N_{mic}:P_{mic}$ ratios detected at the five sites (mean molar ratio: 32:2:1) were comparable to results obtained for tropical mountain forest soils (32:3:1; Tischer et al. 2014) but were lower than documented by a global forest dataset (74:9:1; Cleveland and Liptzin 2007). The $C_{mic}:P_{mic}$ ratios were similar to those found in other beech forests (Joergensen et al. 1995). In addition, Turner and Condrón (2013) observed similar $C_{mic}:P_{mic}$ values based on the same method for P_{mic} analysis. The $C_{mic}:P_{mic}$ values increased with decreasing P status of the soils.

Pools of microbial C, N and P were lowest at site LUE, indicating the general limitation of soil nutrients

and the acidic conditions at this site (Table 8). In a previous study, adaptation of the microbial community to nutrient limitation at the site LUE was shown by a significant higher abundance of oligotrophic taxa such as Acidobacteriales than at BBR (Bergkemper et al. 2016). Surprisingly, P_{mic} reached maximum values at sites MIT and CON, although concentrations of bioavailable P (P_{Resin}) were highest at site BBR (Table 3). In contrast to the $C_{mic}:P_{mic}$ ratio, P_{mic} in the mineral topsoil did not follow the P gradient of soil total P stocks (BBR > MIT \geq VES > CON > LUE). Presumably, microbial growth in BBR is restricted by other soil nutrients or environmental conditions.

While P_{mic} was lowest at LUE, ratios of $C_{mic}:P_{mic}$, $N_{mic}:P_{mic}$ and $C_{mic}:N_{mic}$ were larger than at the other study sites (Table 8). The increased ratio of $C_{mic}:N_{mic}$ indicates a higher fungal contribution to the microbial biomass at site LUE (Mouginot et al. 2014; Fierer et al. 2009). Accordingly, previous data from a metagenomics approach showed a tenfold more fungal sequences at LUE than at BBR, where also the $C_{mic}:N_{mic}$ ratio was much larger (Bergkemper et al. 2016). The large ratio of $C_{mic}:P_{mic}$ in LUE also indicated a strong fungal contribution to the microbial biomass (Heuck et al. 2015). The imbalance of the microbial biomass over the C:P ratio of the soil organic matter ranged from 9 to 45. It increased along the geosequence and was larger at all sites than the average global value for terrestrial ecosystems of 7 (Xu et al. 2013). These results stress the need for P-specific decomposition strategies and for high P uptake efficiency by microorganisms.

Activities of phosphomonoesterase and phosphodiesterase

Phosphatases hydrolyse different phosphate ester bonds to release inorganic P. Therefore, phosphomonoesterases cleave P from monoester forms, such as phospholipids or nucleotides, and phosphodiesterases release P from compounds like nucleic acids (Keller et al. 2012; Paul 2015). Since plants secrete phosphodiesterase only under severe P deficiency, the enzyme is mainly produced by microorganisms (Turner and Haygarth 2005). Phosphodiesterase activity showed more similar patterns among the five soil profiles than the phosphomonoesterase activity (data not shown). However, phosphodiesterase activity was much lower than phosphomonoesterase activity, as already noted

by Turner and Haygarth (2005) for different pasture soils. The absolute activities of the two enzymes were consistent with the species composition of P_{org} forms in the NMR spectra, with a range of 7–34% for diester-P and a range of 36–86% for monoester-P (Table 6). The most striking result was that the LUE site was characterized by high concentrations of diester-P (Table 6) but low phosphodiesterase activities, resulting in large ratio of phosphomonoesterase to phosphodiesterase activity (Fig. 4). This result can be explained by low enzyme production of microorganisms at this site and/or low enzyme stability at low pH values at the LUE site (Table 3; Turner and Haygarth 2005). Limited enzyme concentrations were found as a main reason for recalcitrance of phosphodiesterases in different soils (Jarosch 2016). The retarded hydrolysis of diesters might cause the relative large concentrations of diester P at the LUE site. The large C: P_{org} ratio at the site LUE is in agreement with relative enrichment of diester compounds due to limited enzyme concentrations. Finally, retarded hydrolysis of diesters might also be the reason for the thick forest floor forming at the P-poor sites and the low turnover of organic matter (Table 4).

Leaves

According to Mellert and Göttlein (2012), the observed range of leaf P concentrations (1.2–1.7 mg g⁻¹) are in the “normal range” of European beech. Only beech leaves at sites CON and LUE were in the concentration range of latent P deficiency, with P leaf concentrations of 1.1–1.2 mg g⁻¹. With N:P ratios of 21 and 19 of leaves, respectively, these sites were also close to or even above the critical N:P ratio of 18.9 given for beech leaves (Talkner et al. 2015; Mellert and Göttlein 2012). The increasing difference between the P concentrations of living leaves and the P concentration of leaf litter showed an increasing P re-sorption by trees from senescing leaves with decreasing P contents of the soils. Likely, high P re-sorption is an adaptation of European beech to low P supply (Hofmann et al. 2016; Schmidt et al. 2016; Netzer et al. 2017). In summary, neither standing biomass (Table 1) nor the amount of litter produced (Table 4) seem to be related to the gradient soil P contents. However, based on leaf nutrient concentrations, the sites CON and LUE seem to approach P limitation.

Roots

We observed a strong concentration of root biomass in topsoil horizons at the P-poor sites (Fig. 3), while roots penetrated subsoils especially at P-rich sites (fine root biomass at 80 cm soil depth: BBR 38 mg m⁻²; LUE 2.7 mg m⁻²). Several studies provide evidence that low P concentrations of soils favour the growth of fine roots of different tree species (e.g., Ericsson and Ingestad 1988 for birch, Schneider et al. 2001 for oak). Less information is available on the controls of the depth distribution of the fine root biomass with high depth resolution. Despite large differences in soil P availability, the P concentrations in fine roots showed only a moderate decline with decreasing soil P (Zavišić et al. 2016). Relative differences in fine-root P concentrations at the sites were still bigger than differences in leaf concentrations (Table 4). Obviously, the P supply to roots was facilitated by mycorrhizal fungi, since all vital root tips were completely enwrapped by the fungal mantles. This is in line with observations from other P-poor sites (Schneider et al. 2001). Only in deep soil horizons, the growth of roots seemed to be P-limited, as indicated by very low P concentrations in roots (0.3 mg g⁻¹). Very low P concentrations in coarse roots have also been observed at the P-poorest site LUE (Yang et al. 2016). This finding is well in line with those from isotope exchange kinetics, showing that particularly at site LUE there was no release of mineral-bound inorganic P into soil solution. Thus, roots are limited to the surface layers with elevated microbial activity and P mobilization from organic matter. Of the analyzed biological traits, root distribution between topsoil and subsoil compartments seemed most sensitive to the P status of soils. The ratio of P in fine roots to P_{mic} of the topsoil horizons (0–20 cm) tended to increase with decreasing P status of the soils (BBR: 0.05–LUE: 0.16). This suggests that beech fine roots and/or associated hyphae are more effective at maintaining P concentrations than microorganisms with decreasing P availability. However, it is not clear whether this is simply a result of more effective uptake or other mechanisms, such as uptake from the deeper subsoil.

P nutrition strategies

We defined ecosystem traits that indicate the intensity of P acquisition (acquiring indicators) and the

intensity of P recycling (recycling indicators) (Table 2). We are aware that in ecosystems specific trait values might be achieved by different paths. Thus, our indicator approach provides no concluding proof for underlying mechanisms. Yet, it enables a consistency test for the supply-controlled nutrition strategies of beech forest ecosystems. The indicator values allow for relating nutrition strategies in a quantitative way to the P stocks of soils, thereby providing novel information on how the adjustment of nutrition strategies to changes in P stocks of forest soils occurs.

We assume that ecosystems with P *acquiring strategies* are characterised by spatial (Jobbágy and Jackson 2004) and chemical (Prietz et al. 2016) redistribution of P in soils, for example through P uptake in deeper horizons and fixation of the P mobilized during litter decay by topsoil minerals. Based on this assumption, we identified the following acquiring indicators: (1) P enrichment in topsoil horizons, (2) share of non-stable P in the profile, and (3) kinetics of phosphate mobilization (Table 2).

All the three acquiring indicators highlight more intense spatial and chemical redistribution of P in P-rich than in P-poor soils. We generally observed larger P concentrations in topsoil layers than in subsoil layers (Fig. 1). Published data on P in temperate forest soils confirm this observation (e.g., Talkner et al. 2009; Jobbágy and Jackson 2004). At our study sites, the differences between topsoil and subsoil concentrations were more pronounced at P-rich than at P-poor sites. At sites with large mineral P concentrations, the C investment for establishing and maintaining a deep root system might pay off through gains in nutrient uptake. At P-poor sites, these costs are too high given the low amount of P in the subsoil horizons. The actual depth distribution of fine-root biomass is in line with this explanation, although this does not exclude the possibility that at least some roots also acquire P from deeper depths at P-poor sites. The depth distribution of P_{org} obtained from Hedley fractionation additionally suggests increased P acquisition from subsoil material at P-rich sites. The proportion of P_{org} in the subsoil horizons is higher at P-rich than at P-poor sites (supplementary material S7), which might reflect intense root foraging for P and fine-root turnover in the subsoil horizons of P-rich sites (Cross and Schlesinger 1995). Hedley fractionation results indicate that intense P acquisition occurs preferentially at P-rich sites: In general, the proportion of residual P

seems to be greater at sites low in P than at those having large P stocks (supplementary material S7, see above). This observation corresponds to the P speciation of soil samples by XANES by Prietzel et al. (2016). They observed that P speciation in soils from 10 different forest sites in Germany is strongly related to P cycling in these soils. However, the increasing proportion of stable P with decreasing P stocks might also be explained by increasing fixation of organic and inorganic P by mineral sorbents (Velásquez et al. 2016). Yet the P-poorest site LUE is characterized by low contents of Fe and Al oxyhydroxides. Thus, fixation by mineral sorbents should be of lower relevance than at the other sites. Mobilization kinetics has often been assumed to indicate plant availability, and P mobilization determined by isotopic exchange has been shown to be closely related to plant uptake (Achat et al. 2016). In line with this, P exchangeable within one day from topsoil material was closely related to the P status of the study sites. The rate of P mobilization might be both, the reason for, but also the consequence of P acquisition from mineral material.

The four *recycling indicators* identified were remarkably closely related to the P stocks of the different geosequence sites: (1) accumulation of P in the forest floor (r^2 0.94) (2) fine-root biomass accumulation in the forest floor (r^2 0.95), (3) residence time of the forest floor (r^2 0.99), and (4) contribution of diester-P to total P in A horizons (r^2 0.89; Fig. 3). Also, all the four recycling indicators averaged in the normalized form for aggregated illustration (Fig. 5) confirm decreasing intensity of P recycling with




Fig. 5 Means of normalized indicator values as calculated according to Eq. (3) for the different sites. Bars represent the standard error of normalized values

increasing P supply (r^2 0.96, $p < 0.01$, for the linear regression of recycling indicator values vs. P soil stocks). Thus, most of P seemingly is contained in the surface organic-rich layers at P-poor sites. Slow turnover and high rooting intensity may foster tight P recycling (Fig. 3). Beyond that, we observed an unexpected close, linear, and positive correlation between the diester-P/monoester-P ratio of mineral topsoil horizons and the P stock at a given site (Fig. 3). Despite the uncertainties in the interpretation of NMR data mentioned above, some authors suggested the diester-P/monoester-P ratio as indicator of microbial P transformation (e.g., Turrión et al. 2000). The increase in the diester-P/monoester-P ratio can be seen as a biomarker pointing at increased microbial recycling of organic P with increasing deficiency in overall P supply. This supports our hypothesis that there is a shift to P recycling systems in forest surface soils under extreme P limitations. High phosphomonoesterase-to-diesterase ratios at site LUE likely maintain those chemical signatures.

In summary, the indicator values obtained confirm that European beech forests at silicate sites are characterized by different P nutrition strategies, depending on the P stocks of the soils. Our results imply that with decreasing P contents by the mineral soil, the intensity of recycling processes increases continuously whereas the intensity of acquisition processes decreases. Future work needs to analyze how these different strategies might be impacted by perturbations or disturbances, such as chronic N deposition or climatic extremes. Our results support ideas, which help exemplify the use of such indicators: Interestingly, the peak of biological activity along the P gradient does not seem to be at site BBR (highest P stock) but at sites MIT and VES (intermediate P stocks). These two sites showed the highest P leaf concentrations (Table 4), the lowest $C_{mic}:P_{mic}$ values (Table 8), the largest ratio of P_{org} to total P, and the highest C stocks in soils (Table 3) among the study sites. Under intermediate conditions, P nutrition of vegetation and microbes might be best owing to a mixture of recycling and acquisition strategies.

Ecosystem properties responding to P nutrition strategies

As a consequence of the close relationship between P contents and P recycling indicators, as well as of the




Fig. 6 Phosphorus uptake efficiency as related to the P recycling indicator value determined for the five studied forests

small variation of P leaf concentrations, intense P recycling seems to improve the P use efficiency of P-poor beech forest ecosystems (Fig. 6). The correlation indicates that tight recycling resulting from “well-adjusted” P mobilisation and P uptake by different organisms, thus, also improves the P supply for trees. This, in turn, implies that plant and microbial communities as a whole are adapted to specific environmental conditions, and confirms the concept of ecosystem nutrition (Lang et al. 2016). The results stress that also other ecosystem properties not directly related to P are interacting with P nutrition strategies. Tight cycling appears to be linked to intense P mineralisation, root growth, and P uptake in the forest floor. Particularly, the establishment of the forest floor tightens P cycling and acts as kind of “funnel” to provide available P where required. The formation of the forest floor, in turn, could be facilitated by the depletion in P of the soil organic matter and the P limitation of those organisms inducing depolymerisation of diester-P molecules in the mineral soil. Also, anthropogenic impacts on forest soils, as for example increased N input or climate change and increasing temperatures will affect these interactions and might, thus, disturb the P nutrition strategies of P-poor ecosystems.

Conclusion

We selected a novel P geosequence approach on silicate material to unravel the response of European beech

forests to different soil P stocks. Our results are in line with the hypothesis of a tightening of P cycles with decreasing soil P availability and they provide evidence for the existence of adaptation processes at the ecosystem level. In P-rich but acid soils, plants and soil organisms mobilize P from primary and secondary minerals. In P-poor soils, roots and fungi seem to sustain their P demand more successfully than bacteria, mainly from the forest floor and soil horizons rich in organic matter. In turn, this influences the quality of organic P compounds in soils, which underwent pronounced changes along the P geosequence. Our indicators for P recycling and P acquisition pointed at a continuous change in nutrition strategies with changing P supply. However, elevated proportions of organic P, smallest C:P ratios in microbial biomass, and largest P concentrations in beech leaves were observed at sites of intermediate P stocks, where the combination of different nutrition strategies might favor P uptake. Overall, our results thus underline the high adaptability of beech forest ecosystems to changing P supply. We conclude that within the range of P stocks analyzed, P deficiency in beech forest ecosystems is unlikely the result of a low P supply per se. More likely, sufficient P nutrition depends on supply-specific plant–microorganism–soil interactions.

Acknowledgements This project was carried out in the framework of the priority programme 1685 “Ecosystem Nutrition: Forest Strategies for limited Phosphorus Resources” funded by the DFG (DFG LA 1398/13-1, DFG AM 134/18-1, DFG BA 2821-13-1, DFG KA 1590/12-1, DFG SCHL 446/20-1, DFG KA 1673/9-1, DFG MI 1377/7-1, DFG PO 362/22-1, DFG PR 534/6-1) in cooperation with the Swiss National Foundation (SNF) (Project No. 149130 and Project No. 149138). We thank Dr. Manfred Martin (Landesamt für Geologie, Rohstoffe und Bergbau, Freiburg) for XRF spectroscopy analyses of stony fragments.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Achat DL, Pousse N, Nicolas M, Brédoire F, Augusto L (2016) Soil properties controlling inorganic phosphorus

- availability. General results from a national forest network and a global compilation of the literature. *Biogeochemistry* 127(2–3):255–272. doi:[10.1007/s10533-015-0178-0](https://doi.org/10.1007/s10533-015-0178-0)
- Ad hoc-Arbeitsgruppe Boden (2005) *Bodenkundliche Kartieranleitung*. Ed.: Bundesanstalt für Geowissenschaften und Rohstoffe in Zusammenarbeit mit den Staatlichen Geologischen Diensten. 5th edition. Hannover. ISBN 3-510-95920-5
- Augusto L, Achat DL, Jonard M, Vidal D, Ringeval B (2017) Soil parent material—a major driver of plant nutrient limitations in terrestrial ecosystems. *Glob Change Biol*. doi:[10.1111/gcb.13691](https://doi.org/10.1111/gcb.13691)
- Amelung W, Rodionov A, Urusevskaja I, Haumaier L, Zech W (2001) Forms of organic phosphorus in zonal steppe soils of Russia assessed by ^{31}P NMR. *Geoderma* 103(3–4):335–350. doi:[10.1016/S0016-7061\(01\)00047-7](https://doi.org/10.1016/S0016-7061(01)00047-7)
- Arbeitsgemeinschaft Forsteinrichtung (1996) *Forstliche Standortsaufnahme. Begriffe, Definitionen, Einteilungen, Kennzeichnungen, Erläuterungen*, 5. Aufl. IHW-Verl., Eching bei München
- Backnäs S, Laine-Kaulio H, Kløve B (2012) Phosphorus forms and related soil chemistry in preferential flowpaths and the soil matrix of a forested podzolic till soil profile. *Geoderma* 189–190:50–64. doi:[10.1016/j.geoderma.2012.04.016](https://doi.org/10.1016/j.geoderma.2012.04.016)
- Bergkemper F, Schöler A, Engel M, Lang F, Krüger J, Schloter M, Schulz S (2016) Phosphorus depletion in forest soils shapes bacterial communities towards phosphorus recycling systems. *Environ Microbiol* 18(6):1988–2000. doi:[10.1111/1462-2920.13188](https://doi.org/10.1111/1462-2920.13188)
- Bol R, Amelung W, Haumaier L (2006) Phosphorus-31—nuclear magnetic—resonance spectroscopy to trace organic dung phosphorus in a temperate grassland soil. *J Plant Nutr Soil Sci* 169(1):69–75. doi:[10.1002/jpln.200521771](https://doi.org/10.1002/jpln.200521771)
- Bol R, Julich D, Bröddlin D, Siemens J, Kaiser K, Dippold MA, Spielvogel S, Zilla T, Mewes D, von Blanckenburg F, Puhlmann H, Holzmann S, Weiler M, Amelung W, Lang F, Kuz'yakov Y, Feger K, Gottselig N, Klumpp E, Missong A, Winkelmann C, Uhlir D, Sohr J, von Wilpert K, Wu B, Hagedorn F (2016) Dissolved and colloidal phosphorus fluxes in forest ecosystems—an almost blind spot in ecosystem research. *J Plant Nutr Soil Sci*. doi:[10.1002/jpln.201600079](https://doi.org/10.1002/jpln.201600079)
- Brankatschk R, Töwe S, Kleineidam K, Schloter M, Zeyer J (2011) Abundances and potential activities of nitrogen cycling microbial communities along a chronosequence of a glacier forefield. *ISME J* 5(6):1025–1037. doi:[10.1038/ismej.2010.184](https://doi.org/10.1038/ismej.2010.184)
- Brookes PC, Powelson DS, Jenkinson DS (1982) Measurement of microbial biomass phosphorus in soil. *Soil Biol Biochem* 14(4):319–329. doi:[10.1016/0038-0717\(82\)90001-3](https://doi.org/10.1016/0038-0717(82)90001-3)
- Bünemann EK, Augstburger S, Frossard E (2016) Dominance of either physicochemical or biological phosphorus cycling processes in temperate forest soils of contrasting phosphate availability. *Soil Biol Biochem* 101:85–95. doi:[10.1016/j.soilbio.2016.07.005](https://doi.org/10.1016/j.soilbio.2016.07.005)
- Bünemann EK, Keller B, Hoop D, Jud K, Boivin P, Frossard E (2013) Increased availability of phosphorus after drying and rewetting of a grassland soil. Processes and plant use. *Plant Soil* 370(1–2):511–526. doi:[10.1007/s11104-013-1651-y](https://doi.org/10.1007/s11104-013-1651-y)
- Cade-Menun BJ (2005) Characterizing phosphorus in environmental and agricultural samples by ^{31}P nuclear magnetic resonance spectroscopy. *Talanta* 66(2):359–371. doi:[10.1016/j.talanta.2004.12.024](https://doi.org/10.1016/j.talanta.2004.12.024)
- Cade-Menun BJ (2015) Improved peak identification in ^{31}P -NMR spectra of environmental samples with a standardized method and peak library. *Geoderma* 257–258:102–114. doi:[10.1016/j.geoderma.2014.12.016](https://doi.org/10.1016/j.geoderma.2014.12.016)
- Celi L, Cerli C, Turner BL, Santoni S, Bonifacio E (2013) Biogeochemical cycling of soil phosphorus during natural revegetation of *Pinus sylvestris* on disused sand quarries in Northwestern Russia. *Plant Soil* 367(1–2):121–134. doi:[10.1007/s11104-013-1627-y](https://doi.org/10.1007/s11104-013-1627-y)
- Chen CR, Hou EQ, Condron LM, Bacon G, Esfandbod M, Olley J, Turner BL (2015) Soil phosphorus fractionation and nutrient dynamics along the Cooloola coastal dune chronosequence, southern Queensland, Australia. *Geoderma* 257–258:4–13. doi:[10.1016/j.geoderma.2015.04.027](https://doi.org/10.1016/j.geoderma.2015.04.027)
- Cleveland CC, Liptzin D (2007) C: N: p stoichiometry in soil: is there a “Redfield ratio” for the microbial biomass? *Biogeochemistry* 85(3):235–252. doi:[10.1007/s10533-007-9132-0](https://doi.org/10.1007/s10533-007-9132-0)
- Condron LM, Turner BL, Cade-Menun BJ, Sims JT, Sharpley AN (2005) Chemistry and dynamics of soil organic phosphorus. *Phosphorus* 87–121
- Cross A, Schlesinger W (1995) A literature review and evaluation of the Hedley fractionation: applications to the biogeochemical cycle of soil phosphorus in natural ecosystems. *Geoderma* 64:197–214. doi:[10.1016/0016-7061\(94\)00023-4](https://doi.org/10.1016/0016-7061(94)00023-4)
- Crouse DA, Sierzputowska-Gracz H, Mikkelsen RL (2000) Optimization of sample pH and temperature for phosphorus-31 nuclear magnetic resonance spectroscopy of poultry manure extracts. *Commun Soil Sci Plant Anal* 31(1–2):229–240. doi:[10.1080/00103620009370432](https://doi.org/10.1080/00103620009370432)
- de Vries W, Reinds G, Vel E (2003) Intensive monitoring of forest ecosystems in Europe. *For Ecol Manage* 174(1–3):97–115. doi:[10.1016/S0378-1127\(02\)00030-0](https://doi.org/10.1016/S0378-1127(02)00030-0)
- Eberle J, Allgaier B (2010) *Deutschlands Süden vom Erdmittelalter zur Gegenwart*, 2. Aufl. Spektrum-Sachbuch. Spektrum Akad. Verl., Heidelberg, Neckar
- Ergenzinger P (1967) Die eiszeitliche Vergletscherung des Bayerischen Waldes. *Einzeitalter & Gegenwart* 18(1):152–168
- Ericsson T, Ingestad T (1988) Nutrient and growth of birch seedlings at varied relative phosphorus addition rates. *Plant Physiol* 72:227–235. doi:[10.1111/j.1399-3054.1988.tb05827.x](https://doi.org/10.1111/j.1399-3054.1988.tb05827.x)
- Fardeau JC (1993) Available soil phosphate: its representation by a functional multiple compartment model [conceptual model, pool, buffering capacity]. *Agronomie* 13:317–331
- Fardeau JC (1996) Dynamics of phosphate in soils. An isotopic outlook. *Fertil Res* 45:91–100
- Fiebig M, Ellwanger D, Doppler G (2011) Pleistocene glaciations of Southern Germany. In: Ehlers J, Gibbard PL, Hughes PD (eds) *Quaternary glaciations— extent and chronology: a closer look*, vol 15. Elsevier, Amsterdam, pp 163–173
- Fierer N, Strickland MS, Liptzin D, Bradford MA, Cleveland CC (2009) Global patterns in belowground communities.

- Ecol Lett 12(11):1238–1249. doi:[10.1111/j.1461-0248.2009.01360.x](https://doi.org/10.1111/j.1461-0248.2009.01360.x)
- Gallet A, Flisch R, Ryser J, Frossard E, Sinaj S (2003) Effect of phosphate fertilization on crop yield and soil phosphorus status. *J Plant Nutr Soil Sci* 166(5):568–578. doi:[10.1002/jpln.200321081](https://doi.org/10.1002/jpln.200321081)
- Galván-Tejada NC, Peña-Ramírez V, Mora-Palomino L, Siebe C (2014) Soil P fractions in a volcanic soil chronosequence of Central Mexico and their relationship to foliar P in pine trees. *J Plant Nutr Soil Sci* 177(5):792–802. doi:[10.1002/jpln.201300653](https://doi.org/10.1002/jpln.201300653)
- Gerke J, Hermann R (1992) Adsorption of orthophosphate to humic-Fe-complexes and to amorphous Fe-oxide. *Z Pflanzenernaehr. Bodenk* 155(3):233–236. doi:[10.1002/jpln.19921550313](https://doi.org/10.1002/jpln.19921550313)
- Geyer OF, Gwinner MP (1986) *Geologie von Baden-Württemberg*. Mit 26 Tab, 3., völlig Neubearb. Aufl. Schweizerbart, Stuttgart
- Guckland A, Jacob M, Flessa H, Thomas FM, Leuschner C (2009) Acidity, nutrient stocks, and organic-matter content in soils of a temperate deciduous forest with different abundance of European beech (*Fagus sylvatica* L.). *J Plant Nutr Soil Sci* 172(4):500–511. doi:[10.1002/jpln.200800072](https://doi.org/10.1002/jpln.200800072)
- Guggenberger G, Haumaier L, Zech W, Thomas RJ (1996) Assessing the organic phosphorus status of an Oxisol under tropical pastures following native savanna using ³¹P NMR spectroscopy. *Biol Fertil Soils* 23(3):332–339. doi:[10.1007/BF00335963](https://doi.org/10.1007/BF00335963)
- Haas J, Hagge Ellhöft K, Schack-Kirchner H, Lang F (2016) Using photogrammetry to assess rutting caused by a forwarder—A comparison of different tires and bogie tracks. *Soil Tillage Res* 163:14–20. doi:[10.1016/j.still.2016.04.008](https://doi.org/10.1016/j.still.2016.04.008)
- Hamburg SP (1984) Effects of forest growth on soil nitrogen and organic matter pools following release from subsistence agriculture. In: Stone EL (ed) *Forest soils and treatment impacts*. University of Tennessee, Knoxville, pp 145–158
- Hayes JE, Richardson AE, Simpson RJ (2000) Components of organic phosphorus in soil extracts that are hydrolysed by phytase and acid phosphatase. *Biol Fertil Soils* 32(4):279–286. doi:[10.1007/s003740000249](https://doi.org/10.1007/s003740000249)
- Hedley MJ, Stewart J (1982) Method to measure microbial phosphate in soils. *Soil Biol Biochem* 14(4):377–385. doi:[10.1016/0038-0717\(82\)90009-8](https://doi.org/10.1016/0038-0717(82)90009-8)
- Hendershot WH, Lalonde H, Duquette M (2008) Ion exchange and exchangeable cations. In: Carter MR, Gregorich EG (eds) *Soil sampling and methods of analysis*, vol 2. CRC Press, Boca Raton, pp 197–206
- Heuck C, Weig A, Spohn M (2015) Soil microbial biomass C: N: P stoichiometry and microbial use of organic phosphorus. *Soil Biol Biochem* 85:119–129. doi:[10.1016/j.soilbio.2015.02.029](https://doi.org/10.1016/j.soilbio.2015.02.029)
- Hinedi ZR, Chang AC, Lee RWK (1988) Mineralization of phosphorus in sludge-amended soils monitored by phosphorus-31-nuclear magnetic resonance Spectroscopy. *Soil Sci Soc Am J* 52(6):1593. doi:[10.2136/sssaj1988.03615995005200060014x](https://doi.org/10.2136/sssaj1988.03615995005200060014x)
- Hoffmann G (1968) Eine photometrische Methode zur Bestimmung der Phosphatase-Aktivität in Böden. *Z. Pflanzenernaehr. Bodenk* 118(3):161–172. doi:[10.1002/jpln.19681180303](https://doi.org/10.1002/jpln.19681180303)
- Hofmann K, Heuck C, Spohn M (2016) Phosphorus resorption by young beech trees and soil phosphatase activity as dependent on phosphorus availability. *Oecologia* 181(2):369–379. doi:[10.1007/s00442-016-3581-x](https://doi.org/10.1007/s00442-016-3581-x)
- Jarosch KA (2016) New approaches to characterise soil organic phosphorus and to measure phosphorus transformation rates. Dissertation, ETH
- Jobbágy EG, Jackson RB (2004) The uplift of soil nutrients by plants. *Biogeochemical consequences across scales*. *Ecology* 85(9):2380–2389. doi:[10.1890/03-0245](https://doi.org/10.1890/03-0245)
- Joergensen RG (1996) The fumigation-extraction method to estimate soil microbial biomass. Calibration of the kEC value. *Soil Biol Biochem* 28(1):25–31. doi:[10.1016/0038-0717\(95\)00102-6](https://doi.org/10.1016/0038-0717(95)00102-6)
- Joergensen RG, Kübler H, Meyer B, Wolters V (1995) Microbial biomass phosphorus in soils of beech (*Fagus sylvatica* L.) forests. *Biol Fertil Soils* 19(2–3):215–219. doi:[10.1007/BF00336162](https://doi.org/10.1007/BF00336162)
- Joergensen RG, Mueller T (1996) The fumigation-extraction method to estimate soil microbial biomass. Calibration of the kEN value. *Soil Biol Biochem* 28(1):33–37. doi:[10.1016/0038-0717\(95\)00101-8](https://doi.org/10.1016/0038-0717(95)00101-8)
- John MK (1970) Colorimetric determination of phosphorus in soil and plant materials with ascorbic acid. *Soil Sci* 109(4):214–220
- Jonard M, Fürst A, Verstraeten A, Thimonier A, Timmermann V, Potočić N, Waldner P, Benham S, Hansen K, Merilä P, Ponette Q, de la Cruz AC, Roskams P, Nicolas M, Croisé L, Ingerslev M, Matteucci G, Decinti B, Bascietto M, Rautio P (2015) Tree mineral nutrition is deteriorating in Europe. *Glob Chang Biol* 21:418–430
- Keller M, Oberson A, Annaheim KE, Tamburini F, Mäder P, Mayer J, Frossard E, Bünemann EK (2012) Phosphorus forms and enzymatic hydrolyzability of organic phosphorus in soils after 30 years of organic and conventional farming. *Z Pflanzenernaehr. Bodenk* 175(3):385–393. doi:[10.1002/jpln.201100177](https://doi.org/10.1002/jpln.201100177)
- Kelly EF, Chadwick OA, Hilinski TE (1998) The effect of plants on mineral weathering. In: van Breemen N (ed) *Plant-induced soil changes: processes and feedbacks*. Springer, Dordrecht, pp 21–53
- Laliberté E, Grace JB, Huston MA, Lambers H, Teste FP, Turner BL, Wardle DA (2013) How does pedogenesis drive plant diversity? *Trends Ecol Evol* 28(6):331–340. doi:[10.1016/j.tree.2013.02.008](https://doi.org/10.1016/j.tree.2013.02.008)
- Lang F, Bauhus J, Frossard E, George E, Kaiser K, Kaupenjohann M, Krüger J, Matzner E, Polle A, Prietzel J, Renzenberg H, Wellbrock N (2016) Phosphorus in forest ecosystems: New insights from an ecosystem nutrition perspective. *J Plant Nutr Soil Sci* 179:129–135
- Lorenz M (1995) International co-operative programme on assessment and monitoring of air pollution effects on forests-ICP forests-. *Water Air Soil Pollut* 85(3):1221–1226. doi:[10.1007/BF00477148](https://doi.org/10.1007/BF00477148)
- Lundström U, van Breemen N, Bain D (2000) The podzolization process. A review. *Geoderma* 94(2–4):91–107. doi:[10.1016/S0016-7061\(99\)00036-1](https://doi.org/10.1016/S0016-7061(99)00036-1)
- Margesin R (1996) Phosphodiesterase activity. In: Schinner F, Öhlinger R, Kandeler E, Margesin R (eds) *Methods in soil biology*. Springer, Heidelberg, pp 217–219

- Marschner P, Hatam Z, Cavagnaro TR (2015) Soil respiration, microbial biomass and nutrient availability after the second amendment are influenced by legacy effects of prior residue addition. *Soil Biol Biochem* 88:169–177. doi:[10.1016/j.soilbio.2015.05.023](https://doi.org/10.1016/j.soilbio.2015.05.023)
- McGrath JW, Chin JP, Quinn JP (2013) Organophosphonates revealed: new insights into the microbial metabolism of ancient molecules. *Nature reviews. Microbiology* 11(6):412–419. doi:[10.1038/nrmicro3011](https://doi.org/10.1038/nrmicro3011)
- Mehra OP, Jackson ML (1960) Iron oxide removal from soils and clay by a dithionite-citrate system buffered with sodium bicarbonate. *Clays Clay Miner* 7:317–327
- Mellert KH, Göttlein A (2012) Comparison of new foliar nutrient thresholds derived from van den Burg's literature compilation with established central European references. *Eur J For Res* 131(5):1461–1472. doi:[10.1007/s10342-012-0615-8](https://doi.org/10.1007/s10342-012-0615-8)
- Mooshammer M, Wanek W, Zechmeister-Boltenstern S, Richter A (2014) Stoichiometric imbalances between terrestrial decomposer communities and their resources: mechanisms and implications of microbial adaptations to their resources. *Front Microbiol* 5:22. doi:[10.3389/fmicb.2014.00022](https://doi.org/10.3389/fmicb.2014.00022)
- Mouginot C, Kawamura R, Matulich KL, Berlemont R, Allison SD, Amend AS, Martiny AC (2014) Elemental stoichiometry of Fungi and Bacteria strains from grassland leaf litter. *Soil Biol Biochem* 76:278–285. doi:[10.1016/j.soilbio.2014.05.011](https://doi.org/10.1016/j.soilbio.2014.05.011)
- Murphy J, Riley JP (1962) A modified single solution method for the determination of phosphate in natural waters. *Analytica chimica acta* 27:31–36
- Netzer F, Schmid C, Herschbach C, Rennenberg H (2017) Phosphorus-nutrition of European beech (*Fagus sylvatica* L.) during annual growth depends on tree age and P-availability in the soil. *Environ Exp Botany* 137:194–207. doi:[10.1016/j.envexpbot.2017.02.009](https://doi.org/10.1016/j.envexpbot.2017.02.009)
- Odum EP (1969) The strategy of ecosystem development. *Science* 164(3877):262–270. doi:[10.1126/science.164.3877.262](https://doi.org/10.1126/science.164.3877.262)
- Oehl F, Oberson A, Sinaj S, Frossard E (2001) Organic phosphorus mineralization studies using isotopic dilution techniques. *Soil Sci Soc Am J* 65(3):780. doi:[10.2136/sssaj2001.653780x](https://doi.org/10.2136/sssaj2001.653780x)
- Öhlinger R (1996) Phosphomonoesterase activity with the substrate phenylphosphate. In: Schinner F, Öhlinger R, Kandeler E, Margesin R (eds) *Methods in Soil Biology*. Springer, Heidelberg, pp 210–213
- Ohno T, Zibilske LM (1991) Determination of low concentrations of phosphorus in soil extracts using malachite green. *Soil Sci Soc Am J* 55(3):892. doi:[10.2136/sssaj1991.03615995005500030046x](https://doi.org/10.2136/sssaj1991.03615995005500030046x)
- Paul EA (ed) (2015) *Soil microbiology, ecology and biochemistry*, 4th edn. Elsevier; Academic Press, Amsterdam, London
- Pearson HL, Vitousek PM (2002) Soil phosphorus fractions and symbiotic nitrogen fixation across a substrate-age gradient in Hawaii. *Ecosystems* 5(6):587–596
- Peters R (1997) *Beech forests*, vol 24. Geobotany. Springer, Netherlands
- Pitman R, Bastrup-Birk A, Breda N, Rautio P (2010) Sampling and Analysis of Litterfall. In: *Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests*, Part XIII, pp 1–16
- Porder S, Ramachandran S (2013) The phosphorus concentration of common rocks—a potential driver of ecosystem P status. *Plant Soil* 367(1–2):41–55. doi:[10.1007/s11104-012-1490-2](https://doi.org/10.1007/s11104-012-1490-2)
- Prietz J, Klysubun W, Werner F (2016) Speciation of phosphorus in temperate zone forest soils as assessed by combined wet-chemical fractionation and XANES spectroscopy. *J Plant Nutr Soil Sci* 179(2):168–185. doi:[10.1002/jpln.201500472](https://doi.org/10.1002/jpln.201500472)
- Randriamanantsoa L, Morel C, Rabeharisoa L, Douzet J, Jansa J, Frossard E (2013) Can the isotopic exchange kinetic method be used in soils with a very low water extractable phosphate content and a high sorbing capacity for phosphate ions? *Geoderma* 200–201:120–129. doi:[10.1016/j.geoderma.2013.01.019](https://doi.org/10.1016/j.geoderma.2013.01.019)
- Rautio P, Fürst A, Stefan K, Raitio H, Bartels U (2010) Sampling and Analysis of Needles and Leaves. In: *ICP Forest (ed) Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests*, Part XII, pp 1–19
- Saunders WMH, Williams EG (1955) Observation on the determination of total organic phosphorus in soils. *J Soil Sci* 6(2):254–267. doi:[10.1111/j.1365-2389.1955.tb00849.x](https://doi.org/10.1111/j.1365-2389.1955.tb00849.x)
- Schmidt M, Veldkamp E, Corre MD (2016) Tree-microbial biomass competition for nutrients in a temperate deciduous forest, central Germany. *Plant Soil* 408(1–2):227–242. doi:[10.1007/s11104-016-2923-0](https://doi.org/10.1007/s11104-016-2923-0)
- Schneider K, Turrión M-B, Grieson PF, Gallardo JF (2001) Phosphatase activity, microbial phosphorus, and fine root growth in forest soils in the Sierra de Gata, western central Spain. *Biol Fertil Soils* 34:151–155. doi:[10.1007/s003740100387](https://doi.org/10.1007/s003740100387)
- Schwertmann U (1964) Differenzierung der Eisenoxide des Bodens durch Extraktion mit Ammoniumoxalat-Lösung. *Z Pflanzenernaehr. Dueng Bodenk* 105(3):194–202. doi:[10.1002/jpln.3591050303](https://doi.org/10.1002/jpln.3591050303)
- Selmants PC, Hart SC (2010) Phosphorus and soil development. Does the Walker and Syers model apply to semiarid ecosystems? *Ecology* 91(2):474–484. doi:[10.1890/09-0243.1](https://doi.org/10.1890/09-0243.1)
- Sumann M, Amelung W, Haumaier L, Zech W (1998) Climatic effects on soil organic phosphorus in the North American great plains identified by phosphorus-31 nuclear magnetic resonance. *Soil Sci Soc Am J* 62(6):1580. doi:[10.2136/sssaj1998.03615995006200060015x](https://doi.org/10.2136/sssaj1998.03615995006200060015x)
- Talkner U, Jansen M, Beese FO (2009) Soil phosphorus status and turnover in central-European beech forest ecosystems with differing tree species diversity. *Eur J Soil Sci* 60(3):338–346. doi:[10.1111/j.1365-2389.2008.01117.x](https://doi.org/10.1111/j.1365-2389.2008.01117.x)
- Talkner U, Meiwes KJ, Potočić N, Seletković I, Cools N, de Vos B, Rautio P (2015) Phosphorus nutrition of beech (*Fagus sylvatica* L.) is decreasing in Europe. *Ann For Sci* 72(7):919–928. doi:[10.1007/s13595-015-0459-8](https://doi.org/10.1007/s13595-015-0459-8)
- Ternan NG, Mc Grath JW, Mc Mullan G, Quinn JP (1998) Organophosphonates: occurrence, synthesis and biodegradation by microorganisms. *World J Microbiol Biotechnol* 14(5):635–647. doi:[10.1023/A:1008848401799](https://doi.org/10.1023/A:1008848401799)

- Tiessen H, Moir JO (2008) Characterization of available P by sequential extraction. In: Carter MR, Gregorich EG (eds) Soil sampling and methods of analysis. CRC Press, Boca Raton, pp 293–306
- Tipping E, Somerville CJ, Luster J (2016) The C:N:P: S stoichiometry of soil organic matter. *Biogeochemistry* 130(1–2):117–131. doi:[10.1007/s10533-016-0247-z](https://doi.org/10.1007/s10533-016-0247-z)
- Tischer A, Potthast K, Hamer U (2014) Land-use and soil depth affect resource and microbial stoichiometry in a tropical mountain rainforest region of southern Ecuador. *Oecologia* 175(1):375–393. doi:[10.1007/s00442-014-2894-x](https://doi.org/10.1007/s00442-014-2894-x)
- Turner BL (2004) Optimizing phosphorus characterization in animal manures by solution phosphorus-31 nuclear magnetic resonance spectroscopy. *Journal of Environmental Quality* 33(2):757–766. doi:[10.2134/jeq2004.7570](https://doi.org/10.2134/jeq2004.7570)
- Turner BL, Baxter R, Mahieu N, Sjögersten S, Whitton BA (2004) Phosphorus compounds in subarctic Fennoscandian soils at the mountain birch (*Betula pubescens*)—tundra ecotone. *Soil Biol Biochem* 36(5):815–823. doi:[10.1016/j.soilbio.2004.01.011](https://doi.org/10.1016/j.soilbio.2004.01.011)
- Turner BL, Condron LM (2013) Pedogenesis, nutrient dynamics, and ecosystem development. The legacy of T.W. Walker and J.K. Syers. *Plant Soil* 367(1–2):1–10. doi:[10.1007/s11104-013-1750-9](https://doi.org/10.1007/s11104-013-1750-9)
- Turner BL, Condron LM, Richardson SJ, Peltzer DA, Allison VJ (2007) Soil organic phosphorus transformations during pedogenesis. *Ecosystems* 10(7):1166–1181. doi:[10.1007/s10021-007-9086-z](https://doi.org/10.1007/s10021-007-9086-z)
- Turner BL, Condron LM, Wells A, Andersen KM (2012) Soil nutrient dynamics during podzol development under lowland temperate rain forest in New Zealand. *CATENA* 97:50–62. doi:[10.1016/j.catena.2012.05.007](https://doi.org/10.1016/j.catena.2012.05.007)
- Turner BL, Engelbrecht BMJ (2011) Soil organic phosphorus in lowland tropical rain forests. *Biogeochemistry* 103(1–3):297–315. doi:[10.1007/s10533-010-9466-x](https://doi.org/10.1007/s10533-010-9466-x)
- Turner BL, Haygarth PM (2005) Phosphatase activity in temperate pasture soils: potential regulation of labile organic phosphorus turnover by phosphodiesterase activity. *Sci Total Environ* 344(1–3):27–36. doi:[10.1016/j.scitotenv.2005.02.003](https://doi.org/10.1016/j.scitotenv.2005.02.003)
- Turrión M, Glaser B, Solomon D, Ni A, Zech W (2000) Effects of deforestation on phosphorus pools in mountain soils of the Alay range, Khyrgyzia. *Biol Fertil Soils* 31(2):134–142. doi:[10.1007/s003740050636](https://doi.org/10.1007/s003740050636)
- Väänänen R, Hristov J, Tanskanen N, Hartikainen H, Nieminen M, Ilvesniemi H (2008) Phosphorus sorption properties in podzolic forest soils and soil solution phosphorus concentration in undisturbed and disturbed soil profiles. *Boreal Environ Res* 13(6):553–567
- Vadeboncoeur MA, Hamburg SP, Blum JD, Pennino MJ, Yanai RD, Johnson CE (2012) The quantitative soil pit method for measuring belowground carbon and nitrogen stocks. *Soil Sci Soc Am J* 76(6):2241–2255
- Vance ED, Brookes PC, Jenkinson DS (1987) An extraction method for measuring soil microbial biomass C. *Soil Biol Biochem* 19(6):703–707. doi:[10.1016/0038-0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6)
- Velásquez G, Ngo P, Rumpel C, Calabi-Floody M, Redel Y, Turner BL, Condron LM, de la Luz Mora M (2016) Chemical nature of residual phosphorus in Andisols. *Geoderma* 271:27–31. doi:[10.1016/j.geoderma.2016.01.027](https://doi.org/10.1016/j.geoderma.2016.01.027)
- Vincent AG, Vestergren J, Gröbner G, Persson P, Schleucher J, Giesler R (2013) Soil organic phosphorus transformations in a boreal forest chronosequence. *Plant Soil* 367(1–2):149–162. doi:[10.1007/s11104-013-1731-z](https://doi.org/10.1007/s11104-013-1731-z)
- Winkler JB, Dannenmann M, Simon J, Pena R, Offermann C, Sternad W, Clemenz C, Naumann PS, Gasche R, Kögel-Knabner I, Gessler A, Rennenberg H, Polle A (2010) Carbon and nitrogen balance in beech roots under competitive pressure of soil-borne microorganisms induced by girdling, drought and glucose application. *Funct Plant Biol* 37(9):879. doi:[10.1071/FP09309](https://doi.org/10.1071/FP09309)
- Wood T, Bormann FH, Voigt GK (1984) Phosphorus cycling in a Northern hardwood forest: biological and chemical controls. *Science* 223:391–393. doi:[10.1126/science.223.4634.391](https://doi.org/10.1126/science.223.4634.391)
- WRB (2015) World reference base for soil resources 2014, update 2015. International soil classification system for naming soils and creating legends for soil maps. World soil resources reports. FAO, Rome
- Wu YH, Prietzel J, Zhou J, Bing HJ, Luo J, Yu D, Sun SQ, Liang JH, Sun HY (2014) Soil phosphorus bioavailability assessed by XANES and Hedley sequential fractionation technique in a glacier foreland chronosequence in Gongga Mountain, Southwestern China *Sci China Earth Sci* 57(8):1860–1868. doi:[10.1007/s11430-013-4741-z](https://doi.org/10.1007/s11430-013-4741-z)
- Xu X, Thornton PE, Post WM (2013) A global analysis of soil microbial biomass carbon, nitrogen and phosphorus in terrestrial ecosystems. *Global Ecol Biogeogr* 22(6):737–749. doi:[10.1111/geb.12029](https://doi.org/10.1111/geb.12029)
- Yang N, Zavišić A, Pena R, Polle A (2016) Phenology, photosynthesis, and phosphorus in European beech (*Fagus sylvatica* L.) in two forest soils with contrasting P contents. *J Plant Nutr Soil Sci* 179(2):151–158. doi:[10.1002/jpln.201500539](https://doi.org/10.1002/jpln.201500539)
- Zavišić A, Nassal P, Yang N, Heuck C, Spohn M, Marhan S, Pena R, Kandeler E, Polle A (2016) Phosphorus availabilities in beech (*Fagus sylvatica* L.) forests impose habitat filtering on ectomycorrhizal communities and impact tree nutrition. *Soil Biol Biochem* 98:127–137. doi:[10.1016/j.soilbio.2016.04.006](https://doi.org/10.1016/j.soilbio.2016.04.006)