

# Traces of sunlight in the organic matter biogeochemistry of two shallow subarctic lakes

Marttiina V. Rantala · Carsten Meyer-Jacob · E. Henriikka Kivilä · Tomi P. Luoto · Antti. E. K. Ojala · John P. Smol · Liisa Nevalainen ·

Received: 24 November 2020/Accepted: 9 June 2021/Published online: 18 June 2021  $\ensuremath{\mathbb{C}}$  The Author(s) 2021

Abstract Global environmental change alters the production, terrestrial export, and photodegradation of organic carbon in northern lakes. Sedimentary biogeochemical records can provide a unique means to understand the nature of these changes over long time scales, where observational data fall short. We deployed in situ experiments on two shallow subarctic lakes with contrasting light regimes; a clear tundra lake and a dark woodland lake, to first investigate the photochemical transformation of carbon and nitrogen

**Supplementary Information** The online version contains supplementary material available at https://doi.org/10.1007/s10533-021-00820-9.

Responsible Editor: Klaus-Holger Knorr

M. V. Rantala (⊠) · E. H. Kivilä · T. P. Luoto · L. Nevalainen Faculty of Biological and Environmental Sciences, Ecosystems and Environment Research Programme, University of Helsinki, Lahti, Finland e-mail: marttiina.rantala@helsinki.fi

M. V. Rantala CNR Water Research Institute (IRSA), Unit Verbania, Verbania, Italy

C. Meyer-Jacob · J. P. Smol Paleoecological Environmental Assessment and Research Laboratory (PEARL), Department of Biology, Queen's University, Kingston, ON, Canada elemental (C/N ratio) and isotope ( $\delta^{13}$ C,  $\delta^{15}$ N) composition in lake water particulate organic matter (POM) for downcore inferences. We then explored elemental, isotopic, and spectral (inferred lake water total organic carbon [TOC] and sediment chlorophyll a [CHLa]) fingerprints in the lake sediments to trace changes in aquatic production, terrestrial inputs and photodegradation before and after profound human impacts on the global carbon cycle prompted by industrialization. POM pool in both lakes displayed tentative evidence of UV photoreactivity, reflected as increasing  $\delta^{13}C$  and decreasing C/N values. Through time, the tundra lake sediments traced subtle shifts in primary production, while the woodland lake carried signals of changing terrestrial contributions,

E. H. Kivilä Aquatic Laboratory, Département Des Sciences Fondamentales, Université du Ouébec À Chicou

Fondamentales, Université du Québec À Chicoutimi, Chicoutimi, QC, Canada

A. E. K. Ojala Geological Survey of Finland, Espoo, Finland

A. E. K. Ojala Department of Geography and Geology, University of Turku, Turku, Finland indicating shifts in terrestrial carbon export but possibly also photodegradation rates. Under global human impact, both lakes irrespective of their distinct carbon regimes displayed evidence of increased productivity but no conspicuous signs of increased terrestrial influence. Overall, sediment biogeochemistry can integrate a wealth of information on carbon regulation in northern lakes, while our results also point to the importance of considering the entire spectrum of photobiogeochemical fingerprints in sedimentary studies.

**Keywords** Global change · Paleo-optics · Particulate organic matter · Photobiodegradation · Sediment biogeochemistry · Subarctic

## Introduction

Sunlight governs fundamental biotic and abiotic processes in shallow northern lake ecosystems including those tightly connected to the aquatic carbon balance. Visible wavelengths of the solar spectrum fuel the conversion of inorganic carbon into organic forms through photosynthesis while light, particularly in the ultraviolet (UV) range, also disintegrates organic matter resulting either in partial photodegradation or complete photomineralization (Bertilsson and Tranvik 2000; Chen and Jaffé 2016; Cory and Kling 2018). Carbon sourced both from lake production and from the surrounding terrestrial environment contribute to the aquatic carbon cycle, while their reactivities and pathways within lakes often differ (Kellerman et al. 2015). Growing global human pressures, including warming but also changes in atmospheric deposition chemistry, are increasingly affecting the production, terrestrial export and processing of carbon in shallow northern lakes. These changes reshape underwater optical environments alongside the cycling of carbon (Fig. 1), carrying a threat of amplifying climate feedbacks parallel to cascading ecological regime shifts (Smol et al. 2005; Solomon et al. 2015).

Among the most prominent symptoms of Arctic warming are widespread decreases in the length of the snow and ice cover period (Magnuson et al. 2000; Callaghan et al. 2010; Griffiths et al. 2017) promoting the transfer of solar energy into aquatic ecosystems

(Fig. 1). The ensuing increase in photosynthetically active radiation (PAR) intertwined with nutrient and habitat stimuli are often found to increase the productivity of aquatic communities (Michelutti et al. 2005; Smol and Douglas 2007; Rantala et al. 2017) and may enhance photosynthetic carbon sequestration. There is, however, an ecological tradeoff to improved light availability, as concomitant increases in incident UV radiation may begin to impede algal growth through photoinhibition (Marwood et al. 2000; Vincent et al. 2007). Furthermore, increased UV irradiance can intensify photochemical processing of organic matter, inciting changes in the reactivity of lake organic carbon pools (Berggren et al. 2018) and increasing carbon dioxide (CO<sub>2</sub>) outgassing (Koehler et al. 2014). In addition to shaping internal carbon flows in lakes, global change is altering material transport from surrounding land areas (Fig. 1) with impacts on underwater light regimes and fundamental ecosystem functions (Solomon et al. 2015). Whether through changes in climate or alterations in atmospheric deposition chemistry, affecting the size and mobility of terrestrial carbon stocks, global change is increasing the flow of land-derived organic matter into northern waters (Finstad et al. 2016; Meyer-Jacob et al. 2019; Fig. 1). Colored terrestrial organic carbon is the predominant light attenuating medium in unproductive lakes (Laurion et al. 1997; Nevalainen et al. 2015) diminishing photosynthesis through shading (Karlsson et al. 2009; Ask et al. 2012) while reducing UV stress on aquatic biota (Williamson et al. 2016). Concurrent increases in terrestrial nutrient inputs may further reduce the adverse effects of increased shading and stimulate aquatic productivity (Rantala et al. 2016; Vasconcelos et al. 2018). Simultaneously, increased concentration or photoreactivity of land-derived organic matter may enhance rates of photochemical degradation (Lapierre et al. 2013; Cory and Kling 2018). These myriad environmental drivers often overlap, interact and incite threshold responses in northern lakes challenging predictions of the broad scale effects of global change on their carbon cycling. While the causes and consequences of changing aquatic production and terrestrial carbon export have been the focus of considerable research efforts over the past decades, uncertainties prevail due to the overlapping effects of climate change and other global human disturbances (Meyer-Jacob et al. 2019) and the heterogeneity of consequent lake responses, as



Fig. 1 Schematic depicting prevalent pathways of environmental impact on key lake carbon cycling processes in shallow northern lakes. Depicted are processes promoting or diminishing aquatic primary production, production and export of terrestrial organic matter and nutrients, and coupled

discussed below. As yet, few studies have sought to understand the repercussions of global change on photochemical degradation in shallow northern lakes despite its well-established importance in the aquatic carbon cycle (Cory et al. 2014; Sulzberger et al. 2019).

The impacts of warming are often exacerbated in northern lakes where snow and ice moderate biological activity through much of the year and even modest changes in the cryosphere can result in disproportionate ecological consequences. The effects are further aggravated by the continuous daylight of the summer months, high solar irradiance, and the often small and shallow character of the lake basins linking them tightly to the terrestrial carbon cycle (Rautio et al. 2011). The often nutrient-poor waters can also be considered highly sensitive to climate-mediated increases in nutrient input or atmospheric nitrogen deposition (Wolfe et al. 2006). In spite of these

photochemical and biological mineralization (described in more detail in the main text). Lake sediments integrate information of the environmental drivers as well as the pathways of lake carbon cycling they trigger. (Color figure online)

broadly unifying features, the sensitivity and responses of circumpolar lakes to climate change and atmospheric pollution begin to rapidly depart at regional and local spatial scales. Here the underwater optical environment may be considered as a gauge of environmental change (Williamson et al. 2014) but also as the backdrop for external perturbations. For instance, clear water lakes situated in barren tundra landscapes may be considered particularly vulnerable to changes in UV irradiance because of their low background organic carbon levels (Laurion et al. 1997; Rose et al. 2009) as opposed to wetland and thaw lakes brimming with light attenuating colored carbon (Wauthy et al. 2018). Other local scale features, such as hydrological connection to the surrounding catchment, will further modify the prevalent environmental drivers and the ecosystem response they trigger (Bogard et al. 2019). Moreover, the patterns and controls of lake carbon cycling are unlikely to advance linearly through time as the environmental backdrop is in constant change. The apparent heterogeneity of environmental drivers, rivaled only by the complexity of ensuing lake responses, underscores the need for a spatially and temporally comprehensive assessments of aquatic carbon regulation in northern lakes. Such assessments are severely hampered by the scarce, scattered and discontinuous monitoring data from remote northern environments that can, however, be supplemented with paleoenvironmental data obtained

from lake sediments (Smol and Douglas 2007; Smol

2019). The paleolimnological approach provides a means to overcome data limitations but also to decipher ecosystem-level responses to the multi-scaled and interactive environmental drivers while reducing noise inherent to fine-scale temporal variability (Williamson et al. 2014). Of particular use are easily measured biogeochemical indices that respond sensitively to environmental change, incorporate information of variability occurring within and beyond lake boundaries, and leave traceable fingerprints in lake water and sediment organic matter pools. Carbon and nitrogen elemental ratios (C/N ratio) and isotopic composition ( $\delta^{13}$ C,  $\delta^{15}$ N) in aquatic organic matter provide insights into vital ecosystem processes and are widely used to assess changes in carbon origins and pathways in lake ecosystems (Meyers and Teranes 2001; Talbot 2001). Where frequently used to trace past changes in aquatic primary production (Hodell and Schelske 1998; Wolfe et al. 2006) and terrestrial carbon inputs (Ninnes et al. 2017), they are also known to be influenced by photochemical processing, although the effects are not as well understood. Earlier experimental studies focused on the dissolved organic matter (DOM) pool indicate preferential losses of <sup>12</sup>C and carbon-rich compounds under light exposure, driving increases in  $\delta^{13}$ C (Ophsal and Zepp 2001; Vähätalo and Wetzel 2008) and decreases in C/N (Buffam and McGlathery 2003; Schmitt-Kopplin et al. 1998) of the residual organic matter. Photobiogeochemistry of particulate organic matter (POM) has been much less studied despite its role as a critical intermediate pool between lake water and sediments. Notwithstanding, POM is considered similarly sensitive to photodegradation (Vähätalo 2009) and there are indications of corresponding light-induced elemental and carbon isotope effects as those observed for DOM (Mayer et al. 2009). Overall, the elemental and isotopic composition of organic matter archived in lake sediments could thus integrate information of past variations in some of the most fundamental elements of aquatic carbon cycling in shallow northern lakes (Fig. 1).

Building on this notion, we investigated the biogeochemistry of lake water and sediment organic matter in two shallow lakes at the subarctic Fennoscandian treeline to elucidate the effects of global change on aquatic primary production, terrestrial carbon flows, and photodegradation in this sensitive lake environment. A clear mountainous tundra lake and a dark woodland lake were chosen for the analysis, both of which are typical of the region but represent contrasting carbon and underwater light regimes. Our first objective was to assess whether and how UV photoalteration might contribute to shaping the elemental (C/N) and isotopic ( $\delta^{13}$ C,  $\delta^{15}$ N) composition of lake water particulate organic matter (POM) in the lakes, for potential downcore applications. To this end, we employed in situ experiments on both lakes over four weeks to identify signals of photochemical degradation in the biogeochemistry of lake water POM. We expected to find increases in  $\delta^{13}$ C and declines in C/N of POM in treatments exposed to UV radiation, and further hypothesized that these signals could also be applied in the multiproxy sedimentary context to trace past trends in underwater light and photodegradation rates. Next, we examined elemental (C/N ratio) and isotopic ( $\delta^{13}$ C,  $\delta^{15}$ N) fingerprints in the lake sediments, searching for signals of both changing carbon source (pelagic and benthic aquatic production, terrestrial carbon export) and photochemical processing under natural climate fluctuations and ongoing global change. As the varied source and processing signals often overlap, reflectance spectroscopy was used to depict temporal trends in sediment chlorophyll a (CHLa) and lake water total organic carbon (TOC) concentrations as independent indices of changing aquatic primary production, water transparency, and terrestrial carbon input. We expected to find distinct but disparate signals of environmental change in connection to growing global human influence in the studied lakes. The tundra lake, with its extensive ice cover and low availability of organic substrate in the surrounding catchment, was suspected to be most vulnerable to direct climate impacts on lake ice regimes, and the increases in PAR and UV radiation associated with the lengthening of the ice free season. The dark woodland lake was expected to be more prone to catchment-mediated climate impacts increasing terrestrial carbon export into the lake (for instance, via catchment greening or wetland carbon mobilization). We also suspected that potential increases in the length of the ice free period driven by anthropogenic warming may have increased rates of photodegradation in the lakes. The results are expected to shed light on processes controlling carbon flows in shallow subarctic lakes over decadal to centennial time scales, and to advance the use of sedimentary paleo-optics in the study of lake carbon cycling in the face of growing human influence.

# Material and methods

### Study region

The study was conducted in northwestern Finnish Lapland (Fig. 2) on two lakes that share characteristics typical of lakes in the subarctic region and the circumpolar north more broadly (shallow waters, low productivity) but represent contrasting carbon and underwater light regimes. A tundra lake (Lake Iso-Jehkas, unofficial name) situated  $\sim 300$  m above the mountain birch treeline at an elevation of 925 m.a.s.l. was chosen to portray alpine clear water lakes with low concentrations of organic carbon in the water column and high underwater light exposure. Solar irradiance and the UV radiation dose on the lake are further increased by the high elevation (Rose et al. 2009). To represent a more organic-rich system with low light penetration and UV exposure, we selected a brown colored woodland lake (Lake Mukkavaara, unofficial name) situated at an elevation of 545 m.a.s.l. in the mountain birch woodlands. Rocky outcrops surrounding the tundra lake support mainly lichens, mosses and small shrubs, while the woodland lake is surrounded by patches of mountain birch, low growing dwarf birch and willow, shrubs, and waterlogged soils at the lake margin. Climate in the region is continental, with mean annual air temperature of -2.0 °C, mean July air temperature of 11.8 °C and mean annual precipitation of 530 mm. The growing season length is approximately three months. The lakes are small ( $\sim$  1 ha) with catchment areas of approximately  $\sim$  7 ha. Both lakes are hydrologically isolated with no permanent inlets or outlets. The tundra lake has ultraoligotrophic (total phosphorus 2.2  $\mu$ g L<sup>-1</sup>, mean of measurements performed in July and August 2018 from surface waters at  $\sim 0.5$  m depth) and oligohumic waters with DOC concentrations of 1.6 mg  $L^{-1}$ , while the woodland lake is characterized by oligotrophic (6.7  $\mu$ g L<sup>-1</sup>) and polyhumic waters with DOC of 8.0 mg  $L^{-1}$ . Both lakes have circumneutral waters, with pH values of 6.7 (mean of July and August measurements) in the tundra lake and 7.0 in the woodland lake. Colored dissolved organic matter (CDOM) absorption at 320 nm (a<sub>320</sub>), tightly coupled to underwater light attenuation (Nevalainen et al. 2015), is high in the woodland lake  $(\sim 21 \text{ m}^{-1})$  and low in the tundra lake  $(\sim 2 \text{ m}^{-1})$ . The difference in underwater light conditions is also visibly distinct, as the bottom of the slightly deeper tundra lake (maximum depth  $\sim 4$  m) was clearly visible, while that of the woodland lake ( $\sim 1.2$  m) was not. Iron concentrations were low (below the detection limit of 0.04 mg  $L^{-1}$ ) in both lakes and are not expected to carry large influence on underwater light attenuation. Detailed limnological characterizations of the lakes are provided in Nevalainen et al. (2020a).

# Experimental approach

An experimental analysis was first undertaken to assess whether and how the elemental and isotope biogeochemistry of lake water POM, part of which eventually settles to form lake sediment, displays sensitivity to photochemical alteration in the studied lakes. The experiments were performed in small mesocosms deployed directly onto the lakes to mimic the natural physicochemical and biological environment and ambient fluctuations in sunlight intensity and spectral composition. The experiments accordingly incorporate the effects of both photochemical and biological degradation that are intrinsically coupled in natural waters (Moran et al. 2000; Cory and Kling 2018). Nine treatments were performed on both lakes, comprising triplicate treatments under ambient sunlight (UV + PAR), sunlight without UV wavelengths (PAR), and no light (DARK) as a dark control. Vinyl acetate containers (Insinööritoimisto Haikonen Oy, Paippinen, Finland) with a volume, length and diameter of 340 L, 1.2 m and 0.6 m, respectively, were filled with lake surface water and set afloat on the lakes



**Fig. 2** a Location of the studied lakes in northwestern Finnish Lapland and characteristics of the landscape surrounding the **b** woodland lake ( $68.91^{\circ}$ N,  $21.00^{\circ}$ E) and **c** tundra lake ( $69.07^{\circ}$ N,  $20.88^{\circ}$ E)

from early July to early August in summer 2018. In the woodland lake, the enclosures effectively covered the entire water column and in the tundra lake approximately one-third of the depth of the water column. The containers were wrapped in a double layer of thick white plastic allowing light penetration only through the surface while maximizing the reflection of sunlight. The UV + PAR enclosures were covered with acetate foil (Aclar<sup>®</sup>, Honeywell, Charlotte, USA) transmitting 90-93% of UV (280-390 nm) and 93% of PAR. PAR enclosures were covered with UV blocking foil (no. 311413, Roscolab Ltd, London, UK) transmitting 82% of radiation > 390 nm while screening wavelengths < 390 nm. Dark controls were covered with two layers of thick black plastic. The foils were set up on convex aluminum netting allowing contact with air and preventing condensation on their surface. Wind mixing of the experimental waters was largely inhibited in the enclosures. The experiments were carried out during the Arctic summer exposing the enclosures to continuous daylight, although the highest irradiance period in the region occurs earlier, around mid-June. The incubation time was set at four weeks, estimated sufficient to produce detectable biogeochemical changes based on earlier studies examining isotopic transformation of the DOM pool (Ophsall and Zepp 2001; Chomicki 2009), while trying to minimize disparities between the experimental and natural system growing with time. Water residence time in the closed lake basins is much longer than the duration of the experiments, and the exposure of organic matter to light before burial is likely further promoted by the clear waters in the tundra lake (light reaching the surface sediments) and shallow depth of the woodland lake (increasing potential for sediment resuspension). In both lakes, efficient water column mixing is likely to increase the exposure of organic matter to photochemical processing. Direct scaling of the exposure time between the experimental waters and lake sediments is therefore limited, though neither critical as the primary focus here is on the direction of the elemental and isotopic transformations and on potential differences in their amplitude between the studied tundra and woodland lake.

Following the four week incubation period, the enclosures were sampled for lake water collected into opaque 5 L containers from the surface waters at  $\sim 0.5$  m depth. Concurrent with the installation of the experimental pools in July and collection of the samples in August, lake water was further sampled as reference of pretreatment conditions and natural biogeochemical succession in the lakes. Lake water particulate organic matter (POM) was filtered onto pre-combusted (450 °C for 2 h) GF/F filters. Terrestrial and aquatic vegetation were further sampled on both sites to estimate initial origins of the lake water organic matter pool. To assess changes in the size of the particulate organic carbon pool, we used total and dissolved organic carbon (TOC and DOC) to calculate concentrations of particulate organic carbon (POC) in the lake water and the experimental enclosures, and chlorophyll a concentrations to evaluate changes in the photosynthetic production of POM in the light treatments. Analytical details for the limnological and optical measurements are presented in Nevalainen et al. (2020a) investigating UV effects on the limnological, optical and pigment properties of the dissolved organic matter (DOM) pool in the same experiments.

All samples were kept under cold and dark conditions until further treatment and analysis. The lakes are situated on non-carbonate bedrock and, overall, any influence of carbonate precipitation by photosynthesis in these soft, circumneutral and oligotrophic waters can be considered minor. No measures were therefore taken to remove inorganic carbon since acid treatments can cause nonlinear deviations in  $\delta^{13}$ C and C/N values (Brodie et al. 2011; Schlacher and Connolly 2014). While we thus cannot rule out entirely the influence of carbonates on the elemental and carbon isotope composition of the experimental samples, this should mainly be considered when comparing the absolute values among lakes, whereas our analysis was focused on variability between the treatments. Elemental and isotope analyses were performed on a Delta V Plus continuous flow isotope ratio mass spectrometer linked to a Flash 2000 elemental analyzer with a MAS 200 R autosampler (Thermo-Fisher Scientific, Bremen, Germany) at the NIWA Environmental and Ecological Stable Isotope Facility (Wellington, New Zealand). The isotope ratios are expressed in delta ( $\delta$ ) notation as per mill (%) difference relative to international standard. The reference standards used were Vienna Pee Dee Belemnite for carbon and atmospheric nitrogen. The effects of sunlight on the biogeochemical composition of POM were calculated as respective differences in  $\delta^{13}$ C,  $\delta^{15}$ N and C/N between the different treatments, with particular focus on variability between the two light treatments (PAR and UV + PAR). Changes in POC concentrations were assessed in a similar manner. Notably, both visible and UV radiation contribute to photochemical reactions in the water column, while the UV wavelengths are generally considered as primary driver of organic matter photodegradation. The two were isolated here to better extract the effects of photodegradation from those related to photosynthetic production. Significance of the changes between treatments were tested with one-way analysis of variance (ANOVA) followed by Tukey's pairwise comparison where needed.

# Sedimentary approach

Paleolimnological analyses were subsequently undertaken on the lake sediments to search for biogeochemical evidence of centennial scale variations in aquatic primary production, terrestrial carbon export and photochemical processes. Short sediment cores (15 cm in length) were retrieved with a Limnos gravity corer in July 2018 following the deployment of the experimental enclosures on the lakes. The cores were retrieved from the deepest parts of the morphometrically simple small lake basins. Subsampling was performed in the field at 1 cm intervals and the samples were stored under cold and dark conditions prior to freeze drying and homogenization for elemental and isotope analysis. Radiometric techniques were used to obtain age estimates for the sediment cores, including radiocarbon analysis of sedimentary macrofossils in the deeper sediments and <sup>137</sup>Cs analysis of the surface sequences (topmost 0-10 cm). For <sup>137</sup>Cs, 1-cm thick samples (6-12 g)

were analyzed by direct gamma assay at the Geological Survey of Finland (Espoo, Finland) using a fully digital BrightSpec bMCAUSB pulse height analyzer coupled to a well-type NaI(Tl) detector. The activity concentrations were normalized with sample weights. Radiocarbon analysis was carried out at Beta Analytic (Miami, Florida, USA) on two terrestrial plant macrofossils found at sediment depths of 5 and 12 cm in the woodland lake, and on two moss fragments at depths of 7 and 14 cm in the tundra lake where other suitable plant macro remains were not found. In soft and well-mixed waters, mosses can, however, be considered appropriate for radiocarbon dating (Wolfe et al. 2004; Oswald et al. 2005). Elemental and isotope analyses were performed on freeze dried and homogenized sediment samples at the NIWA Environmental and Ecological Stable Isotope Facility (Wellington, New Zealand) similar to the experiment samples. Visible near-infrared (VNIR) spectroscopic analyses for the inference of past trends in lake water total organic carbon (TOC) and chlorophyll a (CHLa) concentrations were performed on freeze dried and sieved (125 µm) sediment samples. VNIR spectra were analyzed with a Model 6500 series Rapid Content Analyzer (FOSS NIRSystems Inc.) at the Paleoecological Environmental Assessment and Research Laboratory (Kingston, Canada). Scanning was performed through the base of glass vials in diffuse reflectance mode. Measurements were carried out over the wavelength range 400-2500 nm with each obtained spectra representing the mean of 32 scans at 2 nm resolution. Quantification of sediment chlorophyll a concentrations was done following Michelutti et al. (2010) using the peak area between 650 nm and 700 nm. The measurements capture both primary chlorophyll a and its main degradation products making the inferences less sensitive to diagenetic effects (Michelutti and Smol 2016; Rydberg et al. 2020). Lake water TOC estimates were based on a partial least squares regression (PLSR) model between VNIR spectra of surface sediments and corresponding modern surface water TOC concentrations following Meyer-Jacob et al. (2017). A subset of 138 lakes from the universal lake water TOC model for northern lakes (Meyer-Jacob et al. 2017) was used for past lake water TOC inferences in this study. This subset consists of lakes from northern Fennoscandia that are located within < 300 km of the studied lakes and cover a TOC gradient of 0.5 to 14.9 mg  $L^{-1}$ . The resulting 3-component PLSR model had a crossvalidated coefficient of determination of 0.62 and root mean square error of 1.6 mg L<sup>-1</sup>. Pairwise correlations were used to examine interconnection between the sediment biogeochemical indices (C/N ratio,  $\delta^{13}$ C,  $\delta^{15}$ N, TOC, CHLa), applying false discovery rates with Bonferroni correction to account for multiple comparison (q = 0.05).

# Results

Transformation of lake water POM biogeochemistry

The elemental and isotopic composition of POM measured from lake water in July and August in both lakes suggests mixed origins with contributions from pelagic and benthic aquatic production and terrestrial sources (Fig. 3). Mean  $\delta^{13}C_{POM}$  and  $\delta^{15}N_{POM}$  values slightly lower in the tundra were lake  $(-27.5\% \pm 0.7 \text{ SD} / - 0.3\% \pm 0.0 \text{ SD})$  compared to the woodland lake (-  $27.0\% \pm 1.2$  SD), /  $1.6\% \pm 1.7$  SD) where, in turn, C/N values were slightly lower (8.4  $\pm$  1.2 SD) relative to the tundra lake (8.9  $\pm$  1.5 SD). Between the three treatments, the elemental and isotopic composition of POM was distinctly different in both lakes following the fourweek incubation under no light (DARK), light without the UV wavelengths (PAR), and full sunlight (UV + PAR). No statistically significant differences were found based on ANOVA attributable to the relatively large variability among treatment replicates, however, some consistent and conspicuous patterns were observed. In the woodland lake, lowest  $\delta^{13}C_{POM}$ values were found in the DARK enclosures (mean - $29.7\% \pm 1.2$  SD) increasing towards the PAR treatment (- 29.4‰  $\pm$  0.6 SD) and finally UV + PAR  $(-28.4\% \pm 0.3 \text{ SD})$  (Fig. 4). In the tundra lake,  $\delta^{13}$ C patterns were more erratic, however; an increase in  $\delta^{13}C_{POM}$ was observed between PAR  $(-28.0\% \pm 0.9)$ and UV + PARSD)  $(-27.4\% \pm 0.0 \text{ SD})$  consistent with the trend in the woodland lake.  $\delta^{15}N_{POM}$  showed lowest and highest values in the DARK treatments in the woodland  $(-0.0\% \pm 0.5 \text{ SD})$  and tundra  $(2.0\% \pm 1.0 \text{ SD})$ lake, respectively, but no conspicuous and consistent patterns were found between the two light treatments. Both in the woodland and tundra lake, C/N<sub>POM</sub>

-18

-18



**Fig. 3** Carbon ( $\delta^{13}$ C) and nitrogen ( $\delta^{15}$ N) isotopic composition and elemental ratio (C/N) in terrestrial and aquatic vegetation, benthic autotrophic communities, and lake water particulate organic matter (POM; blank square) in the **a** tundra and **b** woodland lake. Corresponding values for POM in the experimental enclosures, following a four-week incubation under no light (DARK; black square), sunlight without the UV wavelengths (PAR; dark grey), and full sunlight (PAR + UV;

declined from DARK (8.4  $\pm$  0.6 and 7.3  $\pm$  0.6, respectively) to PAR (8.1  $\pm$  0.9 and 6.7  $\pm$  0.3, respectively) and UV + PAR (7.6  $\pm$  0.3 and

light grey) exposure are included for reference. Values for surface sediment samples are also indicated (blank diamond). Phytoplankton were not separately measured for the studied lakes but should place towards the upper left corner in the isotope diagrams, with typical values of  $-32.1\% \pm 1.1$  SD and  $3.8\% \pm 1.3$  SD for  $\delta^{13}$ C and  $\delta^{15}$ N, respectively (Kivilä et al. 2020)

 $6.3 \pm 0.2$ , respectively). UV exposure had a more conspicuous effect on POM composition than concentrations both in the tundra and woodland lake, as



**Fig. 4** Carbon ( $\delta^{13}$ C) and nitrogen ( $\delta^{15}$ N) isotopic composition and elemental ratio (C/N) in particulate organic matter (POM) sampled from in situ enclosures comprising treatments under no light (DARK; black squares), sunlight without the UV wavelengths (PAR; dark grey), and full sunlight (PAR + UV;

suggested by the minor variability in POC concentrations (Supplementary Fig. 1) between PAR (0.22 mg  $L^{-1} \pm 0.0$  SD and 0.46 mg  $L^{-1} \pm 0.3$  SD, respectively) and UV + PAR (0.22 mg  $L^{-1} \pm 0.0$  SD and 0.39 mg  $L^{-1} \pm 0.1$  SD, respectively) treatments. Changes in DOC concentrations were similarly very small (Supplementary Fig. 1; Nevalainen et al. 2020a) suggesting little transfer between the particulate and dissolved organic carbon pools during the experiments, although we did not quantify the production of new POC or loss of DOC via mineralization that could mask some of the variability. We focused primarily on changes between the three treatments rather than on changes from the initial values (i.e., before the incubation) as those could be influenced by the experimental setting. Briefly, without convective mixing, part of the more resistant POM fractions may have settled at the bottom of the enclosures over the four weeks. Second, the mixed lake water is likely to incorporate a stronger benthic signal relative to the stagnant experimental waters. Thus, signals of photochemical processing could partially mix with signals of changing organic matter source. Indeed, there was a shift towards more negative  $\delta^{13}C_{POM}$  in the enclosures in both lakes consistent with a relative increase in pelagic contribution (Figs. 3 and 4), even though chlorophyll a concentrations indicated lowered productivity (Nevalainen et al. 2020b, Supplementary Fig. 1). In keeping with the carbon isotope values,  $\delta^{15}N_{POM}$  in the tundra lake enclosures shifted towards

light grey) exposure. Horizontal lines display initial values measured from natural lake water upon deployment of the experiments in early July 2018. Box plots depict upper and lower quartiles (box), median (solid line), mean (dashed line), and minimum and maximum (whiskers) values

a typical pelagic signal (Kivilä et al. 2020) relative to initial values before incubation.

Centennial patterns in sediment biogeochemistry

Radiometric measurement yielded three age estimates for both lakes, with one from each deemed unreliable. Data for the four dates used in the study are presented in Table 1 and displayed in Figs. 5 and 6. Both lakes displayed relatively low <sup>137</sup>Cs activities (Supplementary Fig. 2) with a single peak attributed to the nuclear weapons testing peaking in the 1960s. In this environment, the Chernobyl fallout (1986) is less likely to leave a distinct peak as the lakes are located ~ 700–800 km north of the highest fallout area in central southern Finland (Arvela et al. 1990). The effect of post-fallout re-deposition from the catchment is probably minor in the closed lake basins, however, diffusion (Ojala et al. 2017) may have affected the

 Table 1
 Radiometric age estimates from the sediments of the studied tundra and woodland lake

Lake	Depth (cm)	Radionuclide	Estimated age range
Tundra	7	<sup>14</sup> C	530-452 cal BP
Tundra	14	<sup>14</sup> C	982–904 cal BP
Woodland	5	<sup>137</sup> Cs	~ 1960s
Woodland	12	<sup>14</sup> C	1290-1220 cal BP



Fig. 5 Centennial patterns in sediment biogeochemistry in the tundra lake, including sediment carbon ( $\delta^{13}$ C) and nitrogen ( $\delta^{15}$ N) isotopic composition, their elemental ratio (C/N), inferred lake water total organic carbon (TOC), and sediment

<sup>137</sup>Cs profile of the tundra lake. Specifically, <sup>137</sup>Cs concentrations in the tundra lake peaked at 7 cm inconsistent with the radiocarbon date at the same depth and at 14 cm (Table 1), as well as with age estimates from a long sediment core obtained from the same lake (Nevalainen et al. 2020b). For reference, most of the Holocene history ( $\sim 10500$  years) of the lake was encompassed in a 73-cm sediment sequence, as could be expected for such a sedimentation environment. Mixing of sediment can occur in shallow lake basins, however, this was not supported by the gradual development patterns in the biogeochemical proxies (Fig. 5) and the radiocarbon dating of the long sediment core indicating a stable sedimentation environment for the Holocene (Nevalainen et al. 2020b). The elevated organic content in the uppermost sediments combined with slow accumulation rates in turn may have promoted downward diffusion that could explain the deep (and broad) <sup>137</sup>Cs peak in the lake. In the woodland lake, radiocarbon analysis gave an age

chlorophyll *a* (CHLa; indicative of whole lake primary production). Available age estimates are displayed with horizontal lines and mean values for each sediment variable with vertical dotted lines

estimate of 5130 yr BP for the sample at depth 6 cm possibly related to redeposition of older material (Wolfe et al. 2004). The age is broadly inconsistent with the other age estimates from the profile (Table 1) and dating results from a long sediment core from the same lake (unpublished results). The <sup>137</sup>Cs date at 7 cm for the tundra lake and the <sup>14</sup>C date for the woodland lake were hence rejected. As our work focuses on dynamic relationships between the sedimentary indices, we did not attempt age-depth modelling for either of the cores but use the obtained point in time estimates to provide a coarse chronological framework for the sedimentary inferences.

Through time, variations in sediment biogeochemical proxies were relatively small but largely synchronous throughout the two sediment records (see Supplementary Fig. 3 for pairwise comparisons). Chlorophyll *a* and inferred lake water TOC were generally lower and showed smaller variability in the tundra lake (mean 0.04 mg g<sup>-1</sup>  $\pm$  0.01 SD and



Fig. 6 Centennial patterns in sediment biogeochemistry in the woodland lake, including sediment carbon ( $\delta^{13}$ C) and nitrogen ( $\delta^{15}$ N) isotopic composition, their elemental ratio (C/N), inferred lake water total organic carbon (TOC), and sediment

2.5 mg  $L^{-1}\pm0.3$  SD, respectively) relative to the woodland lake (0.09 mg  $g^{-1} \pm 0.05$  SD and 7.8 mg  $L^{-1} \pm 0.9$  SD, respectively).  $\delta^{13}C$  were slightly more negative in the tundra lake  $(-24.7\% \pm 0.3 \text{ SD})$ compared to the woodland lake ( $-24.0\% \pm 0.5$  SD), while C/N values were very similar between the lakes  $(11.7 \pm 0.8 \text{ and } 11.8 \pm 0.9 \text{ SD}, \text{ respectively})$ . In the tundra lake (Fig. 5), changes in sediment chlorophyll a were positively correlated to variations in inferred lake water TOC (R = 0.73, p = 0.003) and negatively correlated to sediment C/N (R = -0.77) $p = \langle 0.001 \rangle$  and, to lesser extent, to changes in  $\delta^{15}$ N (R = -0.67, p = 0.007). Variations in  $\delta^{13}$ C were more erratic and displayed no conspicuous connections to the other variables. Both chlorophyll a and lake water TOC showed a very subtle increasing trend (by + 0.007 mg g<sup>-1</sup> and + 0.3 mg  $L^{-1}$ , respectively) across the lower half of the sediment sequence matched by declining (-0.5) C/N values.

chlorophyll *a* (CHLa; indicative of whole lake primary production). Available age estimates are displayed with horizontal lines and mean values for each sediment variable with vertical dotted lines

At around 6 cm, chlorophyll a and TOC showed a synchronous decline ( $-0.009 \text{ mg g}^{-1}$  and -0.8 mg $L^{-1}$ , respectively) followed by a more pronounced increase (+ 0.03 mg g<sup>-1</sup> and + 1.2 mg L<sup>-1</sup>, respectively) near the surface. These trends were mirrored by C/N, though with a barely discernible increase (up to + 0.3) after 6 cm and a more distinct surface decline (-1.5). In the woodland lake (Fig. 6), inferred lake water TOC, sediment C/N ratio and  $\delta^{13}$ C showed similar patterns, though only the latter two variables were significantly negatively correlated (R = -0.76, p = 0.001) after correction for multiple comparisons. A transient increase in lake water TOC and C/N  $(+2.1 \text{ mg L}^{-1} \text{ and } +2.3, \text{ respectively})$  coupled to a decline (- 0.8‰) in  $\delta^{13}$ C occurred between around 6 and 7 cm. Below, TOC (mean 8.1 mg  $L^{-1}$ ) and C/N (12.0) were generally more elevated and  $\delta^{13}C$ (-24.0%), more negative compared to the section (6.9 mg  $L^{-1}$ . 11.0‰ and -23.6%, above

respectively), although inferred lake water TOC displayed an increasing trend in the topmost 3 samples  $(+1.1 \text{ mg L}^{-1})$ . Chlorophyll *a* showed little change in the lower half of the sediment sequence followed by an increasing  $(+0.1 \text{ mg g}^{-1})$  trend towards the surface initiated at around 5 cm.

# Discussion

# UV photoalteration of lake water POM biogeochemistry

Our understanding of the control of carbon transformation and mineralization in lake ecosystems has shifted over recent years upon recognition of the key importance of photodegradation in shaping the pathways and fate of organic matter in northern waters (Cory et al. 2014; Sulzberger et al. 2019). However, comparatively little research has been carried out on the effects of photoalteration on lake water POM despite its inherent connection to the sediment carbon pool. Results from our four week experiments suggest that the elemental and carbon isotopic composition of lake water POM in both of the studied subarctic lakes may be sensitive to UV-driven photochemical degradation. Although generally consistent with our expectations, the shifts were comparatively small and should therefore be considered tentative. The experiments were conducted on natural waters for a more accurate depiction of the natural environment, yet with less control over the multiple interacting ecosystem processes that likely account for the scatter between treatment replicates. Whatever combination of processes underlie the observed changes, the direction of the shifts in  $\delta^{13}$ C and C/N of lake water POM in both lakes is consistent with the earlier experimental works. Accordingly, the observed enrichment of the organic matter  $\delta^{13}$ C signature under UV exposure in both lakes (Fig. 4) concurs with the previously observed preferential degradation of the <sup>12</sup>C fraction of carbon under sunlight (Ophsal and Zepp 2001; Vähätalo and Wetzel 2008). Similarly, the decline in C/N conforms with the anticipated faster loss of carbon relative to nitrogen under light exposure (Schmitt-Kopplin et al. 1998; Buffam and McGlathery 2003; Chomicki 2009). These changes have been attributed to the loss of high molecular weight aromatic compounds such as terrestrial lignins (Ophsal and Benner 1998; Vähätalo et al. 1999; Mayer et al. 2009) depleted in <sup>13</sup>C and enriched in carbon (Benner et al. 1987). The process appears plausible also here considering the apparent terrestrial component in the POM pool of both lakes (Fig. 3) corroborated by the spectral characteristics of the lake water DOM (Nevalainen et al. 2020a).

Importantly, the observed changes in POM biogeochemistry between the two light treatments do not depict an isolated signature of photochemical processing but also of tightly coupled biological processes. Although the extent and direction of their influence on elemental and isotopic ratios may be more erratic, there is an element of predictability linked to the origin and biogeochemical characteristics of organic matter. Where primarily autochthonous organic substrate can transform into more biorefractory compounds under light exposure (Benner and Biddanda 1998; Tranvik and Bertilsson 2001), photodegradation generally renders terrestrial organic matter more labile (Moran et al. 2000; Cory and Kling 2018) thereby promoting biodegradation. Consistently, previous research suggests that photoexposure is likely to increase overall microbial activity in clear water and humic oligotrophic lakes due to the low availability of labile carbon produced within-lake (Tranvik and Bertilsson 2001). Bacteria preferentially remove autochthonous organic carbon over lignin and other aromatic terrestrial compounds (Benner et al. 1987) entailing that the effect of photodegradation on organic matter biogeochemistry could in effect be partially muted by the biotic interaction. DOM fluorescence components indicated a considerably stronger microbial signal in the woodland lake  $(\sim 55\%)$  relative to the tundra lake  $(\sim 15\%)$ ; Nevalainen et al. 2020b), in keeping with earlier research suggesting that biodegradation dominates organic matter processing in dark colored lakes (Berggren et al. 2018). In accordance, the low  $\delta^{13}$ C and elevated C/N dark control values (Fig. 4) imply that coupled bacterial processing could be dampening the photochemical signal in the woodland lake. The elevated dark control  $\delta^{13}$ C and C/N in the tundra lake in turn resemble those of the initial lake water POM, suggesting weaker photobiological interaction. Biodegradation aside, we observed lowered chlorophyll a concentrations in the enclosures exposed to full sunlight (Supplementary Fig. 1) potentially related to UV photoinhibition of algal production (Nevalainen et al. 2020a). Reduced demand for inorganic carbon results in higher photosynthetic discrimination against  $^{13}$ C (Hodell and Schelske 1998) and could thus contribute to diminishing the UV effect on carbon isotope signatures. Overall, the consistent shifts towards more enriched  $\delta^{13}$ C values and lowered C/N under UV exposure suggest that the coupled biotic interactions may weaken but not entirely diminish the signal of photochemical processing in the studied lakes, conforming to earlier research considering photochemical and coupled microbial effects on DOM isotope biogeochemistry (Ophsal and Zepp 2001).

Irrespective of their distinct carbon and underwater light regimes, the amplitude of the changes in  $\delta^{13}$ C and C/N between the two light treatments (PAR and UV + PAR) (Fig. 4) did not differ markedly between the studied lakes. In the tundra lake, mean  $\delta^{13}C$ increase under UV exposure was lower (+0.6%)compared to the woodland lake (+1%); however, the decline in C/N was slightly larger (-0.6 vs. - 0.3). The lack of clear distinction between the lakes may relate to confounding biological processes, such as microbial degradation in the woodland lake negating the effects of photodegradation, but also to tradeoffs between substrate lability and availability, and light exposure (Cory et al. 2015). In particular, the effects of photodegradation could be delimited in the woodland lake by the rapid attenuation of light in the dark water column (Williamson et al. 1996; Nevalainen et al. 2015) confining photodegradation to the surface layer, and in the tundra lake by the low availability of photolabile organic matter (Nevalainen et al. 2020a). It should also be noted that, in both lakes, the organic matter has been exposed to natural photochemical and biological processing already prior to the incubation, which is likely to partially account for the small amplitude of the observed biogeochemical changes. Notably, while the results are not directly comparable due to differences in experimental design and environment, our results correspond well, both in direction and extent, with the results of Mayer et al. (2009) examining photolytic transformation of riverine and coastal suspended sediments. The changes in  $\delta^{13}C$ observed by the authors ranged from + 0.4‰ to +0.5% and the declines in C/N from -0.7 to -1.3, following 70 h of constant irradiation by strong light (corresponding to mid-day subtropical sun). This was the only earlier study we found investigating the photolytic transformation of the elemental and isotopic composition of POM precluding further comparisons. For reference, the amplitude of the  $\delta^{13}$ C shifts observed here is at the lower range of variation found in earlier experimental works studying DOM transformation under light exposure (Osburn et al. 2001; Ophsal and Zepp 2001; Vähätalo and Wetzel 2008; Chomicki 2009). Further comparative experiments are evidently needed to decipher the extent to which photochemical and coupled biological processing may contribute to the elemental and isotopic transformation of lake water POM in different lake environments. Although the results from our pioneering study are tentative and do not directly scale with the sediment organic matter that may have been exposed to sunlight for much longer periods of time in the closed lake basins and well-mixed waters, they nevertheless suggest that photodegradation could be a relevant driver of POM biogeochemistry that should not be overlooked in sedimentary studies.

Tracing solar biogeochemical fingerprints through time

In closed lake systems, the fraction of POM not reduced to the dissolved phase and mineralized settles at the bottom of the lake to form a record of its past origins and of subsequent biotic and abiotic alterations. The relative importance of each type of source and alteration varies broadly between lake ecosystems and may vary through time under environmental transitions, whether of natural or human origin. In the studied tundra lake, the most prominent long-term patterns in organic matter biogeochemistry were associated with changes in aquatic primary production, as indicated by the tightly correlated trends in sediment chlorophyll a, C/N ratio and inferred lake water TOC (Fig. 5). Prevalence of the subtle productivity signal in the ultraoligotrophic waters indicates that, in the context of the late Neoglacial climate regime, the lake has remained largely isolated from shifts in terrestrial carbon cycling. Besides recent warming, the sediment sequence covers the Medieval Warm Period ( $\sim$  900–1300 CE) and the following cold period referred to as the Little Ice Age (prevailing until the twentieth century in northern Finland; Luoto et al. 2017) that have left distinct imprints in paleoenvironmental records across the region (Weckström et al. 2006; Kivilä et al. 2020). Although we cannot pinpoint the exact timing of the changes due to chronological constraints, our record suggests that these earlier climate phases have not driven marked changes in the productivity of the lake. The slight chlorophyll a increase towards the sediment surface may suggest a recent increase in aquatic production potentially linked to the observed lengthening of the ice free season over the past century in subarctic Fennoscandia (Callaghan et al. 2010), possibly coupled with atmospheric nitrogen deposition given the declining  $\delta^{15}$ N at the surface (Wolfe et al. 2004). Trends near the sediment surface must, however, always be considered with some caution due to potential diagenetic overprinting. Even though spectral chlorophyll a inferences also incorporate degradation products of chlorophyll a and are therefore less sensitive to post-depositional alteration (Michelutti et al. 2010; Rydberg et al. 2020), some chlorophyll a loss can still occur within the first years after deposition. For example, Rydberg et al. (2020) showed up to 24-45% losses before chlorophyll a levels stabilized 10-15 years after initial sediment deposition in a boreal lake in Sweden. The nearsurface declines in sediment C/N (  $\sim~2$  units) and  $\delta^{15}N$  $(\sim 1.5\%)$  (Fig. 5) could also be partially related to the progressive loss of nitrogen and <sup>14</sup> N with time (Talbot et al. 2001; Gälman et al. 2008). We therefore interpret these recent changes with some caution.

Increases in PAR denote also increasing UV stress (Fig. 1) that can carry cascading effects extending from the cellular level to entire ecosystem functioning (Vincent et al. 2007). UV influence may be considered particularly pronounced in clear and shallow alpine lakes where the underwater light spectrum inclines towards the UV range (Laurion et al. 1997), the thinner atmosphere increases solar exposure (Rose et al. 2009), low temperatures and nutrient deficiency impair photoprotective pigmentation and cellular repair (Beardall et al. 2014), and the shallow waters and continuous summer daylight further limit behavioral adaptations to UV stress. Although primary production in the tundra lake may thus be considered highly sensitive to photoinhibition, the benefits of improved resource availability still appear to have exceeded the adverse effects of UV, even if the potential increase in production was very small. Neither did we detect concomitant opposing patterns in organic matter  $\delta^{13}$ C and C/N values that could have carried indications of enhanced photoprocessing over the recent period, nor earlier in the sediment sequence. It could be that the recorded environmental variability is too subtle to drive changes in photoprocessing rates, however, photodegradation could also be insensitive to increased light in the system due to substrate limitation. In clear waters, already short exposure to UV can rapidly deplete photolabile organic matter fractions (Cory et al. 2015) and thus long-term variability in the rates of photodegradation could be primarily sensitive to changes in the concentrations (or photoreactivity) of terrestrial organic carbon.

In the woodland lake, centennial patterns in sediment biogeochemistry appeared to be linked to varying contributions of terrestrial carbon to the lake water, as indicated by the coeval shifts in sediment  $\delta^{13}$ C values, C/N ratio, and inferred lake water TOC (Fig. 6). Interestingly, while the transient upward shift in C/N and decline in  $\delta^{13}$ C towards typical terrestrial values ( $\sim 28\%$ , Fig. 3, Kivilä et al. 2020) with increasing lake water TOC plausibly reflect increased catchment-derived carbon inputs, the experimental results, corroborated by the earlier research (Ophsal and Zepp 2001; Buffam & McGlathery 2003; Mayer et al. 2009), suggest that lowered photochemical processing could contribute towards a similar elemental and isotopic shift. Correspondingly, <sup>13</sup>C enrichment together with lowered C/N and TOC earlier in the sequence and particularly over the past decades could either point to reduced terrestrial influence or enhanced photodegradation rates. In light of the landscape context and limnological characteristics of the lake, both processes are equally conceivable. For one, carbon flows from the surrounding marshy soils may be expected to be sensitive to changing temperatures and hydrology (Tranvik and Jansson 2002), and we also cannot rule out potential effects of acidification and subsequent recovery on carbon fluxes (Meyer-Jacob et al. 2019) over the past few decades. Unlike in the tundra site, photodegradation in the dark woodland lake is likely to be light limited (Cory et al. 2015) meaning that the rates of photoalteration are more likely to accelerate under increased light than changing concentrations of terrestrial organic carbon. Increased UV exposure under lengthened growing season could thereby either drive or contribute to the generally lowered TOC and prevalent autochthonous signal over the past decades. Unfortunately, we cannot confidently link the earlier transitions in sediment biogeochemistry with the preceding climate phases due to chronological constraints. Additional

sedimentary proxies are also needed to extract the individual or synergistic contributions of changing source (terrestrial influence) and processing (photodegradation rates) on the temporal patterns. Notwithstanding, the results point to a potentially overlooked role of photodegradation in controlling sediment biogeochemical patterns.

Sediment biogeochemistry in the woodland lake carried a strong benthic signature (Fig. 3) in accordance with the shallow and nutrient-poor water column. Benthic primary production is often strongly light limited (Vadeboncoeur et al. 2008; Karlsson et al. 2009) and therefore vulnerable to changes in the concentration of colored terrestrial organic carbon (Ask et al. 2012). At low DOC concentrations (Seekell et al. 2015) or optical depths (Vasconcelos et al. 2018), concurrent nutrient stimuli may, however, override the adverse effect of shading on aquatic biota. In effect, earlier research from northern Lapland indicated that terrestrial nutrients may continue to promote primary production in similar shallow (< 2 m) and dark  $(a_{320} > 20 \text{ m}^{-1})$  lakes despite the apparent low light penetration (Rantala et al. 2016). In the woodland lake, the centennial shifts in lake water TOC, whether related to changes in terrestrial carbon transport or photoprocessing, displayed an ambiguous connection to aquatic primary production. This may be attributed to the dual role of terrestrial organic carbon as a nutrient subsidy and shading agent (Fig. 1), as well as to relative changes in allochthonous and autochthonous contributions to the lake water TOC pool. For one, the transient increase in allochthonous TOC  $(\sim 2 \text{ mg L}^{-1})$  mid-core was little reflected in chlorophyll a concentrations suggesting either that the related shading and nutrient effects were cancelling one another out, or that neither was sufficiently large to influence aquatic production. On the other hand, the increase in TOC ( $\sim 1 \text{ mg L}^{-1}$ ) in the topmost three samples coincided with clearly elevated chlorophyll a and could indicate either increased catchment material fluxes, adding allochthonous TOC and promoting lake production, or higher autochthonous contribution to the TOC pool. The latter appears more plausible here in light of the elevated  $\delta^{13}C$  and low C/N values near the surface, although simultaneous increase in aquatic production could be partially masking allochthonous signals. Diagenetic effects could contribute to the chlorophyll a,  $\delta^{13}$ C and C/N shifts towards the surface but should not account for the trends sustained over several decades as diagenetic effects on chlorophyll a (Rydberg et al. 2020), carbon and nitrogen elemental concentrations (Gälman et al. 2008), and carbon isotope composition (Gälman et al. 2009) should primarily take place within the first few years after deposition.

# Aquatic carbon cycling under global change

Mounting evidence shows that the role of northern lakes in the biogeochemical carbon cycle has changed in response to growing human pressure and will continue to transform under global change (Tranvik et al. 2009). Whether this change will be towards increasing carbon sequestration or release is yet difficult to resolve. A prevalent uncertainty here relates to the complex effects of increased productivity and browning of northern waters (Fig. 1) that often tend to push aquatic carbon balances in opposite directions (Ask et al. 2012). Our analyses depicted disparate controls for lake carbon cycling above and below the subarctic treeline under natural climate fluctuations, yet under global change both of the studied lakes displayed a uniform pattern of increasing aquatic primary production (Figs. 5 and 6) in common to numerous other small and shallow lakes spread across the circumpolar north (Michelutti et al. 2005; Kivilä et al. 2020). The change was, however, very subtle in the tundra lake possibly related to the high elevation of the lake that may delay climate impact on productivity (Griffiths et al. 2017). Although several of the multitudinous effects of global change are simultaneously promoting the browning of northern lakes (Creed et al. 2018; Wauthy et al. 2018), we found no evidence of distinct recent increases in terrestrial carbon loading in either of the lakes. While different in many aspects, the two lakes share certain regional and local features that may contribute to their common deviance from the widespread circumpolar pattern. For one, it may be that no substantial catchment greening (Larsen et al. 2011; Finstad et al. 2016) has occurred around the lakes, as suggested by a recent study from the region (within  $\sim 100$  km of the studied lakes) demonstrating conspicuous plant species shifts over the past four decades across the treeline transect, but no clear expansion in trees nor tall shrubs (Vuorinen et al. 2017). Furthermore, earlier studies have proposed that hydrologically isolated lakes could be less prone to growing terrestrial influence (Bogard et al. 2019) and therefore rather reflect more sensitively changes occurring within-lake. Hydrological disconnectedness could thereby also contribute to the uniform response of these two contrasting lakes. Irrespective of causes, the results illustrate how lake responses to global change may not always align with expectations based on landscape position and lake water carbon regime. This corroborates the importance of identifying key environmental factors shaping lake vulnerability and directing their responses to human perturbation within and across regions, for rigorous global upscaling (Seekell et al. 2018; Bogard et al. 2019). Moreover, our study underscores the utility of the long-term sedimentary perspective that can help us identify ecosystem-level responses to growing human pressures and assess the nature and extent of ongoing changes against background variability prior to anthropogenic influence (Smol 2019). Finally, while ambiguities remain as regards the contribution of photodegradation to the elemental and isotopic patterns, the results suggest that we could be overlooking a relevant driver of organic matter biogeochemistry in sedimentary studies. This can be considered a caveat for the already complex elemental and isotopic carbon indices but also an opportunity for paleo-optical research, and either way calls for further experimental research on POM photobiogeochemistry. In regard to lake sediments, additional techniques such as those drawing on molecular biogeochemistry (Ninnes et al. 2017) for better differentiation of terrestrial influence, or paleopigment applications (Nevalainen and Rautio 2014) shedding further light on underwater UV regimes, will be needed to enable a better distinction of the role of photodegradation. Despite the complexity of intertwined environmental signals, biogeochemical archives retained in lake sediments hold valuable information of aquatic carbon regulation in northern lake ecosystems increasingly exposed to global human perturbation.

Acknowledgements This work was supported by the Academy of Finland (#308954, 314107; 335317) and the Kone Foundation (#090140). We thank the staff of the Kilpisjärvi Biological Station (University of Helsinki) for valuable assistance with organizing the field work, and Milla Rautio and Anssi Vähätalo for insights related to the experimental design. Jörg Tittel and two anonymous reviewers are thanked for their thoughtful comments.

Author contributions Liisa Nevalainen, Marttiina Rantala, Henriikka Kivilä and Tomi Luoto lead the conception of the study and carried out the experimental and sediment material acquisition. All authors contributed to sample processing and analyses, in relation to limnological, elemental and isotopic analyses (Marttiina Rantala, Henriikka Kivilä, Tomi Luoto, Liisa Nevalainen), spectral analyses (Carsten Meyer-Jacob, John Smol), and radiometric dating (Antti Ojala). Marttiina Rantala prepared the first draft of the manuscript that was critically reviewed and commented on by all authors. All authors have read and approved the final manuscript.

**Funding** Open access funding provided by University of Helsinki including Helsinki University Central Hospital. This work was supported by the Academy of Finland (#308954, 314107; 335317) and the Kone Foundation (#090140).

**Data availability** All data generated during this study are included in this published article and its supplementary information files.

Code availability Not applicable.

#### Declarations

**Conflict of interest** The author's declare no conflicts of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

#### References

- Ask J, Karlsson J, Jansson M (2012) Net ecosystem production in clear-water and brown-water lakes. Glob Biogeochem Cy 26:GB1017
- Arvela H, Markkanen M, Lemmelä H (1990) Mobile survey of environmental gamma radiation and fall-out levels in Finland after the chernobyl accident. Radiat Prot Dosimetry 32:177–184
- Beardall J, Stojkovic S, Gao K (2014) Interactive effects of nutrient supply and other environmental factors on the sensitivity of marine primary producers to ultraviolet radiation: implications for the impacts of global change. Aquat Biol 22:5–23

- Benner R, Biddanda B (1998) Photochemical transformations of surface and deep marine dissolved organic matter: effects on bacterial growth. Limnol Oceanogr 43:1373–1378
- Benner R, Fogel ML, Sprague KE, Hodson RE (1987) Depletion of 13C in lignin and its implications for stable carbon isotope studies. Nature 329:708–710
- Berggren M, Klaus M, Selvam BP, Ström L, Laudon H, Jansson M, Karlsson J (2018) Quality transformation of dissolved organic carbon during water transit through lakes: contrasting controls by photochemical and biological processes. Biogeosciences 15:457–470
- Bertilsson S, Tranvik LJ (2000) Photochemical transformation of dissolved organic matter in lakes. Limnol Oceanogr 45:753–762
- Bogard MJ, Kuhn CD, Johnston SE, Striegl RG, Holtgrieve GW, Dornblaser MM, Spencer RGM, Wickland KP, Butman DE (2019) Negligible cycling of terrestrial carbon in many lakes of the arid circumpolar landscape. Nat Geosci 12:180–185
- Brodie CR, Leng MJ, Casford JSL, Kendrick CP, Lloyd JM, Yongqiang Z, Bird MI (2011) Evidence for bias in C and N concentrations and  $\delta^{13}$ C composition of terrestrial and aquatic organic materials due to pre-analysis acid preparation methods. Chem Geol 282:67–83
- Buffam I, McGlathery KJ (2003) Effect of ultraviolet light on dissolved nitrogen transformations in coastal lagoon water. Limnol Oceanogr 48:723–734
- Callaghan TV, Bergholm F, Christensen TR, Jonasson C, Kokfelt U, Johansson M (2010) A new climate era in the sub-Arctic: Accelerating climate changes and multiple impacts. Geophys Res Lett 37:L14705
- Chen M, Jaffé R (2016) Quantitative assessment of photo- and bio-reactivity of chromophoric and fluorescent dissolved organic matter from biomass and soil leachates and from surface waters in a subtropical wetland. Biogeochemistry 129:271–289
- Chomicki K (2009) The use of stable carbon and oxygen isotopes to examine the fate of dissolved organic matter in two small, oligotrophic Canadian Shield lakes. Dissertation, University of Waterloo.
- Cory RM, Harrold KH, Neilson BT, Kling GW (2015) Controls on dissolved organic matter (DOM) degradation in a headwater stream: the influence of photochemical and hydrological conditions in determining light-limitation or substrate-limitation of photo-degradation. Biogeosciences 12:6669–6685
- Cory RM, Kling GW (2018) Interactions between sunlight and microorganisms influence dissolved organic matter degradation along the aquatic continuum. Limnol Oceanogr Lett 3:102–116
- Creed IF, Bergström A, Trick CG, Grimm NB, Hessen DO, Karlsson J, Kidd KA, Kritzberg E, McKnight DM, Freeman EC, Senar OE, Andersson A, Ask J, Berggren M, Cherif M, Giesler R, Hotchkiss ER (2018)
- Finstad AG, Andersen T, Larsen S, Tominaga K, Blumentrath S, de Wit HA, Tømmervik H, Hessen DO (2016) From greening to browning: catchment vegetation development and reduced S-deposition promote organic carbon load on decadal time scales in Nordic lakes. Sci Rep 6:31944

- Gälman V, Rydberg J, Bigler C (2009) Decadal diagenetic effects on  $\delta^{13}$ C and  $\delta^{15}$ N studied in varved lake sediment. Limnol Oceanogr 54:917–924
- Gälman V, Rydberg J, Sjöstedt de-Luna S, Bindler R, Renberg I (2008) Carbon and nitrogen loss rates during aging of lake sediments: changes over 27 years studied in varved lake sediment. Limnol Oceanogr 53:1076–1082
- Griffiths K, Michelutti N, Sugar M, Douglas MSV, Smol JP (2017) Ice-cover is the principal driver of ecological change in High Arctic lakes and ponds. PLoS One 12:e0172989
- Hodell DA, Schelske CL (1998) Production, sedimentation, and isotopic composition of organic matter in Lake Ontario. Limnol Oceanogr 43:200–214
- Karlsson J, Byström P, Ask J, Ask P, Persson L, Jansson M (2009) Light limitation of nutrient-poor lake ecosystems. Nature 460:506–509
- Kellerman AM, Kothawala DN, Dittmar T, Tranvik LJ (2015) Persistence of dissolved organic matter in lakes related to its molecular characteristics. Nat Geosci 8:454–459
- Kivilä EH, Luoto TP, Rantala MV, Nevalainen L (2020) Late-Holocene variability in chironomid functional assemblages and carbon utilization in a tundra lake food web. Hydrobiologia 847:895–911
- Koehler B, Landelius T, Weyhenmeyer GA, Machida N, Tranvik LJ (2014) Sunlight-induced carbon dioxide emissions from inland waters. Glob Biogeochem Cy 28:696–711
- Lapierre J, Guillemette F, Berggren M, del Giorgio PA (2013) Increases in terrestrially derived carbon stimulate organic carbon processing and CO2 emissions in boreal aquatic ecosystems. Nat Commun 4:2972
- Larsen S, Andersen T, Hessen DO (2011) Climate change predicted to cause severe increase of organic carbon in lakes. Glob Change Biol 17:1186–1192
- Laurion I, Vincent WF, Lean DRS (1997) Underwater ultraviolet radiation: development of spectral models for northern high latitude lakes. Photochem Photobiol 65:107–114
- Luoto TP, Kivilä EH, Rantala MV, Nevalainen L (2017) Characterization of the medieval climate anomaly, little ice age and recent warming in northern Lapland. Int J Climatol 37:1257–1266
- Magnuson JJ, Robertson DM, Benson BJ, Wynne RH, Livingstone DM, Arai T, Assel RA, Barry RG, Card V, Kuusisto E, Granin NG, Prowse TD, Stewart KM, Vuglinski VS (2000) Historical trends in lake and river ice cover in the Northern Hemisphere. Science 289:1743–1746
- Marwood CA, Smith REH, Furgal JA, Charlton MN, Solomon KR, Greenberg BM (2000) Photoinhibition of natural phytoplankton assemblages in Lake Erie exposed to solar ultraviolet radiation. Can J Fish Aquat Sci 57:371–379
- Mayer LM, Schick LL, Bianchi TS, Wysocki LA (2009) Photochemical changes in chemical markers of sedimentary organic matter source and age. Mar Chem 113:123–128
- Meyer-Jacob C, Michelutti N, Paterson AM, Monteith D, Yang H, Weckström J, Smol JP, Bindler R (2017) Inferring past trends in lake water organic carbon concentrations in northern lakes using sediment spectroscopy. Environ Sci Technol 51:13248–13255
- Meyer-Jacob C, Michelutti N, Paterson AM, Cumming BF, Keller W, Smol JP (2019) The browning and re-browning

of lakes: divergent lake-water organic carbon trends linked to acid deposition and climate change. Sci Rep 9:16676

- Meyers PA, Teranes JL (2001) Sediment organic matter. In: Last WM, Smol JP (eds) Tracking environmental change using lake sediments, vol 2. Physical and geochemical techniques. Kluwer Academic Publisher, Dordrecht, The Netherlands, pp 239–269
- Michelutti N, Blais JM, Cumming BF, Paterson AM, Rühland K, Wolfe AP, Smol JP (2010) Do spectrally inferred determinations of chlorophyll a reflect trends in lake trophic status? J Paleolimnol 43:205–217
- Michelutti N, Smol JP (2016) Visible spectroscopy reliably tracks trends in paleo-production. J Paleolimnol 56:253–265
- Michelutti N, Wolfe AP, Vinebrooke RD, Rivard B, Briner JP (2005) Recent primary production increases in arctic lakes. Geophys Res Lett 32:L19715
- Moran MA, Sheldon WM, Zepp RG (2000) Carbon loss and optical property changes during long-term photochemical and biological degradation of estuarine dissolved organic matter. Limnol Oceanogr 45:1254–1264
- Nevalainen L, Rantala MV, Kivilä EH, Wauthy M, Rautio M, Luoto TP (2020) Biogeochemical and photobiological responses of subarctic lakes to UV radiation. J Photochem Photobiol B: Biol 209:111932
- Nevalainen L, Rantala MV, Meyer-Jacob C, Kivilä EH, Ojala AEK, Smol JP, Luoto TP (2020) A Holocene record of aquatic bio-optics in subarctic Fennoscandia. Quat Sci Rew 243:106491
- Nevalainen L, Luoto TP, Rantala MV, Galkin A, Rautio M (2015) Role of terrestrial carbon in aquatic UV exposure and photoprotective pigmentation of meiofauna in subarctic lakes. Freshw Biol 60:2435–2444
- Nevalainen L, Rautio M (2014) Spectral absorbance of benthic cladoceran carapaces as a new method for inferring past UV exposure of aquatic biota. Quat Sci Rev 84:109–115
- Ninnes S, Tolu J, Meyer-Jacob C, Mighall TM, Bindler R (2017) Investigating molecular changes in organic matter composition in two Holocene lake-sediment records from central Sweden using pyrolysis-GC/MS. J Geophys Res Biogeosci 122:1423–1438
- Ojala AEK, Luoto TP, Virtasalo JJ (2017) Establishing a highresolution surface sediment chronology with multiple dating methods - testing <sup>137</sup>Cs determination with Nurmijärvi clastic-biogenic varves. Quat Geochron 37:32–41
- Ophsal S, Benner R (1998) Photochemical reactivity of dissolved lignin in river and ocean waters. Limnol Oceanogr 43:1297–1304
- Ophsal SP, Zepp RG (2001) Photochemically-induced alteration of stable carbon isotope ratios ( $\delta^{13}$ C) in terrigenous dissolved organic carbon. Geophys Res Lett 28:2417–2420
- Oswald WW, Anderson PM, Brown TA, Brubaker LB, Hu FS, Lozhkin AV, Tinner W, Kaltenrieder P (2005) Effects of sample mass and macrofossil type on radiocarbon dating of arctic and boreal lake sediments. Holocene 15:758–767
- Rantala MV, Luoto TP, Weckström J, Rautio M, Nevalainen L (2017) Climate drivers of diatom distribution in shallow subarctic lakes. Freshw Biol 62:1971–1985
- Rantala MV, Nevalainen L, Rautio M, Galkin A, Luoto TP (2016) Sources and controls of organic carbon in lakes

across the subarctic treeline. Biogeochemistry 129:235–253

- Rautio M, Dufresne F, Laurion I, Bonilla S, Vincent WF, Christoffersen KS (2011) Shallow freshwater ecosystems of the circumpolar arctic. Écoscience 18:204–222
- Rose KC, Williamson CE, Saros JE, Sommaruga R, Fischer JM (2009) Differences in UV transparency and thermal structure between alpine and subalpine lakes: implications for organisms. Photochem Photobiol Sci 8:1244–1256
- Rydberg J, Cooke CA, Tolu J, Wolfe AP, Vinebrooke RD (2020) An assessment of chlorophyll preservation in lake sediments using multiple analytical techniques applied to the annually laminated lake sediments of Nylandssjön. J Paleolimnol 64:379–388
- Schlacher TA, Connolly RM (2014) Effects of acid treatment on carbon and nitrogen stable isotope ratios in ecological samples: a review and synthesis. MEE 5:541–550
- Schmitt-Kopplin P, Hertkorn N, Schulten H-R, Kettrup A (1998) Structural changes in dissolved soil humic acid during photochemical degradation processes under O<sub>2</sub> and N<sub>2</sub> atmosphere. Environ Sci Technol 32:2531–2541
- Seekell DA, Lapierre J, Cheruvelil KS (2018) A geography of lake carbon cycling. Limnol Oceanogr Lett 3:49–56
- Seekell DA, Lapierre J, Ask J, Bergström A, Deininger A, Rodríguez P, Karlsson J (2015) The influence of dissolved organic carbon on primary production in northern lakes. Limnol Oceanogr 60:1276–1285
- Smol JP (2019) Under the radar: long-term perspectives on ecological changes in lakes. Proc R Soc B 286:20190834
- Smol JP, Douglas MSV (2007) From controversy to consensus: making the case for recent climate change in the Arctic using lake sediments. Front Ecol Environ 5:466–474
- Smol JP, Wolfe AP, Birks HJB, Douglas MSV, Jones VJ, Korhola A, Pienitz R, Rühland K, Sorvari S, Antoniades D, Brooks SJ, Fallu M, Hughes M, Keatley BE, Laing TE, Michelutti N, Nazarova L, Nyman M, Paterson AM, Perren B, Quinlan R, Rautio M, Saulnier-Talbot E, Siitonen S, Solovieva N, Weckström J (2005) Climate-driven regime shifts in the biological communities of arctic lakes. PNAS 102:4397–4402
- Solomon CT, Jones SE, Weidel BC, Buffam I, Fork ML, Karlsson J, Larsen S, Lennon JT, Read JS, Sadro S, Saros JE (2015) Ecosystem consequences of changing inputs of terrestrial dissolved organic matter to lakes: current knowledge and future challenges. Ecosystems 18:376–389
- Sulzberger B, Austin AT, Cory RM, Zepp RG, Paul ND (2019) Solar UV radiation in a changing world: roles of cryosphere–land–water atmosphere interfaces in global biogeochemical cycles. Photochem Photobiol Sci 18:747–774
- Talbot MR (2001) Nitrogen isotopes in palaeolimnology. In: Last WM, Smol JP (eds) Tracking environmental change using lake sediments, vol 2. Physical and geochemical methods. Kluwer Academic Publishers, Dordrecht, pp 401–439
- Tranvik LJ, Downing JA, Cotner JB, Loiselle SA, Striegl RG, Ballatore TJ, Dillon P, Finlay K, Fortino K, Knoll LB, Kortelainen PL, Kutser T, Larsen S, Laurion I, Leech DM, McCallister LS, McKnight DM, Melack JM, Overholt E, Porter JA, Prairie Y, Renwick WH, Sherman BS, Schindler DW, Sobek S, Tremblay A, Vanni MJ, Verschoor AM, Ev W, Weyhenmeyer GA (2009) Lakes and reservoirs as

regulators of carbon cycling and climate. Limnol Oceanogr 54:2298–2314

- Tranvik LJ, Jansson M (2002) Climate change: Terrestrial export of organic carbon. Nature 415:861–862
- Vadeboncoeur Y, Peterson G, Vander Zanden JM, Kalff J (2008) Benthic algal production across lake size gradients: interactions among morphometry, nutrients, and light. Ecology 89:2542–2552
- Vähätalo A, Salonen K, Salkinoja-Salonen M, Hatakka A (1999) Photochemical mineralization of synthetic lignin in lake water indicates enhanced turnover of aromatic organic matter under solar radiation. Biodegradation 10:415–420
- Vähätalo AV, Wetzel RG (2008) Long-term photochemical and microbial decomposition of wetland-derived dissolved organic matter with alteration of <sup>13</sup>C: <sup>12</sup>C mass ratio. Limnol Oceanogr 53:1387–1392
- Vasconcelos FR, Diehl S, Rodríguez P, Karlsson J, Byström P (2018) Effects of terrestrial organic matter on aquatic primary production as mediated by pelagic-benthic resource fluxes. Ecosystems 21:1255–1268
- Vähätalo AV (2009) Light, photolytic reactivity and chemical products. In: Likens GE (ed) Encyclopedia of Inland Waters. Academic Press, pp 761–773
- Vincent WF, Rautio M, Pienitz R (2007) Climate control of biological UV exposure in polar and alpine aquatic ecosystems. In: Ørbaek JV, Kallenborn R, Tombre I, Hegseth EN, Falk-Petersen S, Hoel AH (eds) Arctic alpine ecosystems and people in a changing environment. Springer, Berlin, pp 227–249
- Vuorinen KEM, Oksanen L, Oksanen T, Pyykönen A, Olofsson J, Virtanen R (2017) Open tundra persist, but arctic features decline—vegetation changes in the warming Fennoscandian tundra. Glob Change Biol 23:3794–3807

- Wauthy M, Rautio M, Christoffersen KS, Forsström L, Laurion I, Mariash HL, Peura S, Vincent WF (2018) Increasing dominance of terrigenous organic matter in circumpolar freshwaters due to permafrost thaw. Limnol Oceanogr Lett 3:186–198
- Weckström J, Korhola A, Erästö P, Holmström L (2006) Temperature patterns over the past eight centuries in Northern Fennoscandia inferred from sedimentary diatoms. Quat Res 66:78–86
- Williamson CE, Overholt EP, Brentrup JA, Pilla RM, Leach TH, Schladow SG, Warren JD, Urmy SS, Sadro S, Chandra S, Neale PJ (2016) Sentinel responses to droughts, wildfires, and floods: effects of UV radiation on lakes and their ecosystem services. Front Ecol Environ 14:102–109
- Williamson CE, Brentrup JA, Zhang J, Renwick WH, Hargreaves BR, Knoll LB, Overholt EP, Rose KC (2014) Lakes as sensors in the landscape: Optical metrics as scalable sentinel responses to climate change. Limnol Oceanogr 59:840–850
- Williamson CE, Stemberger RS, Morris DP, Frost TM, Paulsten SG (1996) Ultraviolet radiation in North American lakes: Attenuation estimates from DOC measurements and implications for plankton communities. Limnol Oceanogr 41:1024–1034
- Wolfe AP, Cooke CA, Hobbs WO (2006) Are current rates of atmospheric nitrogen deposition influencing lakes in the Eastern Canadian Arctic? Arct Antarct Alp Res 38:465–476

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.