



Peatland restoration pathways to mitigate greenhouse gas emissions and retain peat carbon

Ülo Mander · Mikk Espenberg · Lulie Melling ·
Ain Kull

Received: 21 June 2023 / Accepted: 4 November 2023 / Published online: 8 December 2023
© The Author(s) 2023

Abstract Peatlands play a crucial role in the global carbon (C) cycle, making their restoration a key strategy for mitigating greenhouse gas (GHG) emissions and retaining C. This study analyses the most common restoration pathways employed in boreal and temperate peatlands, potentially applicable in tropical peat swamp forests. Our analysis focuses on the GHG emissions and C retention potential of the restoration measures. To assess the C stock change in restored (rewetted) peatlands and afforested peatlands with continuous drainage, we adopt a conceptual approach that considers short-term C capture (GHG exchange between the atmosphere and the peatland ecosystem) and long-term C sequestration in peat. The primary criterion of our conceptual model is the capacity of restoration measures to capture C and reduce GHG emissions. Our findings indicate that carbon dioxide (CO_2) is the most influential part of long-term

climate impact of restored peatlands, whereas moderate methane (CH_4) emissions and low N_2O fluxes are relatively unimportant. However, lateral losses of dissolved and particulate C in water can account up to a half of the total C stock change. Among the restored peatland types, Sphagnum paludiculture showed the highest CO_2 capture, followed by shallow lakes and reed/grass paludiculture. Shallow lakeshore vegetation in restored peatlands can reduce CO_2 emissions and sequester C but still emit CH_4 , particularly during the first 20 years after restoration. Our conceptual modelling approach reveals that over a 300-year period, under stable climate conditions, drained bog forests can lose up to 50% of initial C content. In managed (regularly harvested) and continuously drained peatland forests, C accumulation in biomass and litter input does not compensate C losses from peat. In contrast, rewetted unmanaged peatland forests are turning into a persistent C sink. The modelling results emphasized the importance of long-term C balance analysis which considers soil C accumulation, moving beyond the short-term C cycling between vegetation and the atmosphere.

Responsible Editor: Klaus Butterbach-Bahl.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10533-023-01103-1>.

Ü. Mander · M. Espenberg · A. Kull
Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia
e-mail: ulo.mander@ut.ee

L. Melling
Sarawak Tropical Peat Research Institute, Kuching, Sarawak, Malaysia

Keywords Carbon dioxide · Methane · Nitrous oxide · Paludiculture · Peatland restoration · Rewetting

Introduction

Importance of peatlands in a changing climate

Peatlands cover only about 3% of the Earth's terrestrial surface (Gorham 1991) but play a crucial role in the global carbon (C) cycle. They act as significant C stores and are sources or sinks for greenhouse gases (GHG) like carbon dioxide (CO_2), methane (CH_4) and nitrous oxide (N_2O ; Frolking et al. 2011). These gases contribute to climate change and are considered crucial anthropogenic GHGs. Due to their high C density, peatlands are globally recognized as vital C reservoirs (Gallego-Sala et al. 2018) accounting for about 21% of the global soil organic C stock, estimated $\sim 3000 \text{ Pg}$ (Leifeld and Menichetti 2017). Peatlands also serve as substantial stores of organic N, with Northern peatlands alone accumulating 8–15 Pg N. When including tropical peatlands, the global estimations reached up to 26 Pg N (Swenson et al. 2019).

Undisturbed peatlands are currently a C sink ($\sim 0.1 \text{ Pg C yr}^{-1}$), a moderate source of CH_4 ($\sim 0.03 \text{ Pg CH}_4 \text{ yr}^{-1}$), and a very weak source of N_2O ($\sim 0.00002 \text{ Pg N}_2\text{O-N yr}^{-1}$) (Frolking et al. 2011). However, anthropogenic activities, primarily agriculture and forestry drainage (up to 20% of global peatlands), result in net CO_2 emissions ($\sim 0.1 \text{ Pg C yr}^{-1}$), reduced CH_4 emissions (10% smaller than in natural conditions), and increased N_2O emissions (~ 20 times higher than in natural peatlands). Consequently, subsidence and soil degradation contribute nearly 6% of global anthropogenic GHG emissions (Wichtmann et al. 2016). Most likely, the global peatland's GHG balance has turned to a C source, a slightly smaller CH_4 source, and a larger (but still small) N_2O source (Frolking et al. 2011). In Europe, 46% of the remaining peatlands have degraded to the point where peat is no longer actively forming (Swenson et al. 2019).

As typical wetlands, peatlands are severely threatened by drainage, climate change, fires and groundwater extraction (Fluet-Chouinard et al. 2023). However, their restoration is beneficial, enabling C capture and sequestration, and minimizing N_2O emissions (Leifeld and Menichetti 2017).

The northern hemisphere has experienced the highest warming during winter and early spring (Ljungqvist et al. 2016), leading to more frequent freeze-thaw cycles. Annual precipitation has also increased, particularly during the cold half-year in

northern regions (Ljungqvist et al. 2016). Conversely, there are many indications of growing frequency of droughts during the warm season (Chiang et al. 2021). An increase in flash floods is also predicted (Zheng et al. 2022). These changing climatic conditions, droughts, rapid fluctuations in groundwater level (Mander et al. 2021), flash floods (Schindler et al. 2020), and soil moisture conditions (Pärn et al. 2018; Evans et al. 2021; Huang et al. 2021) can create hot spots and hot moments of GHG emission in peatland ecosystems.

Peatland restoration

Rewetting is a crucial step for conservation or sequestration of C in peatlands previously affected by drainage (Günther et al. 2020). Likewise, rewetting can affect nitrogen cycle and reduce cumulative $\text{N}_2\text{O-N}$ emissions by up to 70% in European peatlands (Liu et al. 2022). Usually, peatland restoration and rewetting are considered as synonyms, but in this paper we consider rewetting as increasing water table level compared with previously drained status. In case of restoration we expect that both water table and vegetation are manipulated in the way that enables to achieve ecosystem status similar to pre-drainage. Restoration pathways depend on factors such as initial vegetation of the drained area (Heger et al. 2022; Schaller et al. 2022), nutrient status of the residual peat layer (Kreylig et al. 2021) and the expected time frame for achieving planned ecological and socioeconomic benefits of the restoration.

Rewetting may transform former peat extraction sites into mires, paludicultural land, wet forests or shallow waterbodies. Ecologically, the preferred pathway is to restore them as mires (Wilson et al. 2022), which can be achieved by restoring water level, by a combination of rewetting and the application of peat moss layer transfer technique (Gonzalez-Sargas & Rochefort 2019), or to establish shallow waterbodies in hydrologically complex sites (Christen et al. 2016).

Peatland rewetting for agricultural use (including paludiculture) is challenging but may have a positive short-term effect on CO_2 capture and GHG mitigation (Maljanen et al. 2010; Wilson et al. 2016a, b) but may not stop long-term peat mineralisation. Rewetting for paludiculture (Wichtmann et al. 2016) usually encompasses *Sphagnum*-based C farming (Gaudig et al. 2017/2018), energy crop cultivation (Hyvönen

et al. 2009; Mander et al. 2012; Järveoja et al. 2016a; Kandel et al. 2020); wet forestry (Anadon-Rosell et al. 2022), or cultivation of the reed (Martens et al. 2021) or cattail (De Jong et al. 2021). Berries and other wetland plants are also suitable for paludiculture, but cranberry (*Vaccinium*) is the most suitable pioneer species for mire restoration and long-term C capture (Freeman et al. 2022).

Restoration of drained forests is usually achieved by blocking ditches and restoring water levels (Grand-Clement et al. 2015). Regulating the water regime in restored sites is effective for mitigation of GHG emissions (Järveoja et al. 2016b).

Peatland restoration is vulnerable to hydroclimatic conditions, particularly in the temperate zone. Changing precipitation pattern, increasing temperature and decreasing snowpack are expected to contribute to more frequent extreme events like droughts and torrential rains, resulting in increased vulnerability and interannual variability (Alm et al. 1999; Drollinger et al. 2019). In addition, it is also important to consider the potential substantial losses of dissolved and particulate C from drained and restored peatlands when estimating C budgets (Billett et al. 2010; Ross et al. 2022).

Due to global warming, northern peatlands are projected to experience increased GHG emissions, particularly during non-growing period (Rafat et al. 2021), while Garcin et al. (2022) highlight a lack of knowledge of hydroclimatic vulnerability of peat C in tropical peatlands. Wet-dry seasonality of GHG fluxes is expected from tropical peatland soils (Inubushi et al. 2003). However, the overall impact of climate change on GHG fluxes needs to be better understood. Current understanding suggests that changes in soil temperature, photosynthesis, and soil moisture drive alterations in net C fluxes (Rafat et al. 2021).

Restoration versus afforestation of peatlands

Peatland restoration involves rewetting, whereas afforestation of drained peatlands and maintaining their drained condition cannot be considered equivalent to restoration. The key issue lies in the difference in the time scale and the discrepancy in distinguishing between short-term C capture in ecosystem (often observed in studies on GHG exchange between the atmosphere and ecosystem) and long-term C sequestration in soil. When peatlands are drained, C loss

from the peat can offset the benefit of long-term CO₂ sequestration achieved by afforestation (Jurasinski et al. 2020, 2023).

Currently, there is insufficient evidence on the long-term benefits of active afforestation of drained peatlands to mitigate climate change (Jurasinski et al. 2023). Afforestation on drained peatland forests and some former peat extraction areas may provide short-term benefits for climate change mitigation (Mäkiranta et al. 2007; Samariks et al. 2023). However, this approach does not account for the value of long-term C storage in peat. Similarly, intensive forestry on drained peatlands will not restore the peatland ecosystem's flora, fauna, and functions (Haapalehto et al. 2017; Loisel & Gallego-Sala 2022).

In addition, afforested drained peatlands are more susceptible to wildfires (Kohlenberg et al. 2018; Zheng et al. 2023). These risks are exacerbated by more frequent and more intense droughts in the boreal zone (Walker et al. 2019), resulting in losses of burnt wood and substantial C losses from burnt and burning peat layers (Liu et al. 2023). The impacts of severe fires have been devastating drained areas of formerly tropical peatland forests in Southeast Asia (Page et al. 2009).

Restoration-versus-afforestation of peatlands is being debated during the legislation procedure of the European Union Nature Restoration Law (NRL) (Jurasinski et al. 2023).

The general aim of this paper is to assess short-term C capture (GHG exchange between the atmosphere and the peatland ecosystem) and analyse the fluxes in the context of long-term C sequestration in peat of restored (rewetted) peatlands and afforested drained peatlands. As a specific objective we implement a conceptual model that compares changes in the long-term C budget and climate warming mitigation potential in the restored and afforested peatlands.

Conceptual framework

To provide a comprehensive understanding of the framework and to organize the rich source material, we have adopted a three-stage system. These stages refer to peatlands with different water regimes: natural (pristine), drained, and restored (rewetted) ecosystems. Within each stage of water regime, we further divided them into blocks based on primary

land use types (Fig. 1). Estimated fluxes of all three GHGs—CO₂, CH₄ and N₂O—are presented using a general three tier scale (high, medium and low fluxes). Flux values are represented by arrows, with upward-pointing arrows indicating emission and downward-pointing arrows indicating capture or uptake of the corresponding gas. In addition, we have indicated the averaged lateral losses of dissolved organic C in water (Fig. 1).

Although our main focus lies on temperate and boreal peatlands, we also briefly discuss tropical peatland forests due to their significance as global hot spots of GHGs, and their large-scale escalating disturbance and destruction. In characterizing GHG fluxes in natural and drained peatlands, we primarily relied on literature sources supplemented by unpublished original data from our research group. However, for




Fig. 1 Conceptual figure showing main generalized types of peatland ecosystems from natural and drained to potentially restored peatlands. Proposed set-up for greenhouse gas emission from the main compartments of ecosystems is shown. Arrows indicate literature-based mean values of greenhouse gases CO₂, CH₄ and N₂O fluxes (upward and downward arrows indicate emissions and captures of the corresponding gas,

respectively). Question marks indicate scarce data without representative values. Aggregated lateral losses of dissolved organic carbon (DOC) in water are based on Wilson et al. 2016a. Rewetting is displayed as a necessary presumption for peatland restoration. Paludiculture—Sphagnum, berries and energy crops. Open restored areas – bog- and fen-type peatlands

tropical wetlands, data is limited or sporadic, resulting in the use of question marks.

For restored (rewetted) peatlands, due to the scarcity and less systematic nature of the data, we present a selection of available GHG flux data in Table 1. The categories of restored (rewetted) peatlands are as in Fig. 1: wet forests, paludiculture, shallow lakes and open mires. Upon no accurate distinction between the mentioned categories in the source data we added a general category – rewetted peatlands – to both Table 1 and Fig. 2. The latter one presents average flux values (Fig. 2A–C) and GHG balances in CO₂ equivalents (Fig. 2D and Fig. 2E). In the more general Fig. 1, paludiculture is shown as a single category. To compile this information, we referred to both literature sources and results from local research projects in Estonia. For our literature search, we utilized well-known indexing systems such as Web of Science (Clarivate Analytics), Scopus, and Google Scholar. We used following keywords and their combinations: “rewetted peatland(s)”, “restored wetland(s)”, “restored peatland(s)”, “rewetted forest(s)”, “paludiculture”, “Sphagnum farming”, “greenhouse gases”, “carbon dioxide”, “methane”, “nitrous oxide”, “tropical peatlands”. In addition, we looked for references from cited rewetting-related papers.

To explain the dilemma of peatland restoration and afforestation in a long-term perspective and mitigation of climate warming, we created a conceptual model that helps to characterize the changes in the C budget under different management practices (Fig. 3).

The resulting climatic effects of these management options are strongly time-dependent and rely on difference in forming above or below ground carbon pool and carbon turnover rate.

Our conceptual model of long-term C stock dynamics over 300-year period is based on the approach proposed by Minkkinen & Laine (1998). We developed it further to include C stock dynamics in nutrient-poor mires (open bog and bog forest), drained bog forest, and potential land use scenarios after milled peat extraction (restored bog, naturally regenerated unmanaged forest and managed forest). In modelling, we assumed equal C stock 686 t ha⁻¹ in any peatland as initial status, corresponding to C stock with 2 m peat depth in pristine bog (Supplementary Table 1). As approximation, the stem, branches and coarse root biomass were assumed to have stable ratio (branches 12% of stem biomass and

coarse roots 19% of above ground biomass) in any age group, wood density (kg m³) and C content (%) values were derived from literature (Kask and Pikk 2009; Külla 1997). Initial value of stem biomass was set 3 m³ ha⁻¹ for pristine open bog, 100 m³ ha⁻¹ yr⁻¹ for pristine bog forest and 0 m³ ha⁻¹ for any other land use classes (initially harvested peatland forest or treeless peat extraction site) with annual stem biomass increment by 0 m³ ha⁻¹ yr⁻¹ for open bog, 0.1 m³ ha⁻¹ yr⁻¹ for stable continuous cover bog forest, 4 m³ ha⁻¹ yr⁻¹ for drained peatland forest and naturally regenerated forest on peat extraction site (initial value 0 m³ ha⁻¹ yr⁻¹ until year 20, thereafter gradually increasing until 4 m³ ha⁻¹ yr⁻¹ by year 50 and starting to decline to 0.1 m³ ha⁻¹ yr⁻¹ by year 160 as equal to pristine bog forest) while mean annual increment of 6 m³ ha⁻¹ yr⁻¹ was assigned to managed forest on drained former peat extraction site because of higher fertility of deeper peat layers (usually fen peat) after removal of Sphagnum peat. In pristine bog forest, restored bog and unmanaged peatland forest on former peat extraction site no harvest and timber removal is considered. Harvesting cycle is assumed to be 100 years in both restored managed peatland forest and drained managed peatland forests, while only stem biomass is removed. Fine root biomass was considered as part of C turnover and part of net ecosystem exchange (NEE). NEE values for modelled ecosystems are based on original data from Estonian study sites (Table 1), Estonian national GHG inventory, and literature (Salm et al. 2012; Wilson et al. 2016a). A conceptual scheme presents modelled long-term C stock dynamics in pristine mire ecosystems, restored and afforested drained peatlands (Fig. 3).

Results and discussion

Greenhouse gas fluxes in restored peatlands

Figure 1 illustrates a conceptual view of estimated GHG fluxes in natural, drained and restored (rewetted) peatlands. In restored peatlands, CO₂ fluxes play the main role in climate impacts, while CH₄ emissions are moderate and N₂O fluxes are low making them less significant. Lateral losses of dissolved and particulate C in water can account up to a half of total C budget. Among the different restored peatlands, paludiculture has the highest estimated carbon dioxide

Table 1 Greenhouse gas fluxes in main types of boreal and temperate rewetted peatlands

Type	Name	Coordinates	Country	Period of study	Nutrient status	Plant community/ treatment	NEE (CO ₂ -C)	CH ₄ -C	N ₂ O-N	Source
Rewetted peatlands	Twitchell Island old wetland (26 yrs), former agricultural land	38°06'N, 121°38'W	USA, CA	Aug. 2012–Aug. 2013	Rich	<i>Schoenoplectus acutus</i> , <i>Typha latifolia</i> , <i>T. domingensis</i> , <i>T. angustifolia</i> , <i>Ludwigia peploides</i> , <i>Lemna</i> sp.)	−3970	387		Knox et al. 2015
	Sherman Island young wetland, 13 yrs), former peatland pasture	38°05'N, 121°42'W	USA, CA	March 2012–Apr. 2013	Rich	<i>Schoenoplectus acutus</i> , <i>Typha</i> spp.,	−3680	530		Knox et al. 2015
Boreal zone aggregated			Global review			All rewetted peatlands (restored from drained forests, grasslands and croplands, and peat extraction sites)	−1300	123.6	0	Günther et al. 2020
Temperate zone aggregated	Rewetted organic soils		Global review Germany			All rewetted peatlands (see above)	−400	205.9	0	Günther et al. 2020
						Average of both nutrient-poor and nutrient-rich rewetted organic soils	−0.4	279	0.1	Tiemeyer et al. 2020
Rewetted peatlands			Global review			Average of 20 literature sources	675	131	1.2	Bianchi et al. 2021
Rewetted peatland forest	Rewetted boreal forests		Global review	Before 2016	Poor	Coniferous and mixed peatland forests (soil fluxes) Selected examples from IPCC 2014 Wetlands Supplement	−1520	73	0.16	Wilson et al. 2016a
						Coniferous and mixed peatland forests (soil fluxes)	−1930	220	0.16	Wilson et al. 2016a

Table 1 (continued)

Type	Name	Coordinates	Country	Period of study	Nutrient status	Plant community/ treatment	NEE (CO ₂ -C)	CH ₄ -C	N ₂ O-N	Source
Rewetted temperate forests		Global review		Before 2016	Poor	Coniferous, deciduous and mixed peatland forests (soil fluxes)	-1220	160	0.16	Wilson et al. 2016a
28 rewetted peatland forests		Finland	1 year (2008–2019)	Rich	Fen forests (soil fluxes)	960	418	0.16	Wilson et al. 2016a	
Wöpkendorf <i>Alnus glutinosa</i>	54°7'36"N, 12°29'04"E	Germany	May 2018–Apr. 2020	Rich	Bog forests (soil fluxes)			0.8	Minkkinen et al. 2020	
					Rewetted peatland alder forest; <i>Carex acutiformis</i> , <i>C. riparia</i> , <i>Houttonia palustris</i> , <i>Solanum dulcamara</i>	15 to 565		0.5	Köhne et al. 2021	
Paludiculture										
<i>Reed/grasses</i>	Lavassaare <i>Phalaris</i>	58°34'20"N, 24°23'15"E	Estonia	May 2010–May 2011	Rich	<i>Phalaris arundinacea</i> fertilized	3.56	-0.006		Mander et al. 2012
					Rich	<i>P. arundinacea non-fertilized</i>	1.75	-0.03		Mander et al. 2012
				Jan-Dec 2014	Rich	<i>P. arundinacea fertilized</i>	790	0.14	2.39	Järveoja et al. 2016a
					Rich	<i>P. arundinacea non-fertilized</i>	2010	0.18	1.08	Järveoja et al. 2016a
Keressaare <i>Phalaris</i>	58°36'56"N, 27°00'36"E	Estonia	Apr. 2015–Mar. 2018	Poor	<i>P. arundinacea</i> high water table, fertilized, lined	-892	1.14	19.0	Maddison et al. 2018	
					<i>P. arundinacea</i> low water table, fertilized, lined	-249	3.07	28.1	Maddison et al. 2018	
Gersloot <i>Typha latifolia</i>	53°01'N, 5°55"E	Netherlands	Feb–June 2017	Rich	Fen peat, lab experiments		72.9	2.18	Vroom et al. 2018	

Table 1 (continued)

Type	Name	Coordinates	Country	Period of study	Nutrient status	Plant community/ treatment	NEE (CO ₂ -C)	CH ₄ -C	N ₂ O-N	Source
Øby <i>P. arundinacea</i> , <i>Poa</i> spp	56°27'32"N, 9°40'40"E	Denmark	Mar. 2015–Mar. 2017	Rich	<i>P. arundinacea</i> flooded plots	–3970 to –5970	450 to 1,110	4.0 to 5.5	Kandl et al. 2020	
Emergent crops		Global review	Rich	<i>P. arundinacea</i> semi-flooded plots	–2650 to –4990	160 to 710	4.0 to 6.0	Kandl et al. 2020		
<i>Vaccinium</i> spp.	46°15'N, 72°00'W	Canada, Quebec	Growing season 2012, 2013	Poor	Average of 6 literature sources	174	1.68	Bianchi et al. 2021		
Maima	58°35'54"N, 24°22'36"E	Estonia	(a)	Poor	Cultivated cranberry (Vaccinium macrocarpon)	27	0.61	Lloyd et al. 2019		
Laiuse	58°47'17"N, 26°31'47"E	Estonia	(a)	Poor	Naturally regenerated cranberry (<i>Oxycoccus palustris</i>)	–799	68.5	2.0	Burdun et al. 2021	
Ess-soo	57°54'51"N, 26°41'26"E	Estonia	(a)	Poor	Naturally regen. Cranberry (<i>Oxycoccus palustris</i>)	–837	31.4	–0.002	Viru et al. 2020; Burdun et al. 2021	
<i>Sphagnum</i> spp	53°02' N, 7°29'E	Germany	June 2010–Dec. 2011	Poor	Naturally regen. cranberry (<i>Oxycoccus palustris</i>)	–707	5.4	0.08	Viru et al. 2020; Burdun et al. 2021	
Moore Rewetted former peat mining area					<i>Sphagnum cuspidatum</i> , <i>Eriophorum angustifolium</i> , <i>Molinia caerulea</i>	–949	233	–2.58	Beyer & Höper 2015	
Sphagnum site					<i>S. papillosum</i> , <i>S. cuspидatum</i> , <i>S. palustre</i> , <i>S. fallax</i> , <i>E. angustifolium</i> , <i>Erica tetralix</i> , <i>Juncus effusus</i> , <i>Betula pendula</i> , <i>Drosera</i> spp.	–987	24	0.55	Beyer & Höper 2015	
Hankhauser Moor, rewetted former bog grassland	53°16' N, 08°18' E	Germany	Sep. 2011–Aug. 2013	Poor	<i>Sphagnum palustre</i> cultivation	–5470 to –6290	10–14	0.01	Giinthner et al. 2017	

Table 1 (continued)

Type	Name	Coordinates	Country	Period of study	Nutrient status	Plant community/ treatment	NEE (CO ₂ -C)	CH ₄ -C	N ₂ O-N	Source
Ess-soo	57°54'51"N, 26°41'26"E	Estonia	(a)	Sep. 2011– Aug. 2013	Poor	<i>Sphagnum papillosum</i> cultivation Dredged hollow with naturally regenerated <i>Sphagnum</i>	-8750 to -8980	12–27	-1.0 to 1.0	Günther et al. 2017
Maima	58°35'54"N, 24°22'36"E	Estonia	(a)		Poor	Dredged hollow with naturally regenerated <i>Sphagnum</i>	-740	116	-0.11	Viru et al. 2020; Bur- dun et al. 2021
Provinzial-moor	52°40' N, 07°06' E	Germany		Mar. 2017– Mar. 2019	Rich	<i>S. papillosum</i> <i>S. palustre</i> ; ditch irrigation	-600 to 2200	3 to 53	0.5 to 1.3	Oestmann et al. 2022
Drents	52°41' N, 07°05' E	Germany			Rich	<i>S. papillosum</i> drip irrigation	700 to 900	1 to 4	1.9 to 12	Oestmann et al. 2022
Hankhauser Moor	53°16' N, 08°18' E	Germany		2015–2022	Rich	<i>Sphagnum spp.</i> , produc- tion field	-1,609±739	22.7±6.7	-0.47±1.57	Daun et al. 2023
Shallow lake	Giel'cikau Kasy	52°38'N, 25°21'E	Belarus	Aug. 2010– Aug. 2012	Rich	<i>Phragmites australis</i> , <i>Lemna spp</i>	-4,453 to -8,242	480 to 1470	0 to 0.005	Minke et al. 2016
Västkärr	59°06' N, 14°45' E	Sweden		2010–2014, snow-free periods (65% yr)	Rich	Graminoids, <i>Carex</i> spp., <i>Typha</i> spp		44	4.3	Jordan et al. 2020
Zilakalna, Tevgarsu	57°36' N, 25°10' E; 57°40' N, 24°57' E	Latvia		Dec. 2016– Dec. 2022	Poor	Permanently flooded peat extraction area; no vegetation	550±5	251±77	0.00±0.06	Bardule et al. 2023
Laiuse	58°47'17"N, 26°31'47"E	Estonia		Sep. 2019– June 2023	Poor	14 ha, open water, <i>P.</i> <i>australis</i> , <i>Typha</i> & sedges in littoral	1,079	292	0.17	Burdun et al. 2021
						Open bog/ fen				

Table 1 (continued)

Type	Name	Coordinates	Country	Period of study	Nutrient status	Plant community/ treatment	NEE (CO ₂ -C)	CH ₄ -C	N ₂ O-N	Source
Bois-des-Bel, restored bog	47°57'N, 69°26'W	Canada, Quebec	May–Oct. 2000, 2001, 2002	Poor	<i>Sphagnum spp.</i> , <i>Polytrichum spp.</i> , <i>Chamaedaphne calyculata</i> , <i>Vaccinium angustifolium</i> , <i>Ledum groenlandicum</i> , <i>E. vaginatum</i> , <i>T. latifolia</i>	–4,330 to –10,253				Waddington et al. 2010
Wandering River restored bog (former horticultural area)	55°11'N, 112°29'W	Canada, Alberta	July–Sep. 2011, May–July 2012	Poor	<i>Sphagnum spp.</i> , <i>E. vaginatum</i> , <i>Salix pedicellaris</i> , <i>Polytrichum strictum</i> , <i>Pohlia nutans</i>	–11,647 to 1,095	–3,23 to 540			Strack et al. 2014
Trebel River valley	54°06'N, 12°44'E	Germany	March 2011–March 2012	Rich	<i>Phragmites australis</i>	–830 to 320	110			Günther et al. 2015
Phragmites				Rich	<i>Carex spp.</i>	–430 to 710	105			Günther et al. 2015
Typha				Rich	<i>Typha spp.</i>	–30 to 290	590–630			Günther et al. 2015
Carex				Rich	<i>Erica tetralix</i> , <i>Calluna vulgaris</i> , <i>V. oxyococcus</i> , <i>Andromeda polifolia</i>	84	478	–0.16		Vanselow-Algan et al. 2015
Himmel-moor Rewetted in 1981 Heath site	53°44'20"N, 9°50'58"E	Germany	Aug. 2010–Jan. 2012	Poor	<i>Sphagnum spp.</i>	16	748	0.34		Vanselow-Algan et al. 2015
Sphagnum				Aug. 2010–Jan. 2012	Poor	<i>Molinia caerulea</i>	67	1.114	0.26	Vanselow-Algan et al. 2015
Purple moor grass				Aug. 2010–Jan. 2012	Poor					

Table 1 (continued)

Type	Name	Coordinates	Country	Period of study	Nutrient status	Plant community/ treatment	NEE (CO ₂ -C)	CH ₄ -C	N ₂ O-N	Source
Burns Bog Ecol. Cons. Area (BBECA)	49°06'37"N 123°00'03"W	Canada, British Columbia	June–Aug. 2014	Poor	<i>Ledum groenlandicum</i> , <i>Betula pendula</i> , <i>V. corymbosum</i> , <i>Sphagnum capillifolium</i> ,		240	0.0075		Christen et al. 2016
Restored peat mining area										
Rewetted Cleared										
Rewetted Sedge	49°07'09"N 123°00'01"W	Canada, British Columbia	June–Aug. 2014	Poor	<i>Rhynchospora alba</i> , <i>Dulichium arundinaceum</i> , <i>Sphagnum spp.</i> , <i>L. groenlandicum</i> , <i>V. uliginosum</i>		669	0.005		Christen et al. 2016
Bellacorik	54°07'30" N, 09°33'22" W	Ireland	Nov2008–Dec 2013	Poor	<i>Sphagnum spp.</i> , <i>Juncus effusus</i> , <i>Eriophorum spp.</i>	–1,040±800	90±20	0		Wilson et al. 2016b
Tässä	58°32'16" N, 25°51'43" E	Estonia	Mar. 2014–Mar. 2015	Poor	Bryophytes, Sphagnum; 50 high water level	2.0±0.9		–0.01±0.02	Järveoja et al. 2016b	
Seba Beach	53°27'17" N, 114°52'50" W	Canada, Alberta	May–Sep. 2015	Poor	Bryophytes, Sphagnum, herbs, shrubs; low water level	–247	0.9±0.5	0.2±0.1	Järveoja et al. 2016b	
Seba Beach	53°33'N, 114°44'W	Canada, Alberta	May–Aug. 2016, May–Aug. 2017	Poor	All plots (moss, bare, <i>Eriophorum</i>)			–0.135	Brummell et al. 2017	
					<i>Sphagnum spp.</i> , sedges (rest. 2009)	21			Bieniada & Strack 2021	
Bois-des-Bel	47°58'2" N 69°25'43" W	Canada, Quebec	Nov. 2013–Oct. 2014	Poor	Graminoids, <i>Polygonum</i> moss, <i>Sphagnum spp.</i> , sedges (rest. 2012)		703			Bieniada & Strack 2021
Uchter Moor		Germany	Jan–Dec. 2017	Rich	<i>Sphagnum spp.</i> , sedges	–900±180	44±2			
Rewetted since 1999 former peat mining area					<i>Eriophorum vaginatum</i> , <i>Molinia caerulea</i> E. <i>angustifolium</i>	262±33	48.8±9.8	0.57±0.21	Nugent et al. 2018	
									Schaller et al. 2022	

Table 1 (continued)

Negative values correspond to sequestration, positive values indicate emission to the atmosphere. Units: kg C ha ⁻¹ yr ⁻¹ ; kg N ha ⁻¹ yr ⁻¹	
Knox et al.: g CO ₂ –C m ⁻² yr ⁻¹ ; g CH ₄ –C m ⁻² yr ⁻¹ ; eddy covariance (FMA analyser).	
Günther et al. 2020: t CO ₂ ha ⁻¹ yr ⁻¹ ; kg CH ₄ –C ha ⁻¹ yr ⁻¹ ; kg N ₂ O–N ha ⁻¹ yr ⁻¹ .	
Bianchi et al.: t CO ₂ –eq ha ⁻¹ yr ⁻¹ , conversion factor of climate–carbon feedbacks: CH ₄ =34 and N ₂ O=298 (Myhre et al. 2013).	
Tiemeyer et al.: t CO ₂ –C ha ⁻¹ yr ⁻¹ ; kg CH ₄ –C ha ⁻¹ yr ⁻¹ ; kg N ₂ O–N ha ⁻¹ yr ⁻¹ , annual average.	
Wilson et al. 2016a: t CO ₂ –eq ha ⁻¹ yr ⁻¹ ; conversion factor of climate–carbon feedbacks: CH ₄ =34 and N ₂ O=298 (Myhre et al. 2013).	
Minkkinen et al.: g N ₂ O–N m ⁻² yr ⁻¹ ; chambers, GC; average annual values.	
Köhne et al.: kg CH ₄ ha ⁻¹ yr ⁻¹ ; chambers, GC, Los Gatos laser analyser.	
Mander et al.: mg CO ₂ –C m ⁻² h ⁻¹ ; µg CH ₄ –C m ⁻² h ⁻¹ ; µg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC, annual median values.	
Járveoja et al. 2016a: g CO ₂ –C m ⁻² h ⁻¹ ; g CH ₄ –C m ⁻² h ⁻¹ ; chambers, GC, average annual values.	
Maddison et al.: mg CO ₂ –C m ⁻² h ⁻¹ ; µg CH ₄ –C m ⁻² h ⁻¹ ; µg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC; average annual values.	
Vroom et al.: mg CH ₄ m ⁻² h ⁻¹ ; mg N ₂ O m ⁻² d ⁻¹ ; chambers on mesocosms in lab, Picarro G2308.	
Kandel et al.: Mg CO ₂ –C ha ⁻¹ yr ⁻¹ ; Mg CH ₄ –C ha ⁻¹ h ⁻¹ ; mg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC, range between average annual values.	
Lloyd et al.: µg CO ₂ –C m ⁻² h ⁻¹ ; µg CH ₄ –C m ⁻² h ⁻¹ ; µg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC, vegetation period average values.	
Burdun et al.: mg CO ₂ –C m ⁻² h ⁻¹ ; µg CH ₄ –C m ⁻² h ⁻¹ ; chambers, GC, average annual values.	
Viru et al.: mg CO ₂ –C m ⁻² h ⁻¹ ; µg CH ₄ –C m ⁻² h ⁻¹ ; µg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC, average monthly and annual values.	
Oestmann et al. 2022: t CO ₂ ha ⁻¹ yr ⁻¹ ; g CH ₄ –C m ⁻² yr ⁻¹ ; mg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC, range between average annual values of sub-sites.	
Beyer & Höper 2015: g CO ₂ C m ⁻² yr ⁻¹ ; g CH ₄ –C m ⁻² yr ⁻¹ ; mg N ₂ O–N m ⁻² yr ⁻¹ ; chambers, GC, average annual values.	
Günther et al. 2017: g CO ₂ m ⁻² yr ⁻¹ ; g CH ₄ –C m ⁻² yr ⁻¹ ; mg N ₂ O m ⁻² yr ⁻¹ ; chambers, GC, average annual.	
Daun et al.: g CO ₂ m ⁻² yr ⁻¹ ; g CH ₄ –C m ⁻² yr ⁻¹ ; g N ₂ O m ⁻² yr ⁻¹ ; chambers, GC, average annual ± sd.	
Minke et al.: mg CO ₂ –C m ⁻² h ⁻¹ ; µg CH ₄ –C m ⁻² h ⁻¹ ; µg N ₂ O–N m ⁻² h ⁻¹ ; chambers, LiC or (CO ₂); GC (CH ₄ , N ₂ O), average annual values.	
Jordan et al.: mmol CH ₄ m ⁻² h ⁻¹ ; µmol N ₂ O m ⁻² h ⁻¹ ; average annual values.	
Bardule et al.: mg CO ₂ –C m ⁻² h ⁻¹ ; mg CH ₄ –C m ⁻² h ⁻¹ ; µg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC, average annual ± se.	
Waddington et al.: g CO ₂ m ⁻² d ⁻¹ ; chambers, IRGA analyser, growing season average values.	
Strack et al.: g CO ₂ m ⁻² d ⁻¹ ; mg CH ₄ m ⁻² d ⁻¹ chambers, IRGA analyser, growing season average values.	
Günther et al.: g CO ₂ C m ⁻² yr ⁻¹ ; g CH ₄ –C m ⁻² yr ⁻¹ ; chambers, GC, average annual values.	
Vanselow-Algan et al.: g CO ₂ m ⁻² yr ⁻¹ ; µg CH ₄ –C m ⁻² s ⁻¹ ; µg N ₂ O–N m ⁻² s ⁻¹ , chambers, GC, average annual values.	
Christen et al.: g CO ₂ m ⁻² d ⁻¹ with EC; nmol CH ₄ m ⁻² s ⁻¹ and nmol N ₂ O–N m ⁻² s ⁻¹ with chambers & GC, June–August median values.	
Wilson et al. 2016b: mg CO ₂ –C m ⁻² h ⁻¹ ; mg CH ₄ –C m ⁻² h ⁻¹ ; µg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC, average annual ± sd.	
Járveoja et al. 2016b: mg CO ₂ –C m ⁻² h ⁻¹ ; µg CH ₄ –C m ⁻² h ⁻¹ ; µg N ₂ O–N m ⁻² h ⁻¹ ; chambers, GC, average annual values.	
Brummell et al.: mg N ₂ O–N m ⁻² d ⁻¹ ; chambers, GC, average annual values.	
Bieniada & Strack: µg CH ₄ –C m ⁻² h ⁻¹ ; Los Gatos; average annual values, chambers.	
Nugent et al.: g CO ₂ –C m ⁻² yr ⁻¹ ; eddy covariance; mean ± sd.	
Schaller et al.: EC LiCor g CO ₂ m ⁻² yr ⁻¹ ; EC LosGatos g CH ₄ m ⁻² yr ⁻¹ and mg N ₂ O–N m ⁻² yr ⁻¹ , average annual ± standard error values.	
^a NEE CO ₂ measurements from July 2021 until June 2023; CO ₂ Reco, CH ₄ and N ₂ O measurements from August 2017 until June 2023.	





Fig. 2 Average annual values of greenhouse gas fluxes in main groups of restored (rewetted) peatlands. **A–C** Annual average values of CO_2 -C (NEE), CH_4 -C and N_2O fluxes. **D**: greenhouse gas balance in CO_2 equivalent GWP₁₀₀ values (25

for CH_4 , 298 for N_2O ; Myhre et al. 2013). **E**: greenhouse gas balance in CO_2 equivalent GWP₅₀₀ values (7.6 for CH_4 , 153 for N_2O ; Forster et al. 2007)

(CO_2) sequestration rate ($> 1000 \text{ kg C ha}^{-1} \text{y}^{-1}$) followed by rewetted forests, open fens and bogs. However, some rewetted peatlands are potential source of both CO_2 and CH_4 , at least during the first 20 years after restoration (Vanselow-Algan et al. 2021).

According to our analysis, all restored peatlands were C sinks. However, there was significant variation of the data, mainly due to limited data availability and differences in the age since rewetting, as well as variation in plant cover development (Table 1). Average annual CO_2 -C stored in rewetted forests,

Fig. 3 Conceptual figure of C balance in hemiboreal ombrotrophic afforested peatland sites in a 300-year time-span. For drained and restored peatlands, time since drainage is shown. The tabs on the managed peatland forests graphs represent 100-year logging cycles. See Supplementary Table 1 for input parameters and modelling assumptions



open peatlands and paludicultural sites was -928 , -534 , and $-528 \text{ kg CO}_2\text{-C ha}^{-1} \text{ yr}^{-1}$, respectively (Table 1). These findings are consistent with the CO_2 flux values reported by Günther et al. (2020) for rewetted peatlands. Due to their diversity, paludicultural ecosystems showed the most significant variation in $\text{CO}_2\text{-C}$ fluxes ranging from high C capturing to moderate emissions, with the highest values observed in *Phalaris arundinacea* and *Poa* spp. plantation on rich fen peat (Kandel et al. 2020). Shallow lakes established on flooded peat extraction sites generally emitted low to moderate levels of CO_2 except extensively vegetated eutrophic sites (Minke et al. 2016).

In all rewetted peatlands, the average annual CH_4 fluxes were at moderate level, with the highest values observed in shallow lakes ($451 \text{ kg CH}_4\text{-C ha}^{-1} \text{ yr}^{-1}$) and followed by rewetted forests, open bogs/fens and paludiculture (218 , 163 , and $122 \text{ kg CH}_4\text{-C ha}^{-1} \text{ yr}^{-1}$, respectively; Table 1). Average annual N_2O fluxes ranged from low to moderate levels (0.01 to $4.45 \text{ kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$), except in *Phalaris* paludiculture plantations on poor and acid peat where liming and fertilization resulted in fluxes of up to $19 \text{ kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$ under high groundwater levels and $28.1 \text{ kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$ under low water levels (Maddison et al. 2016; Table 1).

In management of drained peatlands, global warming potential (GWP) of GHG-s should be considered to avoid making decisions based solely on

short-period benefits that may overlook the long-term climate cooling effect. Drained peatlands are known to be persistent CO_2 emitters over the long term, while rewetted peatlands as resilient re-established mire ecosystems effectively contribute to mitigating climate change, even considering radiative forcing of increased CH_4 emissions (Günther et al. 2020) and decreased N_2O emissions. Figure 2D and E demonstrate the climate effect of rewetted peatland ecosystems in GWP₁₀₀ and GWP₅₀₀ timeframes. Mires are the only terrestrial ecosystems capable to continuously sequester atmospheric carbon in the long term (Gorham et al. 2012; Cobb et al. 2020), i.e., the carbon sequestration itself is more important than its compound (methane vs carbon dioxide).

For restored (rewetted) peatlands see literature sources in Table 1.

For natural reference and drained peatlands the following literature sources served as the basis of this figure: Abdalla et al. 2016; Aitova et al. 2023; Bardule et al. 2023; Bianchi et al. 2021; Bieniada & Strack 2021; Brummell et al. 2017; Burdun et al. 2021; Busman et al. 2023; Clement et al. 2020; Couwenberg et al. 2010; Daun et al. 2023; Frolking et al. 2011; Griffis et al. 2020; Günther et al. 2020; Hergoualc'h and Verschot 2014; Hyvönen et al. 2009; Inubushi et al. 2003; Järveoja et al. 2016a,b; Jauhainen et al. 2008, 2012, 2019, 2023; Jordan et al. 2020; Kandel et al. 2020; Kull 2016; Maddison et al. 2016; Mander

et al. 2008, 2012, 2016; Melling et al. 2007; Nugent et al. 2018; Oestmann et al. 2022; Oktarita et al. 2017; Pärn et al. 2023; Petrescu et al. 2015; Rosset et al. 2022; Sakata et al. 2015; Salm et al. 2012; Sjögestren et al. 2011; Takakai et al. 2006; Tang et al. 2018; Toma et al. 2011; Truu et al. 2020; Turetsky et al. 2014; Veber et al. 2021, 202X; Wilson et al. 2016a,b; Wong et al. 2018.

Modelling approach

To estimate the carbon balance of afforested drained and restored rewetted peatlands we developed a conceptual model. Based on the availability of data from our research projects and similar studies in other countries we chose hemiboreal ombrotrophic peatlands as a modelling example. Figure 3 presents estimated long-term dynamics of C stock of afforested drained and restored rewetted hemiboreal ombrotrophic peatlands and their reference pristine bog ecosystem. In a 300-year perspective, considering stable climate conditions, drained forest peat (former bogs) will lose 1.8 and 1.0 t C ha⁻¹ yr⁻¹, respectively, due to peat mineralization. In afforested peatlands under continuous drainage, C losses from peat reach 0.9 t C ha⁻¹ yr⁻¹, whereas ecosystem C losses (the peat+vegetation budget and 100-year timber-harvesting regime) are 0.4–0.6 t C ha⁻¹ yr⁻¹. In comparison, fifty years after rewetting the naturally regenerated unmanaged peatland forest show decreased C loss, which is due to lower peat mineralization and C accumulation in biomass. In 160 years they achieve C dynamics similar to the pristine bog forest—with a moderate C sequestration rate of 0.1 t C ha⁻¹ yr⁻¹. Restoring a peat extraction site to bog ecosystem would become C neutral in nearly 20 years, and onwards continuous mean annual C sequestration of 0.22 t C ha⁻¹ yr⁻¹ is assumed (Supplementary Table 1).

Peatland forest drainage can give a significant increase in short-term C capture in biomass (Lohila et al. 2011), however, long-term dynamics in C stock and peat mineralisation remain largely unknown.

Restoration of crop plantations established on tropical peatlands and other dramatically altered peatlands differs from restoration of temperate and boreal bogs because the source community is predominantly swamp forest where peat-forming material is predominantly wood. Above ground and below ground litter

formation under the reforestation and management with moderate drainage can more easily compensate peat mineralization than in temperate and boreal areas (Couwenberg et al. 2010).

Conclusions

Understanding the dynamics of GHG fluxes caused by land-use change is essential for successful peatland restoration. Our analysis identified several contradictory research results and gaps in a deep understanding of these processes. Notably, there is a lack of GHG flux data for most of tropical drained and restored peatlands, with the exception of oil palm plantations. Another important issue is retention of C in restored peatlands, where we can differentiate between short-term C capture (GHG exchange between the peatland and atmosphere) and long-term C capture (accumulation in soil). Based on our analysis of literature sources and own research results from Estonia, rewetting of former peat extraction areas for further management or conservation is the only viable approach for long-term C sequestration. In contrast, afforestation combined with continuous peatland drainage may have short-term economic benefits but leads to C losses in the long term.

Uncertainties in long-term estimations of C storage and GHG flux dynamics are remarkably high and do not allow for exact predictions. However, even educated guesses can be valuable for decision making on further management of peatlands. To make accurate estimations, it is crucial to investigate the full combined impact of hydroclimate change, microbial processes, and vegetation on GHG emissions from restored peatlands.

Acknowledgements The concept for this paper was developed at the workshop titled "Peatlands for climate change mitigation in agriculture" that took place in Aarhus, Denmark, on 4–5 October 2022, and which was sponsored by the Organisation for Economic Co-operation and Development (OECD) Co-operative Research Programme: Sustainable Agricultural and Food Systems. This study was supported by the European Research Council (ERC) under grant agreement No 101096403 (MLTOM23415R), the Estonian Research Council (PRG352, and MOBERC20 and MOBERC44), the European Commission through the European Regional Development Fund (Center of Excellence EcolChange, TK-131), and the EU programmes: HORIZON-CSA project No 101079192 "Living Labs for Wetland Forest Research (LiWeFor)", LIFE21-IPC-EE-LIFE-SIP AdaptEst (MLTOM23090 (101069566)

"Implementation of national climate change adaptation activities in Estonia", and the Estonian State Forest Management Centre financed project LLTOM17250 "Water level restoration in cut-away peatlands: development of integrated monitoring methods and monitoring".

Author contributions UM: conceptualized the study, acquired funding, analyzed the data, and drafted the manuscript. ME: conceptualized the study, analyzed the data, generated visualizations, and contributed to reviewing the manuscript. LM: contributed to reviewing the manuscript. AK: conceptualized the study, acquired funding, collected original data, analyzed the data, and contributed to reviewing the manuscript.

Funding OECD Cooperative Research Programme, European Research Council, 101096403, Ülo Mander, HORIZON EUROPE Widening Participation and Strengthening the European Research Area, 101079192, Ülo Mander, Eesti Teadusagentuur, MOBERC20, Ülo Mander, MOBERC44, Ülo Mander, PRG352, Ülo Mander, European Regional Development Fund, EcolChange TK131, Ülo Mander, LIFE21-IPC-EE-LIFE-SIP, 101069566, Ain Kull, State Forest Management Centre, LLTOM17250, Ain Kull

Data availability Data are available within the article and its supplementary materials. For additional data or questions, please contact the authors.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Abdalla M, Hastings A, Truu J, Espenberg M, Mander Ü, Smith P (2016) Emissions of methane from northern peatlands: a review of management impacts and implications for future management options. *Ecol Evol* 6:7080–7102. <https://doi.org/10.1002/ece3.2469>
- Aitova E, Morley T, Wilson D, Renou-Wilson F (2023) A review of greenhouse gas emissions and reovals from Irish peatlands. *Mires Peat* 29(04):17. <https://doi.org/10.19189/MaP.2022.SNPG.StA.2414>
- Alm J, Schulman L, Walden J, Nykänen H, Martikainen P, Silvolu J (1999) Carbon balance of a boreal bog during year with an exceptionally dry summer. *Ecology* 80(1):161–174. <https://doi.org/10.2307/176987>
- Adanón-Rosell A, Scharnweber T, von Arx G, Peters RL, Smiljanić M, Weddell S, Wilmking M (2022) Growth and wood trait relationships of *Alnus glutinosa* in peatland forest stands with contrasting water regimes. *Front Plant Sci* 12:788106. <https://doi.org/10.3389/fpls.2021.788106>
- Bārdule A, Butlers A, Spalva G, Ivanovs J, Melniķis RN, Līcīte I, Andis Lazdiņš A (2023) The surface-to-atmosphere GHG fluxes in rewetted and permanently flooded former peat extraction areas compared to pristine peatland in hemiboreal Latvia. *Water* 15(10):1954. <https://doi.org/10.3390/w15101954>
- Beyer C, Höper H (2015) Greenhouse gas exchange of rewetted bog peat extraction sites and a *Sphagnum* cultivation site in Northwest Germany. *Biogeosciences* 12:2101–2117. <https://doi.org/10.5194/bg-12-2101-2015>
- Bianchi A, Larmola T, Kekkonen H, Saarnio S, Lång K (2021) Review of greenhouse gas emissions from rewetted agricultural soils. *Wetlands* 41:108. <https://doi.org/10.1007/s13157-021-01507-5>
- Bieniada A, Strack M (2021) Steady and ebullitive methane fluxes from active, restored and unrestored horticultural peatlands. *Ecol Eng* 169:106324. <https://doi.org/10.1016/j.ecoleng.2021.106324>
- Billet MF, Charman DJ, Clark JM, Evans CD, Evans MG, Ostle NJ, Worrall F, Burden A, Dinsmore KJ, Jones T, McNamara NP, Parry L, Rowson JG, Rose R (2010) Carbon balance of UK peatlands: current state of knowledge and future research challenges. *Clim Res* 45:13–29. <https://doi.org/10.3354/cr00903>
- Brummell ME, Lazcano C, Strack M (2017) The effects of *Eriophorum vaginatum* on N₂O fluxes at a restored, extracted peatland. *Ecol Eng* 106:287–295. <https://doi.org/10.1016/j.ecoleng.2017.06.006>
- Burdun I, Kull A, Maddison M, Veber G, Karasov O, Sagris V, Mander Ü (2021) Remotely sensed land surface temperature can be used to estimate ecosystem respiration in intact and disturbed northern peatlands. *J Geophys Res Biogeosciences*. <https://doi.org/10.1029/2021JG006411>
- Busman NA, Melling L, Gohe KJ, Imran Y, Sangok FE, Watanabe A (2023) Soil CO₂ and CH₄ fluxes from different forest types in tropical peat swamp forest. *Sci Total Environ* 858:159973. <https://doi.org/10.1016/j.scitotenv.2022.159973>
- Chiang F, Mazdiyasni O, AghaKouchak A (2021) Evidence of anthropogenic impacts on global drought frequency, duration, and intensity. *Nat Commun* 12:2754. <https://doi.org/10.1038/s41467-021-22314-w>
- Christen A, Jassal RS, Black TA, Grant NJ, Hawthorne I, Johnson MS, Lee SC, Merkens M (2016) Summertime greenhouse gas fluxes from an urban bog undergoing restoration through rewetting. *Mires Peat* 17(03):1–24
- Clément R, Pärn J, Maddison M, Henine H, Chaumont C, Tournebize J, Uri V, Espenberg M, Günther T, Mander Ü (2020) Frequency-domain electromagnetic induction for upscaling greenhouse gas fluxes in two hemiboreal drained peatland forests. *J Appl Geophys* 173:103944. <https://doi.org/10.1016/j.jappgeo.2020.103944>

- Cobb AR, Dommain R, Tan FY, Heng NHE, Harvey CF (2020) Carbon storage capacity of tropical peatlands in natural and artificial drainage networks. *Environ Res Lett* 15:114009. <https://doi.org/10.1088/1748-9326/aba867>
- Couwenberg J, Dommain R, Joosten H (2010) Greenhouse gas fluxes from tropical peatlands in South-East Asia. *Glob Change Biol* 16:1715–1732. <https://doi.org/10.1111/j.1365-2486.2009.02016.x>
- Daun C, Huth V, Gaudig G, Günther A, Krebs M, Jurasiczki G (2023) Full-cycle greenhouse gas balance of a Sphagnum paludiculture site on former bog grassland in Germany. *Sci Total Environ* 877:162943. <https://doi.org/10.1016/j.scitotenv.2023.162943>
- De Jong M, van Hal O, Pijlman J, van Eekeren N, Junginger M (2021) Paludiculture as paludifuture on Dutch peatlands: An environmental and economic analysis of Typha cultivation and insulation production. *Sci Total Environ* 792:148161. <https://doi.org/10.1016/j.scitotenv.2021.148161>
- Drollinger S, Maier A, Glatzel S (2019) Interannual and seasonal variability in carbon dioxide and methane fluxes of a pine peat bog in the Eastern Alps, Austria. *Agr Forest Meteorol* 275:69–78. <https://doi.org/10.1016/j.agrformet.2019.05.015>
- Evans CD, Peacock M, Baird AJ, Artz RRE, Burden A, Callaghan N, Chapman PJ, Cooper HM, Coyle M, Craig E, Cumming A, Dixon S, Gauci V, Grayson RP, Helfter C, Heppell CM, Holden J, Jones DL, Kaduk J, Levy P, Matthews R, McNamara NP, Misselbrook T, Oakley S, Page SE, Rayment M, Ridley LM, Stanley KM, Williamson JL, Worrall F, Morrison R (2021) Overriding water table control on managed peatland greenhouse gas emissions. *Nature* 593:548–552. <https://doi.org/10.1038/s41586-021-03523-1>
- Fluet-Chouinard E, Stocker BD, Zhang Z, Malhotra A, Melton JR, Poulter B, Kaplan JO, Goldewijk KK, Siebert S, Minayeva T, Hugelius G, Joosten H, Barthelmes A, Prigent C, Aires F, Hoyt AM, Davidson N, Finlayson CM, Lehner B, Jackson RB, McIntyre PB (2023) Extensive global wetland loss over the last three centuries. *Nature* 614:281–286. <https://doi.org/10.1038/s41586-022-05572-6>
- Forster P, Ramaswamy V, Artaxo P, Berntsen T, Betts TR, Fahey DW, Haywood J, Lean J, Lowe DC, Myhre G, Nganga J, Prinn R, Raga G, Schulz M, Van Dorland R (2007) Changes in Atmospheric Constituents and in Radiative Forcing. In: Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. In: Solomon S et al (eds) Cambridge University Press. United Kingdom and New York, Cambridge, p 234
- Freeman TR, Evans CD, Musarika S, Morrison R, Newman TR, Page SE, Wiggs GFS, Bell NGA, Styles D, Wen Y, Chadwick DR, Jones DL (2022) Responsible agriculture must adapt to the wetland character of mid-latitude peatlands. *Glob Change Biol* 28:3795–3811. <https://doi.org/10.1111/gcb.16152/>
- Frolking S, Talbot J, Jones MC, Treat CC, Kauffman JB, Tuittila ES, Roulet N (2011) Peatlands in the Earth's 21st century climate system. *Environ Rev* 19:371–396. [https://doi.org/10.1139/A11-014/](https://doi.org/10.1139/A11-014)
- Gallego-Sala AV, Charman DJ, Brewer S, Page SE, Prentice IC, Friedlingstein P, Moreton S, Amesbury MJ, Beilman DW, Björck S, Blyakharchuk T, Bochicchio C, Booth RK, Bunbury J, Camill P, Carless D, Chimner RA, Clifford M, Cressey C, Courtney-Mustaphi C, De Vleeschouwer F, de Jong R, Fialkiewicz-Koziel B, Finkelstein SA, Garneau M, Githumbi E, Hribjlan J, Holmquist J, Hughes PDM, Jones C, Jones MC, Karofeld E, Klein ES, Kokfelt U, Korhola A, Lacourse T, Le Roux G, Lamentowicz M, Large D, Lavoie M, Loisel J, Mackay H, MacDonald GM, Makila M, Magnan G, Marchant R, Marcisz K, Martínez Cortizas A, Massa C, Mathijssen P, Mauquoy D, Mighall T, Mitchell FJG, Moss P, Nichols J, Oksanen PO, Orme L, Packalen MS, Robinson S, Roland TP, Sanderson NK, Sannel ABK, Silva-Sánchez N, Steinberg N, Swindles GT, Turner TE, Uglow J, Välijanta M, van Bellen S, van der Linden M, van Geel B, Wang GP, Yu ZC, Zaragoza-Castells J, Zhao Y (2018) Latitudinal limits to the predicted increase of the peatland carbon sink with warming. *Nat Clim Change* 8:907–9813. <https://doi.org/10.1038/s41558-018-0271-1>
- Garcin Y, Schefuß E, Dargie GC, Lewis S et al (2022) Hydro-climatic vulnerability of peat carbon in the central Congo Basin. *Nature* 612:277–282. <https://doi.org/10.1038/s41586-022-05389-3>
- Gaudig G, Krebs M, Prager A, Wichmann S, Barney M, Caporn SJM, Emmel M, Fritz C, Grobe A, Gutierrez-Pacheco S, Hogue-Hugron S, Holzträger S, Irrgang S, Kämäriinen A, Karofeld E, Koch G, Koebbing JF, Kumari S, Matcutadze I, Oberpaur C, Oestmann J, Raabe P, Rammes D, Rochefort L, Schmilewski C, Sendzikaitė J, Smolders A, St-Hilaire B, van de Riet B, Wright B, Wright N, Zoch L, Joosten H (2017/18) Sphagnum farming from species selection to the production of growing media: a review. *Mires Peat* 20(13):1–30
- Gonzalez-Sargas E, Rochefort L (2019) Declaring success in Sphagnum peatland restoration: Identifying outcomes from readily measurable vegetation descriptors. *Mires Peat* 24:19. <https://doi.org/10.19189/MaP.2017.OMB.305>
- Gorham E (1991) Northern peatlands: Role in the carbon cycle and probable responses to climatic warming. *Ecol Appl* 1:182–195. <https://doi.org/10.2307/194181>
- Gorham E, Lehman C, Dyke A, Clymo D, Janssens J (2012) Long-term carbon sequestration in North American peatlands. *Quat Sci Rev* 58:77–82. <https://doi.org/10.1016/j.quascirev.2012.09.018>
- Grand-Clement E, Anderson K, Smith D, Angus M, Luscombe DJ, Gatis N, Bray LS, Brazier RE (2015) New approaches to the restoration of shallow marginal peatlands. *J Environ Manage* 161:417e430. <https://doi.org/10.1016/j.jenvman.2015.06.023>
- Griffis TJ, Roman DT, Wood JD, Deventer J, Fachin L, Rengifo J, Del Castillo D, Lilleskov E, Kolka R, Chimner RA, del Aguilá-Pasquel J, Wayson C, Hergoualc'h K, Baker JM, Cadillo-Quiroz H, Ricciuto DM (2020) Hydrometeorological sensitivities of net ecosystem carbon dioxide and methane exchange of an Amazonian palm swamp

- peatland. *Agr Forest Meteorol* 295:108167. <https://doi.org/10.1016/j.agrformet.2020.108167>
- Günther A, Jurasiński G, Albrecht K, Gaudig G, Krebs M, Glatzel S (2017) Greenhouse gas balance of an establishing Sphagnum culture on a former bog grassland in Germany. *Mires Peat* 20(02):1–16
- Günther A, Barthelmes A, Huth V, Joosten H, Jurasiński G, Koebisch F, Couwenberg J (2020) Prompt rewetting of drained peatlands reduces climate warming despite methane emissions. *Nat Commun* 11:1644. <https://doi.org/10.1038/s41467-020-15499-z>
- Haapalehto T, Juutinen R, Kareksela S, Kuitunen M, Tahvanainen T, Vuori H, Kotiaho JS (2017) Recovery of plant communities after ecological restoration of forestry-drained peatlands. *Ecol Evol* 7(19):7848–7858. <https://doi.org/10.1002/ece3.3243>
- Heger T, Jeschke JM, Febria C, Kollmann J, Murphy S, Rochefort L, Shackelford N, Temperton VM, Higgs E (2022) Mapping and assessing the knowledge base of ecological restoration. *Restor Ecol.* <https://doi.org/10.1111/rec.13676>
- Hergoualc'h K, Verschot LV, (2014) GHG emission factors for land use and land-use change in Southeast Asian peatlands. *Mitig Adapt Strat Gl* 19:789–807. <https://doi.org/10.1007/s11027-013-9511-x>
- Huang YY, Ciais P, Luo YQ, Zhu D, Wang YP, Qiu CJ, Goll DG, Guenet B, Makowski D, Graaf DE, Leifeld J, Kwon MJ, Hu J, Qu LY (2021) Tradeoff of CO₂ and CH₄ emissions from global peatlands under water-table drawdown. *Nat Clim Change* 11:618–622. <https://doi.org/10.1038/s41558-021-01059-w>
- Hyvönen NP, Huttunen JT, Shurpali NJ, Tavi NM, Repo ME, Martikainen PJ (2009) Fluxes of nitrous oxide and methane on an abandoned peat extraction site: effect of reed canary grass cultivation. *Biores Technol* 100(20):4723–4730. <https://doi.org/10.1016/j.biortech.2009.04.043>
- Inubushi K, Furukawa Y, Hadi A, Purnomo E, Tsuruta H (2003) Seasonal changes of CO₂, CH₄ and N₂O fluxes in relation to land-use change in tropical peatlands located in coastal area of South Kalimantan. *Chemosphere* 52:603–608. [https://doi.org/10.1016/S0045-6535\(03\)00242-X](https://doi.org/10.1016/S0045-6535(03)00242-X)
- Järveoja J, Peichl M, Maddison M, Teemusk A, Mander Ü (2016a) Full carbon and greenhouse gas balances of fertilized and non-fertilized reed canary grass cultivations on an abandoned peat extraction area in a dry year. *GCB Bioenergy* 8:952–968. <https://doi.org/10.1111/gcbb.12308>
- Järveoja J, Peichl M, Maddison M, Soosaar K, Vellak K, Karofeld E, Teemusk A, Mander Ü (2016b) Impact of water table level on annual carbon and greenhouse gas balances of a restored peat extraction area. *Biogeosciences* 13:2637–2651. <https://doi.org/10.5194/bg-13-2637-2016>
- Jauhainen J, Limin S, Silvennoinen H, Vasander H (2008) Carbon dioxide and methane fluxes in drained tropical peat before and after hydrological restoration. *Ecology* 89(12):3503–3514. <https://doi.org/10.1890/07-2038.1>
- Jauhainen J, Silvennoinen H, Hämäläinen R, Kusin K, Limin S, Raison RJ, Vasander H (2012) Nitrous oxide fluxes from tropical peat with different disturbance history and management. *Biogeosciences* 9:1337–1350. <https://doi.org/10.5194/bg-9-1337-2012>
- Jauhainen J, Alm J, Bjarnadottir B, Callesen I, Christiansen JR, Clarke N, Dalgaard L, He HX, Jordan S, Kazanaviciute V, Klemedtsson L, Laurén A, Lazdins A, Lehtonen A, Lohila A, Lupikis A, Mander Ü, Minkkinen K, Kasimir Å, Olsson M, Ojanen P, Óskarsson H, Sigurdsson BD, Søgaard G, Soosaar K, Vesterdal L, Laiho R (2019) Reviews and syntheses: Greenhouse gas exchange data from drained organic forest soils – a review of current approaches and recommendations for future research. *Biogeosciences* 16:4687–4703. <https://doi.org/10.5194/bg-16-4687-2019>
- Jauhainen J, Heikkilä J, Clarke N, He HX, Dalgaard L, Minkkinen K, Ojanen P, Vesterdal L, Alm J, Butlers A, Callesen I, Jordan S, Lohila A, Mander Ü, Óskarsson H, Sigurdsson BD, Søgaard G, Soosaar K, Kasimir Å, Bjarnadottir B, Lazdins A, Laiho R (2023) Greenhouse gas emissions from drained organic forest soils – synthesizing data for site-specific emission factors for boreal and cool temperate regions. *Biogeosciences Discuss.* <https://doi.org/10.5194/bg-2023-89>
- Jordan S, Strömgen M, Fiedler J, Lode E, Nilsson T, Lundin L (2020) Methane and nitrous oxide emission fluxes along water level gradients in littoral zones of constructed surface water bodies in a rewetted extracted peatland in Sweden. *Soil Syst* 4(1):17. <https://doi.org/10.3390/soilsystems4010017>
- Jurasiński G, Ahmad S, Anadon-Rosell A, Berendt J, Beyer F, Bill R, Blume-Werry G, Couwenberg J, Günther A, Joosten H, Koebisch F, Köhn D, Koldrack N, Kreyling J, Leinweber P, Lennartz B, Liu HJ, Michaelis D, Mrotzek A, Negassa W, Schenk S, Schmacka F, Schwieger S, Smiljanic M, Tanneberger F, Teuber L, Urich T, Wang HT, Weil M, Wilmking M, Zak D, Wrage-Mönnig N (2020) From understanding to Sustainable use of peatlands: the WETSCAPES approach. *Soil Syst* 4(1):14. <https://doi.org/10.3390/soilsystems4010014>
- Jurasiński G, Byrne K, Chojnicki BH, Christiansen JR, Huth V, Joosten H, Juszczak R, Juutinen S, Kasimir Å, Klemedtsson L, Kotowski W, Kull A, Lamantowicz M, Lindgren A, Linkovičienė R, Lohila A, Mander Ü, Manton M, Minkkinen K, Peters J, Renou-Wilson F, Sendžikaitė J, Šimanauskienė R, Tanneberger F, van Diggelen R, Vasander H, Wilson D, Zak DH, Couwenberg J (2023) Active afforestation of drained peatlands is not a viable option under the EU Nature Restoration Law. Zenodo. <https://doi.org/10.5281/zenodo.7831174>
- Kandel TP, Karki S, Elsgaard L, Labouriau R, Lærke PE (2020) Methane fluxes from a rewetted agricultural fen during two initial years of paludiculture. *Sci Total Environ* 713(15):136670. <https://doi.org/10.1016/j.scitotenv.2020.136670>
- Kask R, Pikk J (2009) Second thinning Scots pine wood properties in different forest site types in Estonia. *Balt for* 15(1):97–104
- Knox SH, Sturtevant C, Matthes JH, Koteen L, Verfaillie J, Baldocchi D (2015) Agricultural peatland restoration: effects of land-use change on greenhouse gas (CO₂ and CH₄) fluxes in the Sacramento-San Joaquin Delta. *Glob*

- Change Biol 21(2):750–765. <https://doi.org/10.1111/gcb.12745>
- Kohlenberg AJ, Turetsky MR, Thompson DK, Branfireun BA, Mitchell CPJ (2018) Controls on boreal peat combustion and resulting emissions of carbon and mercury. Environ Res Lett 13:035005. <https://doi.org/10.1088/1748-9326/aa9ea8>
- Köhönen D, Günther A, Schwabe I, Jurassiński G (2021) Short-lived peaks of stem methane emissions from mature black alder (*Alnus glutinosa* (L.) Gaertn.)—Irrelevant for ecosystem methane budgets? Plant Environ Interact 2:16–27. <https://doi.org/10.1002/pei3.10037>
- Kreyling J, Tanneberger F, Jansen F, van der Linden S, Aggenbach C, Blüml V, Couwenberg J, Emsens W-J, Joosten H, Klimkowska A, Kotowski W, Kozub L, Lennartz B, Liczner Y, Liu H, Michaelis D, Oehmke C, Parakennings K, Pleyl E, Poyda A, Raabe S, Röhl M, Rücker K, Schneider A, Schrautzer J, Schröder C, Schug F, Seeber E, Thiel F, Thiele S, Tiemeyer B, Timmermann T, Urich T, van Diggelen R, Vegelin K, Verbruggen E, Wilmking M, Wrage-Mönnig N, Wołejko L, Zak D, Jurassiński G (2021) Rewetting does not return drained fen peatlands to their old selves. Nat Commun 12:5693. <https://doi.org/10.1038/s41467-021-25619-y>
- Kull A (2016) Buffer zones to limit and mitigate harmful effects of long-term anthropogenic influence to maintain ecological functionality of bogs, stage II. Estonian Environmental Investment Centre (KIK), 183 p. (In Estonian). https://4ce0b57b-a630-4e1d-8a88-d1e1a7a51b96.filesusr.com/ugd/6b6658_446958f4118b44a2a68812820c3119b.pdf
- Külla T (1997) Below-ground and above-ground structure of a middle-aged Scots Pine stand and Norway Spruce stands. MSc Thesis, Estonian University of Life Sciences, Tartu, 100 p. (In Estonian).
- Leifeld J, Menichetti L (2017) The underappreciated potential of peatlands in global climate change mitigation strategies. Nat Commun 9:1071. <https://doi.org/10.1038/s41467-018-03406-6>
- Liu W, Fritz C, Weideveld STJ, Aben RCH, van den Berg M, Velthuis M (2022) Annual CO₂ budget estimation from chamber-based flux measurements on intensively drained peat meadows: Effect of gap-filling strategies. Front Environ Sci 10:803746. <https://doi.org/10.3389/fenvs.2022.803746>
- Liu HJ, Zak D, Zableckis N, Cossmer A, Langhammer N, Meermann B, Lennartz B (2023) Water pollution risks by smoldering fires in degraded peatlands. Sci Total Environ 871:161979. <https://doi.org/10.1016/j.scitotenv.2023.161979>
- Ljungqvist F, Krusic P, Sundqvist H, Zorita E, Brattström G, Frank D (2016) Northern Hemisphere hydroclimate variability over the past twelve centuries. Nature 532:94–98. <https://doi.org/10.1038/nature17418>
- Lloyd K, Madramootoo CA, Edwards KP, Grant A (2019) Greenhouse gas emissions from selected horticultural production systems in a cold temperate climate. Geoderma 349:45–55. <https://doi.org/10.1016/j.geoderma.2019.04.030>
- Lohila A, Minkkinen K, Aurela M, Tuovinen JP, Penttilä T, Ojanen P, Laurila T (2011) Greenhouse gas flux measurements in a forestry-drained peatland indicate a large carbon sink. Biogeosciences 8:3203–3218. <https://doi.org/10.5194/bg-8-3203-2011>
- Loisel J, Gallego-Sala A (2022) Ecological resilience of restored peatlands to climate change. Commun Earth Environ 3:208. <https://doi.org/10.1038/s43247-022-00547-x>
- Maddison M, Järveoja J, Kuller R, Muñoz M, Ostonen I, Teder H, Teemusk A, Torga R, Viru B, Mander Ü (2016) Reed canary grass (*Phalaris arundinacea*) cultivation as bioenergy crop on an abandoned peat extraction area with low soil pH. 20th EGU General Assembly, EGU2018, Proceedings from the conference held 4–13 April, 2018 in Vienna, Austria, p.6310. 2018EGUGA.20.6310M
- Mäkiranta P, Hytönen J, Aro L, Maljanen M, Pihlatie M, Potila H, Shurpali NJ, Laine J, Lohila A, Martikainen PJ, Minkkinen K (2007) Soil greenhouse gas emissions from afforested organic soil croplands and cutaway peatlands. Boreal Environ Res 12(2):159–175
- Maljanen M, Sigurdsson BD, Guðmundsson J, Óskarsson H, Huttunen JT, Martikainen PJ (2010) Greenhouse gas balances of managed peatlands in the Nordic countries – present knowledge and gaps. Biogeosciences 7:2711–2738. <https://doi.org/10.5194/bg-7-2711-2010>
- Mander Ü, Löhmus K, Teiter S, Uri V, Augustin J (2008) Gaseous nitrogen and carbon fluxes in riparian alder stands. Boreal Environ Res 13:231–241
- Mander Ü, Järveoja J, Maddison M, Soosaar K, Aavola R, Ostonen I, Salm J (2012) Reed canary grass cultivation mitigates greenhouse gas emissions from abandoned peat extraction areas. GCB Bioenergy 4:462–474. <https://doi.org/10.1111/j.1757-1707.2011.01138.x>
- Mander Ü, Krasnova A, Escuer-Gatius J, Espenberg M, Schindler T, Machacova K, Pärn J, Maddison M, Megonigal P, Pihlatie M, Kasak K, Niinemets Ü, Junninen H, Soosaar K (2021) Forest canopy mitigates soil N₂O emission during hot moments. Npj Clim Atmos Sci 4:39. <https://doi.org/10.1038/s41612-021-00194-7>
- Mander Ü (2016) Carbon and nitrogen cycle in drained peatland forests. Research project report. Appendix 1. Estonian State Forest Management Centre (RMK), 20 p. (In Estonian). https://media.rmk.ee/files/Rakendusuuringu%20lopparuanne_Kodusoometsad.pdf
- Martens M, Karlsson MPE, Ehde PM, Mattsson M, Weissenauer SEB (2021) The greenhouse gas emission effects of rewetting drained peatlands and growing wetland plants for biogas fuel production. J Environ Manage 277:111391. <https://doi.org/10.1016/j.jenvman.2020.111391>
- Melling L, Hatano R, Goh KJ (2007) Nitrous oxide emissions from three ecosystems in tropical peatland of Sarawak, Malaysia. Soil Sci Plant Nutr 53:792–805. <https://doi.org/10.1111/j.1747-0765.2007.00196.x>
- Minke M, Augustin J, Burlo A, Yarmashuk T, Chuvashova H, Thiele A, Freibauer A, Tikhonov V, Hoffmann M (2016) Water level, vegetation composition, and plant productivity explain greenhouse gas fluxes in temperate cutover fens after inundation. Biogeosciences 13:3945–3970. <https://doi.org/10.5194/bg-13-3945-2016>

- Minkkinen K, Laine J (1998) Long-term effect of forest drainage on the peatcarbon stores of pine mires in Finland. *Can J For Res* 28(9):1267–1275. <https://doi.org/10.1139/x98-104>
- Minkkinen K, Ojanen P, Koskinen M, Penttilä T (2020) Nitrous oxide emissions of undrained, forestry-drained, and rewetted boreal peatlands. *Forest Ecol Manag* 478:118494. <https://doi.org/10.1016/j.foreco.2020.118494>
- Myhre G, Shindell D, Bréon F-M, Collins W, Fuglestvedt J, Huang J, Koch D, Lamarque J-F, Lee L, Mendoza B, Nakajima T, Robock A, Stephens G, Takemura T, Zhang H (2013) Anthropogenic and natural radiative forcing. In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (eds) Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge United Kingdom and New York, pp 659–740
- Nugent KA, Strachan IB, Strack M, Roulet NT, Rochefort L (2018) Multi-year net ecosystem carbon balance of a restored peatland reveals a return to carbon sink. *Glob Chang Biol* 24(12):5751–5768. <https://doi.org/10.1111/gcb.14449>
- Oestmann J, Tiemeyer B, Düvel D, Grobe A, Dettmann U (2022) Greenhouse gas balance of Sphagnum farming on highly decomposed peat at former peat extraction sites. *Ecosystems* 25:350–371. <https://doi.org/10.1007/s10021-021-00659-z>
- Oktarita S, Hergoualc'h K, Anwar S, Verchot LV (2017) Substantial N₂O emissions from peat decomposition and N fertilization in an oil palm plantation exacerbated by hotspots. *Environ Res Lett* 12:104007. <https://doi.org/10.1088/1748-9326/aa80f1>
- Page S, Hosculo A, Langner A, Tansey K, Siegert F, Limin S, Rieley J (2009) Tropical peatland fires in Southeast Asia. In: Cochrane M (ed) Tropical Fire Ecology. Climate Change, Land Use and Ecosystem Dynamics. Praxis Publishing Ltd, Chichester, pp 263–287
- Pärn J, Verhoeven JTA, Butterbach-Bahl K, Dise NB, Ullah S, Aasa A, Egorov S, Espenberg M, Järveoja J, Jauhainen J, Kasak K, Klemmedsson L, Kull A, Laggoun-Défarge F, Lapshina ED, Lohila A, Löhman K, Maddison M, Mitsch WJ, Müller C, Niinemets Ü, Osborne B, Pae T, Salm JO, Sgouridis F, Sohar K, Soosaar K, Storey K, Teemusk A, Tenywa MM, Tournebize J, Truu J, Veber G, Villa JA, Zaw SS, Mander Ü (2018) Nitrogen-rich organic soils under warm well-drained conditions are global nitrous oxide emission hotspots. *Nat Commun* 9(9):1–8. <https://doi.org/10.1038/s41467-018-03540-1>
- Pärn J, Soosaar K, Schindler T, Machacova K, Alegria-Muñoz W, Fachin-Malaverri LM, Jibaja-Aspajo JL, Negron-Juarez R, Maddison M, Rengifo-Marin JE, Garay-Dinis DJ, Arista-Oversluijs AG, Avila-Fucos MC, Chávez-Vásquez R, Wampuch RH, Peas-García E, Sohar K, Cordova-Horna S, Pacheco-Gómez T, Urquiza-Muñoz JD, Tello-Espinoza R, Mander Ü (2023) Effects of water table fluctuation on greenhouse gas emissions from wetland soils in the Peruvian Amazon. In press, *Wetlands*
- Petrescu AMR, Lohila A, Tuovinen J-P, Baldocchi DD, Desai AR, Roulet NT, Vesala T, Dolman AJ, Oechel WC, Marcolla B, Friberg T, Rinne J, Matthes JH, Merbold L, Meijide A, Kiely G, Scottcornola ST, Zona D, Varlagin A, Lair DYF, Veenendaals E, Parmentier F-JW, Skiba U, Lund M, Hensen A, van Huissteden J, Flanagan LB, Shurpali NJ, Grünwald T, Humphreys ER, Jackowicz-Korczynski M, Aurela MA, Laurila T, Grüning C, Corradi CAR, Schrier-Uijls AP, Christensen TR, Tamstorf MP, Mastepanov M, Martikainen PJ, Verma SB, Bernhofer C, Cescatti A (2015) The uncertain climate footprint of wetlands under human pressure. *P Natl Acad Sci USA* 112(15):4594–4599. <https://doi.org/10.1073/pnas.1416267112>
- Rafat A, Rezanezhad F, Quinton WL, Humphreys ER, Webster K, Van Cappellen P (2021) Non-growing season carbon emissions in a northern peatland are projected to increase under global warming. *Commun Earth Environ* 2:111. <https://doi.org/10.1038/s43247-021-00184-w>
- Renou-Wilson F, Mueller C, Moser G, Wilson D (2016) To graze or not to graze? Four years greenhouse gas balances and vegetation composition from a drained and a rewetted organic soil under grassland. *Agric Ecosyst Environ* 222:156–170. <https://doi.org/10.1016/j.agee.2016.02.011>
- Rosset T, Binet S, Rigal F, Gandois L (2022) Peatland dissolved organic carbon export to surface waters: Global significance and effects of anthropogenic disturbance. *Geophys Res Lett*. <https://doi.org/10.1029/2021GL096616>
- Sakata R, Shimada S, Arai H, Yoshioka N, Yoshioka R, Aoki H, Kimoto N, Sakamoto A, Melling L, Inubushi K (2015) Effect of soil types and nitrogen fertilizer on nitrous oxide and carbon dioxide emissions in oil palm plantations. *Soil Scie Plant Nutr* 61:48–60. <https://doi.org/10.1080/00380768.2014.960355>
- Salm JO, Maddison M, Tammik S, Soosaar K, Truu J, Mander Ü (2012) Emissions of CO₂, CH₄ and N₂O from undisturbed, drained and mined peatlands in Estonia. *Hydrobiologia* 694(1):41–55. <https://doi.org/10.1007/s10750-011-0934-7>
- Samariks V, Lazdinš A, Bardule A, Kaleja S, Butlers A, Spalva G, Jansons A (2023) Impact of former peat extraction field afforestation on soil greenhouse gas emissions in hemiboreal region. *Forests* 14:184. <https://doi.org/10.3390/f142020184>
- Schaller C, Hofer B, Klemm O (2022) Greenhouse gas exchange of a NW German peatland, 18 years after rewetting. *J Geophys Res Biogeosciences*. <https://doi.org/10.1029/2020JG005960>
- Schindler T, Mander Ü, Machacova K, Espenberg M, Krasnov D, Escuer-Gatius J, Veber G, Pärn J, Soosaar K (2020) Short-term flooding increases CH₄ and N₂O emissions from trees in a riparian forest soil-stem continuum. *Sci Rep* 10(1):3204. <https://doi.org/10.1038/s41598-020-60058-7>
- Sjögersten S, Cheesman AW, Lopez O, Turner BL (2011) Biogeochemical processes along a nutrient gradient in a tropical ombrotrophic peatland. *Biogeochemistry* 104:147–163. <https://doi.org/10.1007/s10533-010-9493-7>
- Strack M, Keith AM, Xu B (2014) Growing season carbon dioxide and methane exchange at a restored peatland on

- the Western Boreal Plain. *Ecol Eng* 64:231–239. <https://doi.org/10.1016/j.ecoleng.2013.12.013>
- Swenson MM, Regan S, Bremmers DTH, Lawless J, Saunders M, Gill LW (2019) Carbon balance of a restored and cutover raised bog: implications for restoration and comparison to global trends. *Biogeosciences* 16:713–731. <https://doi.org/10.5194/bg-16-713-2019>
- Takakai F, Morishita T, Hashidoko Y, Darung U, Kuramochi K, Dohong S, Limin SH, Hatano R (2006) Effects of agricultural land-use change and forest fire on N₂O emission from tropical peatlands, Central Kalimantan, Indonesia. *Soil Sci Plant Nutr* 52:662–674. <https://doi.org/10.1111/j.1747-0765.2006.00084.x>
- Tang ACI, Stoy PC, Hirata R, Musin KK, Aeries EB, Wenceslaus J, Melling L (2018) Eddy covariance measurements of methane flux at a tropical peat forest in Sarawak, Malaysian Borneo. *Geophys Res Lett* 45:4390–4399. <https://doi.org/10.1029/2017GL076457>
- Tiemeyer B, Freibauer A, Borraz EA, Augustin J, Bechtold M, Beetz S, Beyer C, Ebliie M, Eickenscheidt T, Fiedler S, Förster C, Gensior A, Giebels M, Glatzel S, Heinichen J, Hoffmann M, Höper H, Jurasinski G, Lagner A, Leiber-Sauheitl K, Peichl-Brak M, Drösler M (2020) A new methodology for organic soils in national greenhouse gas inventories: Data synthesis, derivation and application. *Ecol Indic* 109:105838. <https://doi.org/10.1016/j.ecolind.2019.105838>
- Toma Y, Takakai F, Darung U, Kuramochi K, Limin SH, Dohong S, Hatano R (2011) Nitrous oxide emission derived from soil organic matter decomposition from tropical agricultural peat soil in central Kalimantan, Indonesia. *Soil Sci Plant Nutr* 57:436–451. <https://doi.org/10.1080/00380768.2011.587203>
- Truu M, Nölvak H, Ostonen M, Oopkaup K, Maddison M, Ligi T, Espenberg M, Uri V, Mander Ü, Truu J (2020) Soil bacterial and archaeal communities and their potential to perform N-cycling processes in soils of boreal forests growing on well-drained peat. *Front Microbiol* 11:591358. <https://doi.org/10.3389/fmicb.2020.591358>
- Turetsky MR, Kotowska A, Bubier J, Dise NB, Crill P, Hornbrook ERC, Minkkinen K, Moore TR, Myers-Smith IH, Nykänen H, Olefeldt D, Rinne J, Saarnio S, Shurpalov N, Tuittila E-S, Waddington JM, White JR, Wickland KP, Wilmking M (2014) A synthesis of methane emissions from 71 northern, temperate, and subtropical wetlands. *Glob Change Biol* 20:2183–2197. <https://doi.org/10.1111/gcb.12580>
- Vanselow-Algan M, Schmidt SR, Greven M, Fiencke C, Kutzbach L, Pfeiffer EM (2021) High methane emissions dominate annual greenhouse gas balances 30 years after bog rewetting. *Biogeosciences Discuss* 12:2809–2842. <https://doi.org/10.5194/bgd-12-2809-2015>
- Veber G, Kull A, Paal J (2021) Spatio-temporal variability of greenhouse gases along drainage gradient in 17 peatlands across Estonia. Proceedings of the 16th International Peatland Congress 2021, Tallinn, Estonia. Publcon PCO, pp. 120–126. ISBN 978-9916-4-0652-6
- Veber G, Paal J, Läänelaid A, Sohar K, Tampuu T, Salm JO, Maddison M, Teemusk A, Mander Ü, Kull A (202X) Simple vegetation and soil parameters determine greenhouse gas fluxes along drainage gradient in peatlands. Unpublished manuscript.
- Viru B, Veber G, Jaagus J, Kull A, Maddison M, Muuhel M, Espenberg M, Teemusk A, Mander Ü (2020) Wintertime greenhouse gas fluxes from hemiboreal drained peatlands. *Atmosphere* 11:731. <https://doi.org/10.3390/atmos11070731>
- Vroom RJE, Xi FJ, Geurts JJM, Chojnowska A, Smolders AJP, Lamers LPM, Fritz C (2018) Typha latifolia paludiculture effectively improves water quality and reduces greenhouse gas emissions in rewetted peatlands. *Ecol Eng* 124:88–98. <https://doi.org/10.1016/j.ecoleng.2018.09.008>
- Waddington JM, Strack M, Greenwood MJ (2010) Toward restoring the net carbon sink function of degraded peatlands: Short-term response in CO₂ exchange to ecosystem-scale restoration. *J Geophys Res Biogeosciences* 115(G1):G01008. <https://doi.org/10.1029/2009JG001090>
- Walker XJ, Baltzer JL, Cumming SG, Day NJ, Ebert C, Goetz S, Johnstone JF, Potter S, Rogers BM, Schuur EAG, Turetsky MR, Mack MC (2019) Increasing wild fires threaten historic carbon sink of boreal forest soils. *Nature* 572:520–523. <https://doi.org/10.1038/s41586-019-1474-y>
- Wichtmann W, Schröder C, Joosten H (2026) Paludiculture - Productive Use of Wet Peatlands: Climate Protection - Biodiversity - Regional Economic Benefits. Schweizerbart Science Publishers, Stuttgart, p 272
- Wilson D, Blain D, Couwenberg J, Evans CD, Murdiyarso D, Page SE, Renou-Wilson F, Rieley JO, Sirin A, Strack M, Tuittila ES (2016a) Greenhouse gas emission factors associated with rewetting of organic soils. *Mires Peat* 17:1–28. <https://doi.org/10.19189/MaP.2016.OMB.222>
- Wilson D, Farrell CA, Fallon D, Moser G, Müller C, Renou-Wilson F (2016b) Multiyear greenhouse gas balances at a rewetted temperate peatland. *Glob Change Biol* 22:4080–4095. <https://doi.org/10.1111/gcb.13325>
- Wilson D, Mackin F, Tuovinen JP, Moser G, Farrell C, Renou-Wilson F (2022) Carbon and climate implications of rewetting a raised bog in Ireland. *Glob Change Biol* 28(21):6349–6365. <https://doi.org/10.1111/gcb.16359>
- Wong GX, Hirata R, Hirano T, Kiewa F, Aeries EB, Musin KK, Waili JW, Lo KS, Melling L (2018) Micrometeorological measurement of methane flux above a tropical peat swamp forest. *Agr Forest Meteorol* 256–257:353–361. <https://doi.org/10.1016/j.agrformet.2018.03.025>
- Zhang W, Furtado K, Zhou TJ, Wu PL, Chen XL (2022) Constraining extreme precipitation projections using past precipitation variability. *Nat Commun* 13:6319. <https://doi.org/10.1038/s41467-022-34006-0>
- Zheng B, Ciais P, Chevallier P, Yang H, Canadell JG, Chen Y, van der Felde IR, Aben I, Chuvieco E, Davis SJ, Deeter M, Hong CP, Kong YW, Li HY, Li H, Lin X, He KB, Zhang QA (2023) Record-high CO₂ emissions from boreal fires in 2021. *Science* 379:912–917. <https://doi.org/10.1126/science.adc0805>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.