

Quality of life in noncompliant adults with phenylketonuria after resumption of the diet

**M. Bik-Multanowski · B. Didycz · R. Mozrzyimas ·
M. Nowacka · L. Kaluzny · W. Cichy · B. Schneiberg ·
J. Amilkiewicz · A. Bilar · M. Gizewska ·
A. Lange · E. Starostecka · A. Chrobot ·
B. I. Wojcicka-Bartłomiejczyk · A. Milanowski**

Published online: 12 January 2009
© SSIEM and Springer 2009

**Erratum to: J Inherit Metab Dis Short Report #133
(2008) Online
DOI 10.1007/s10545-008-0978-7**

The following Acknowledgement was omitted from the text:

The study was sponsored by the Nutricia Foundation Research Grant and by government grant for scientific research (NN402329233).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10545-008-0978-7>.

M. Bik-Multanowski (✉) · B. Didycz
Chair of Pediatrics, Jagiellonian University,
ul. Wielicka 265, 30-663 Krakow, Poland
e-mail: m.bik@wp.pl

R. Mozrzyimas
Voivodeship Hospital, Wrocław, Poland

M. Nowacka · A. Milanowski
Department of Pediatrics, Institute of Mother and Child,
Warsaw, Poland

L. Kaluzny · W. Cichy
Department of Pediatric Gastroenterology and Metabolism,
Medical University, Poznan, Poland

B. Schneiberg
Chair and Department of Pediatrics,
Children Nephrology and Endocrinology,
Silesian Medical Academy, Zabrze, Poland

J. Amilkiewicz
Children's Clinical Hospital,
Lublin, Poland

A. Bilar · M. Gizewska
II Department of Pediatrics,
Pomeranian Medical Academy,
Szczecin, Poland

A. Lange · E. Starostecka
Metabolic Department,
Polish Mother's Health Memorial Institute,
Łódź, Poland

A. Chrobot
Voivodeship Hospital, Bydgoszcz, Poland

B. I. Wojcicka-Bartłomiejczyk
Polish PKU Working Group,
Warsaw, Poland