

Evaluation of ecological and in vitro effects of boron on prostate cancer risk (United States)

**Wade T. Barranco · Paul F. Hudak ·
Curtis D. Eckhert**

Published online: 1 March 2007
© Springer Science+Business Media B.V. 2007

**Erratum to: Cancer Causes Control 18: 71–77
DOI 10.1007/s10552-006-0077-8**

Figure 1 (page 74) of this article was corrupted. The correct version and legend is given overleaf. The publishers apologise for this error.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10552-006-0077-8>

W. T. Barranco · C. D. Eckhert (✉)
Department of Environmental Health Sciences, University
of California, Los Angeles, CA 90095-1772, USA
e-mail: ceckhert@ucla.edu

P. F. Hudak
Department of Geography and Environmental Science
Program, University of North Texas, Post Office Box
305279, Denton, TX 76203-5279, USA
e-mail: hudak@unt.edu

Fig. 1 Scatter plots of mean groundwater B (**A, B**) and Se (**C, D**) concentrations in Texas state planning regions, as a function of prostate cancer incidence and mortality. y-axis represents the average annual age-adjusted (to the 2000 U.S. standard population) prostate cancer incidence and mortality rates per 100,000 people. Dotted line represents 95% confidence interval, while solid line represents the linear regression

