Correction to: Cannabis use is associated with patient and clinical factors in a population-based sample of colorectal cancer survivors

P. A. Newcomb^{1,2} · M. Ton^{1,2} · R. C. Malen¹ · J. L. Heffner¹ · J. Labadie¹ · A. I. Phipps^{1,2} · A. N. Burnett-Hartman^{1,3}

Published online: 19 October 2021
© Springer Nature Switzerland AG 2021

Correction to: Cancer Causes & Control
<https://doi.org/10.1007/s10552-021-01468-4>

Following publication of the original article, the author has identified an error in Table 1.

The correct version of Table 1 is given below.

The original article has been revised.

The original article can be found online at <https://doi.org/10.1007/s10552-021-01468-4>.

✉ P. A. Newcomb
newcomb@fredhutch.org

¹ Public Health Sciences Division, Fred Hutchinson Cancer Research Center, 1100 Fairview Avenue N, Seattle, WA 98109, USA

² Department of Epidemiology, University of Washington, Seattle, WA, USA

³ Institute for Health Research, Kaiser Permanente Colorado, Denver, CO, USA

Table 1 Odds ratios comparing current cannabis use among CRC patients in the Puget Sound region of Washington State according to participant characteristics, clinical factors, and quality of life (FACT-C) scores ($n=1,433$)

	Never/Former Cannabis Users $n=1,094$ n (%)	Current Cannabis Users $n=339$ n (%)	Unadjusted OR (95% CI)	Adjusted* OR (95% CI)
Age (years)				
65+	374 (34.2)	66 (19.5)	1.00 (ref)	1.00 (ref)
55–64	329 (30.1)	118 (34.8)	1.75 (1.26, 2.43)	1.85 (1.30, 2.62)
45–54	286 (26.1)	103 (30.4)	1.96 (1.40, 2.75)	2.20 (1.53, 3.17)
18–44	105 (9.6)	52 (15.3)	2.86 (1.87, 4.37)	3.30 (2.07, 5.26)
Sex				
Female	514 (47.0)	139 (41.0)	1.00 (ref)	1.00 (ref)
Male	580 (53.0)	200 (59.0)	1.28 (1.00, 1.63)	1.17 (0.89, 1.55)
Race				
Non-Hispanic White	854 (78.1)	275 (81.1)	1.00 (ref)	1.00 (ref)
Black	31 (2.8)	15 (4.4)	1.50 (0.80, 2.83)	1.53 (0.76, 3.07)
Hispanic	24 (2.2)	8 (2.4)	1.04 (0.46, 2.33)	0.77 (0.32, 1.86)
Asian & Pacific Islander	84 (7.7)	10 (3.0)	0.37 (0.19, 0.72)	0.29 (0.14, 0.58)
Other	58 (5.3)	18 (5.3)	0.96 (0.56, 1.66)	0.77 (0.43, 1.38)
Education				
College graduate or higher	505 (46.2)	138 (40.7)	1.00 (ref)	1.00 (ref)
Some College	374 (34.2)	114 (33.6)	1.11 (0.84, 1.47)	1.10 (0.81, 1.51)
HS graduate or less	205 (18.7)	84 (24.8)	1.49 (1.09, 2.05)	1.35 (0.95, 1.92)
BMI				
< 25.0	365 (33.4)	155 (45.7)	1.00 (ref)	1.00 (ref)
25.0–29.9	340 (31.1)	97 (28.6)	0.70 (0.52, 0.94)	0.65 (0.48, 0.90)
30.0+	359 (32.8)	81 (23.9)	0.55 (0.41, 0.75)	0.49 (0.35, 0.68)
Smoking				
Never	647 (59.1)	151 (44.5)	1.00 (ref)	1.00 (ref)
Former	359 (32.8)	145 (42.8)	1.72 (1.33, 2.24)	1.90 (1.42, 2.54)
Current	83 (7.6)	41 (12.1)	2.11 (1.39, 3.18)	1.82 (1.15, 2.88)
Alcohol				
Non-/Occasional Drinkers (< 1 g/day)	576 (52.7)	132 (38.9)	1.00 (ref)	1.00 (ref)
Light-to-Moderate Drinkers (1–28 g/day)	394 (36.0)	138 (40.7)	1.50 (1.15, 1.96)	1.36 (1.01, 1.83)
Heavy Drinkers (> 28 g/day)	109 (10.0)	63 (18.6)	2.48 (1.73, 3.55)	1.97 (1.32, 2.96)
FACT-C				
Lowest quartile	230 (21.0)	114 (33.6)	1.91 (1.46, 2.51)	1.51 (1.12, 2.02)
Stage				
Localized	447 (40.9)	75 (22.1)	1.00 (ref)	1.00 (ref)
Regional	455 (41.6)	162 (47.8)	2.08 (1.54, 2.79)	2.11 (1.54, 2.90)
Distant	167 (15.3)	96 (28.3)	3.35 (2.37, 4.73)	3.12 (2.15, 4.52)
Site				
Proximal	359 (32.8)	101 (29.8)	1.00 (ref)	1.00 (ref)
Distal	309 (28.2)	107 (31.6)	1.26 (0.92, 1.71)	1.11 (0.79, 1.56)
Rectal	408 (37.3)	128 (37.8)	1.14 (0.85, 1.53)	0.84 (0.60, 1.16)
Ostomy				
No Ostomy	887 (81.1)	252 (74.3)	1.00 (ref)	1.00 (ref)
Ostomy	203 (18.6)	86 (25.4)	1.49 (1.12, 1.99)	1.09 (0.80, 1.49)

*Adjusted for age, sex, race, BMI, smoking status, alcohol, education, and tumor stage

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.