

## **Exercise tissue Doppler echocardiography with strain rate imaging in healthy young individuals: feasibility, normal values and reproducibility**

**Björn Goebel · Raoul Arnold · Eric Koletzki ·  
Herbert E. Ulmer · Joachim Eichhorn ·  
Martin Borggreffe · Hans R. Figulla  
Tudor C. Poerner**

Published online: 27 September 2006  
© Springer Science+Business Media B.V. 2006

**Erratum to: Int J Cardiovasc Imaging  
DOI 10.1007/s10554-006-9130-7**

In the original publication of this article, the following information was missing:

The authors B. Goebel and R. Arnold contributed equally to this work.

---

The online version of the original article can be found at  
<http://www.dx.doi.org/10.1007/s10554-006-9130-7>

---

B. Goebel · H. R. Figulla · T. C. Poerner (✉)  
Klinik für Innere Medizin I, Universitätsklinikum  
Jena, Friedrich-Schiller-Universität Jena, Erlanger-  
Allee 101, D-07743 Jena, Germany  
e-mail: poerner@t-online.de

E. Koletzki · M. Borggreffe · B. Goebel ·  
T. C. Poerner  
I. Med. Klinik, Universitätsklinikum Mannheim,  
Theodor-Kutzer-Ufer 1-3, D-68135 Mannheim,  
Germany

R. Arnold · H. E. Ulmer · J. Eichhorn  
Department of Pediatric Cardiology,  
Universitätskinderklinik Heidelberg, Neuenheimer  
Feld 120, D-69120 Heidelberg, Germany

R. Arnold  
Department of Pediatric Cardiology,  
Universitätskinderklinik Freiburg, Mathildenstraße 1,  
D-79106 Freiburg, Germany