

Erratum to: Comparison of Clinical Efficacy and Safety of Clopidogrel Resinate With Clopidogrel Bisulfate in Patients Undergoing Percutaneous Coronary Intervention

Mahn-Won Park · Seung Hee Jeong · Sung Ho Her · Pum-Joon Kim · Jung Sun Cho · Chan Joon Kim · Wook-Sung Chung · Ki-Bae Seung · Hyeon-Woo Yim · Kiyuk Chang

Published online: 31 August 2013
© Springer Science+Business Media New York 2013

Erratum to: Cardiovasc Drugs Ther
DOI 10.1007/s10557-013-6466-9

The original version of this article unfortunately contained a mistake. P values for In-hospital death are missed in Table 1. The missing data are shown below:

Table 1 Baseline characteristics of the overall and propensity score-matched population according to clopidogrel type

Variables	Overall population			Propensity matched population		
	Clopidogrel bisulfate (<i>n</i> =9,127)	Clopidogrel resinate (<i>n</i> =1,360)	<i>p</i> value	Clopidogrel bisulfate (<i>n</i> =1,235)	Clopidogrel resinate (<i>n</i> =1,235)	<i>p</i> value
In-hospital death	136 (1.5)	30 (1.6)	0.24	10 (0.8)	15 (1.2)	0.12

The online version of the original article can be found at <http://dx.doi.org/10.1007/s10557-013-6466-9>.

M.-W. Park · S. H. Her · J. S. Cho · C. J. Kim
The Catholic University of Korea, Daejeon St. Mary's Hospital,
Daejeon, Republic of Korea

S. H. Jeong · H.-W. Yim
Department of Preventive Medicine, College of Medicine,
The Catholic University of Korea, Seoul, Republic of Korea

S. H. Jeong · H.-W. Yim
Clinical Research Coordinating Center of Catholic Medical Center,
Seoul, Republic of Korea

P.-J. Kim · W.-S. Chung · K.-B. Seung · K. Chang (✉)
The Catholic University of Korea, Seoul St. Mary's Hospital, 505
Banpo-Dong, Seocho-Gu, Seoul 137-040, Republic of Korea
e-mail: kiyuk@catholic.ac.kr