Cardiovascular Drugs and Therapy (2023) 37:613
https://doi.org/10.1007/s10557-022-07333-z

CORRECTION

Correction to: Effect of Insulin Resistance on Recurrence After Radiofrequency Catheter Ablation in Patients with Atrial Fibrillation

Zhe Wang[1,2] · Yi-Jia Wang[1] · Zhi-Yu Liu[3] · Qing Li[4] · Ya-Wei Kong[3] · Ying-Wei Chen[3] · Yi-Hong Sun[2] · Jian-Zeng Dong[3,5]

Published online: 21 March 2022

Correction to: Cardiovascular Drugs and Therapy
https://doi.org/10.1007/s10557-022-07317-z

The original article has been corrected. Certain texts under the section Risk Factors Associated with AF Recurrence have to be corrected. Please see below.

From: In the univariable analysis, patients with recurrence of AF after ablation had higher HOMAIR values; HR 1.264, 95% CI 1.096–1.457, P = 0.001] and LAD [(41.7 ± 6.6) mm vs. (39.1±6.4) mm; HR 1.046, 95% CI 1.009–1.084, P = 0.015].

To: In the univariable analysis, patients with recurrence of AF after ablation had higher HOMA-IR values (2.5 (IQR 1.9, 3.6) vs. 1.8 (IQR 1.4, 2.6); HR 1.264, 95% CI 1.096–1.457, $P = 0.001$) and LAD ((41.7 ± 6.6) mm vs. (39.1 ± 6.4) mm; HR 1.046, 95% CI 1.009–1.084, $P = 0.015$).

The original article can be found online at https://doi.org/10.1007/s10557-022-07317-z.

✉ Ying-Wei Chen
zzyingweichen@126.com

1 Graduate School of Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing 100730, China

2 Department of Cardiology, China-Japan Friendship Hospital, No.2 East Yinghua Road, Beijing 100029, Chaoyang District, China

3 Department of Cardiology, The First Affiliated Hospital of Zhengzhou University, No.1 Jianshe East Road, Erqi District, Zhengzhou 450052, Henan, China

4 Department of Cardiology, Peking University China-Japan Friendship School of Clinical Medicine, Beijing 100029, China

5 Department of Cardiology, Beijing Anzhen Hospital, Capital Medical University, Beijing 100029, China