## CORRECTION



## Correction to: Impact of storage time prior to cryopreservation on mechanical properties of aortic homografts

Ida Axelsson · Anna Gustafsson · Hanna Isaksson · Johan Nilsson · Torsten Malm

Published online: 17 April 2023 © Springer Nature B.V. 2023

Correction to: Cell Tissue Bank

https://doi.org/10.1007/s10561-023-10079-z

In the original publication of the article, the Fig. 5c was missed.

The complete Fig. 5 is given below. The original article has been corrected.

The original article can be found online at https://doi.org/10.1007/s10561-023-10079-z.

I. Axelsson (⊠) · T. Malm Tissue Bank Lund, Baravägen 37, 22242 Lund, Sweden e-mail: ida.axelsson@med.lu.se

T. Malm e-mail: torsten.malm@skane.se

I. Axelsson · J. Nilsson
Department of Cardiothoracic Surgery, Skane University
Hospital, Lund, Sweden
e-mail: johan.nilsson@med.lu.se

I. Axelsson · J. Nilsson · T. Malm
Department of Clinical Science, Cardiothoracic Surgery,
Lund University, Lund, Sweden
e-mail: johan.nilsson@med.lu.see-mail: torsten.malm@skane.se

A. Gustafsson · H. Isaksson Department of Biomedical Engineering, Lund University, Lund, Sweden e-mail: anna.gustafsson@bme.lth.se

H. Isaksson e-mail: hanna.isaksson@bme.lth.se


I Nilceon

Department of Translational Medicine, Artificial Intelligence and Bioinformatics in Cardiothoracic Sciences, Lund University, Lund, Sweden

T. Malm Pediatric Cardiac Surgery Unit, Children's Hospital, Skane University Hospital, Lund, Sweden



Fig. 5 a Elastic modulus at different time intervals of decontamination. The middle line is median, the lower and upper axis correspond to the first and third quartiles. The upper whisker extends from the axis to the largest value no further than 1.5 x interquartile range (IQR) from the axis, and the lower whisker extend to the smallest value no further than 1.5 x IQR from the axis. Data beyond whiskers are outlying points that are plotted individually. *p*-values represent results from Wilcoxon Signed Rank test, with individual comparisons between the group of 2-4 days and other groups, b Yield stress at different time points. See Fig. 5a for further description, c Energy at yield stress at different time points. See Fig. 5a for further description









Fig. 5 (continued)



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly

from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

