Correction to: Role of Delayed Neuroglial Activation in Impaired Cerebral Blood Flow Restoration Following Comorbid Injury

Zareen Amtul¹ · Jasmine Randhawa² · Abdullah N. Najdat² · David J. Hill^{3,4} · Edith J. Arany⁵

Published online: 10 October 2019

© Springer Science+Business Media, LLC, part of Springer Nature 2019

Correction to: Cellular and Molecular Neurobiology
<https://doi.org/10.1007/s10571-019-00735-y>

The original version of this article contained a random order of part labels for Fig. 4. The correct caption of Fig. 4 with correct order of part labels is given below.

The original article can be found online at <https://doi.org/10.1007/s10571-019-00735-y>.

✉ Zareen Amtul
zareen.amtul@gmail.com

¹ Department of Anatomy and Cell Biology, University of Western Ontario, London N6A 5C1, Canada

² Department of Biology, University of Western Ontario, London N6A 5B7, Canada

³ Departments of Medicine, Physiology, and Pharmacology, and Pediatrics, University of Western Ontario, London N6A 5C1, Canada

⁴ Lawson Health Research Institute, London, ON N6A 4V2, Canada

⁵ Department of Pathology and Laboratory Medicine, University of Western Ontario, London N6A 5C1, Canada

Fig. 4 Neuroglia bordering the lesion core: Coronal images demonstrate OX6 positive cells restricted to the lesion core (a). On d28, OX6 positive ramified microglial cells border the lesion core in ET1 and A β +ET1 rats (b). Plot shows a percentage of OX6 positive neuroglial cells bordering the lesion core in ipsilateral striatum of ET1 and A β +ET1 rats from d1 to d28 (c). Coronal images demonstrate the loss of GFAP immunoreactivity in the lesion core, which is filled with edematous fluid on d1 in A β +ET1 rats, d1 and d7 reprinted by permission from Springer Nature, Amtul et al. 2018C (d). On d28, GFAP positive active astrocytes border the lesion core in ET1 and A β +ET1 rats (e). High-resolution images are showing astrocytes surrounding the small pockets of lesion core (separated by black dotted lines) in ET1 and A β +ET1 rats on d28 (f). Plot shows a percentage of GFAP positive neuroglial cells bordering the lesion core in ipsilateral striatum of ET1 and A β +ET1 rats from d1 to d28 (g)