Correction to: Complex patterns of genetic and morphological differentiation in the Smallmouth Bass subspecies (*Micropterus dolomieu dolomieu* and *M. d. velox*) of the Central Interior Highlands

Joe C. Gunn¹ · Leah K. Berkman² · Jeff Koppelman² · Andrew T. Taylor³ · Shannon Brewer⁴ · James M. Long⁴ · Lori S. Eggert¹

Published online: 24 August 2020
© Springer Nature B.V. 2020

Correction to: Conservation Genetics

<https://doi.org/10.1007/s10592-020-01295-1>

In the original publication of the article, Table 2 was published incorrectly. The correct Table 2 is given in this Correction.

The original article can be found online at <https://doi.org/10.1007/s10592-020-01295-1>.

✉ Joe C. Gunn
jcg5g9@mail.missouri.edu

¹ Division of Biological Sciences, University of Missouri, 226 Tucker Hall, Columbia, MO 65211, USA

² Missouri Department of Conservation, 3500 E. Gans Rd., Columbia, MO 65201, USA

³ Department of Natural Resource Ecology and Management, Oklahoma State University, Stillwater, OK 74078, USA

⁴ U.S. Geological Survey, Oklahoma Cooperative Fish and Wildlife Research Unit, Department of Natural Resource Ecology and Management, Oklahoma State University, Stillwater, OK 74078, USA

Table 2 Raw and adjusted trait means of standard length (SL), orbital length (OL), head length (HL), and body depth (BD) for Neosho and Northern Smallmouth Bass samples assessed by (a) taxonomic range, and (b) genetic assignment

Trait	Raw mean (mm)		Adjusted mean (mm)		ANCOVA results			
	Neosho	Northern	Neosho	Northern	F_{int}	F_{cat}	p_{int}	p_{cat}
(a) Taxonomic range								
SL	234.864	246.726	232.274	238.232	1.200	2.100	0.271	0.166
OL	14.135	14.474	14.060	14.322	30.628	0.110	< 0.001	0.744
HL	79.966	76.463	79.616	73.282	0.220	14.070	0.636	0.001
BD	66.803	68.340	65.766	65.013	1.130	0.160	0.289	0.696
(b) Genetic assignment								
SL	206.588	244.809	232.809	234.963	1.800	0.300	0.181	0.620
OL	13.925	14.721	14.521	14.488	2.041	0.100	0.158	0.756
HL	71.921	78.259	79.433	74.645	0.120	3.140	0.728	0.097
BD	58.026	68.065	66.834	63.973	0.160	0.180	0.693	0.681

Trait means were back-transformed to provide values in millimeters. ANCOVA results are based on coefficients calculated from log-transformed trait values and include F -statistics and p -values for interactions (int) and categories (cat; taxonomic range or genetic assignment)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.