



Correction to: How do close others to those with anxiety feel about treatment? Development and validation of the Treatment Concerns Questionnaire–Close Others

Olivia A. Merritt¹ · Karen Rowa^{2,3} · Christine L. Purdon¹

Published online: 9 September 2022
© Springer Science+Business Media, LLC, part of Springer Nature 2022

Correction to: Cognitive Therapy and Research (2022)
<https://doi.org/10.1007/s10608-022-10318-9>

The original version of this article unfortunately contained a mistake. The corrected details are given below.

(a). The alignment of numbers in Table 2 was published incorrectly. The correct Table 2 is given below.

(b). The table notes for Table 4 were published in-text in error. The correct Table 4 is given below.

The original article has been updated.

The original article can be found online at <https://doi.org/10.1007/s10608-022-10318-9>.

✉ Olivia A. Merritt
oliviamerritt@outlook.com

¹ Psychology Department, University of Waterloo, 200 University Ave W, Waterloo, ON N2L 3G1, Canada

² Anxiety Treatment and Research Centre, St. Joseph's Healthcare, Hamilton, ON, Canada

³ Department of Psychiatry and Behavioural Neurosciences, McMaster University, Hamilton, ON, Canada

Table 2 Demographic information

	Participant (close other)	Person with anxiety
Age (years)		
Mean	35.8	36.9
Standard deviation	13.2	17.7
Gender, n (%)		
Male	94 (32.8%)	114 (39.7%)
Female	188 (65.5%)	164 (57.1%)
Gender non-binary, gender non-conforming	3 (1.0%)	7 (2.4%)
Declined to answer	1 (0.3%)	2 (0.7%)
Ethnicity, n (%)		
White	200 (69.7%)	195 (67.9%)
East Asian	21 (7.3%)	16 (5.6%)
South Asian	18 (6.3%)	20 (7.0%)
Black/African American	14 (4.9%)	15 (5.2%)
Other	34 (11.8%)	41 (14.3%)
Highest level of education, n (%)		
High school or less	203 (70.7%)	144 (50.2%)
Some or completed university/college degree	41 (14.3%)	45 (15.7%)
Some or completed graduate degree	4 (1.4%)	9 (3.1%)
Other or declined to answer		

Table 4 Factor statistics

Item #	Item	Factor Loadings				
		Communalities	F1	F2	F3	F4
3	My loved one will feel like a ‘failure’ if they do not ‘succeed’ in treatment	0.41	0.53	0.00	− 0.08	0.11
10	Treatment will be overwhelming for my loved one	0.60	0.75	− 0.05	− 0.14	− 0.07
12	My loved one will feel hopeless or depressed if treatment does not help	0.41	0.61	− 0.06	− 0.03	0.08
14	Treatment will create too much pressure for my loved one	0.75	0.81	0.00	− 0.02	0.08
15	Through treatment, my loved one may learn about symptoms or fears others have, and will take on these fears or symptoms as their own	0.45	0.55	0.21	− 0.01	− 0.02
17	Treatment will lead to worsened symptoms	0.61	0.62	0.21	0.03	0.10
5	Treatment will negatively change the way my loved one sees me	0.71	0.01	0.79	− 0.06	0.11
8	I will lose my relationship with my loved one because of this treatment	0.68	− 0.10	0.82	− 0.10	0.09
11	If my loved one’s anxiety improves, other issues in our relationship/family will become more prominent	0.34	0.09	0.56	0.00	− 0.14
13	Treatment will change my loved one in a negative way	0.57	0.32	0.51	0.16	0.18
7	My loved one will not use the coping tools that they learn in treatment	0.68	− 0.06	− 0.03	-0.73	0.23
16	My loved one will not be fully committed to improving	0.70	− 0.01	0.08	-0.83	− 0.04
18	My loved one will not complete all the required components of the treatment	0.67	0.04	0.03	-0.78	0.01
19	If treatment is difficult, my loved one will become discouraged and give up	0.74	0.30	− 0.03	− 0.70	− 0.04
2	Treatment will not work	0.71	0.19	− 0.11	0.01	0.76
4	Treatment will be a waste of time and/or money	0.54	− 0.03	0.20	− 0.09	0.64
6	Treatment will not be potent or comprehensive enough to help my loved one	0.60	0.10	− 0.02	− 0.14	0.64

Extraction communalities are reported. Factor loadings are represented by pattern coefficients. F1 = Adverse Reactions (AR); F2 = Personal/Family Consequences (PC); F3 = Lack of Commitment (LC); F4 = Ineffectiveness (IE). See Appendix for final measure.