

## Suppression of growth of highly-metastatic human breast cancer cells by norcantharidin and its mechanisms of action

Yan Huang · Qian Liu · Kun Liu ·  
Kazumi Yagasaki · Guoying Zhang

Published online: 2 August 2009  
© Springer Science+Business Media B.V. 2009

### Erratum to: Cytotechnology DOI 10.1007/s10616-009-9210-3

In the original version of this article, in the “Materials and methods” section, under the subsection “Animal experimentation and preparation of sera from NCTD-treated rabbits”, the third sentence currently reads:

‘NCTD was given via the intravenous injection to the rabbits once daily at a dose of  $10 \text{ mg mL}^{-1}$   $100 \text{ g}^{-1}$  body weight for 3 days’

Should read:

‘NCTD was given via the intravenous injection to the rabbits once daily at a dose of  $1 \text{ mg}$   $0.2 \text{ mL}^{-1} \text{ kg}^{-1}$  body weight for 3 days’.

---

The online version of the original article can be found under doi:[10.1007/s10616-009-9210-3](https://doi.org/10.1007/s10616-009-9210-3).

---

Y. Huang · Q. Liu · K. Liu · G. Zhang (✉)  
Laboratory of Molecular Pharmacology, School of  
Pharmacy, Yantai University, No. 30, Qing Quan Lu,  
Lai Shan Qu, 264005 Yantai, Shandong Province, China  
e-mail: zhang\_zhang6173@yahoo.com.cn

K. Yagasaki  
Department of Applied Biological Science, Tokyo Noko  
University, Saiwai-cho 3-5-8, Fuchu, Tokyo 183-8509,  
Japan