

Erratum to: Inhibitory effect of hot-water extract of quince (*Cydonia oblonga*) on immunoglobulin E-dependent late-phase immune reactions of mast cells

Takeshi Kawahara · Tatsuhiro Iizuka

Published online: 21 July 2011
© Springer Science+Business Media B.V. 2011

Erratum to: *Cytotechnology* (2011) 63:143–152 DOI 10.1007/s10616-010-9323-8

Under the heading “Preparation and cultivation of mouse bone marrow-derived mast cells (BMMCs)”, the sentence “The centrifuged cells were suspended in complete RPMI1640 medium supplemented with 10% FBS, 1 mM non-essential amino acids (Invitrogen), 5 ng/mL recombinant mouse IL-3 (Peprotech, Rocky Hill, NJ, USA), 50 mM 2-mercaptoethanol,

100 IU/mL penicillin, and 100 µg/mL streptomycin” should be replaced by “The centrifuged cells were suspended in complete RPMI1640 medium supplemented with 10% FBS, 0.1 mM non-essential amino acids (Invitrogen), 5 ng/mL recombinant mouse IL-3 (Peprotech, Rocky Hill, NJ, USA), 50 µM 2-mercaptoethanol, 100 IU/mL penicillin, and 100 µg/mL streptomycin”.

The corrected version of the Fig. 1 is given below.

The online version of the original article can be found under doi:[10.1007/s10616-010-9323-8](https://doi.org/10.1007/s10616-010-9323-8).

T. Kawahara (✉) · T. Iizuka
Laboratory of Food Bioscience, Faculty of Agriculture,
Shinshu University, 8304, Minamiminowa, Kamiina,
Nagano 399-4598, Japan
e-mail: tkawafb@shinshu-u.ac.jp

Fig. 1 Effects of quince HW on cytokine expression in RBL-2H3 cells. The expression of all cytokines was analyzed after 2 h of stimulation with IgE + Ag. Control means no stimulus (without IgE + Ag). Data are represented as the relative mean value \pm standard deviation (SD) against the unstimulated control ($n = 3$). * $p < 0.05$ and ** $p < 0.01$ vs. quince HW-untreated cells

