

Erratum to: Role of Nociceptors/Neuropeptides in the Pathogenesis of Visceral Hypersensitivity of Nonerosive Reflux Disease

Norimasa Yoshida · Masaaki Kuroda · Takahiro Suzuki ·
Kazuhiro Kamada · Kazuhiko Uchiyama · Osamu Handa ·
Tomohisa Takagi · Toshikazu Yoshikawa · Hirofumi Kuramoto

Published online: 9 December 2012
© Springer Science+Business Media New York 2012

Erratum to: Dig Dis Sci
DOI 10.1007/s10620-012-2337-7

Figure 1a of the article was published with graphic error of negative bar below zero in normal group. Corrected figure is given below.

The online version of the original article can be found under
doi:[10.1007/s10620-012-2337-7](https://doi.org/10.1007/s10620-012-2337-7).

N. Yoshida (✉) · T. Suzuki
Department of Gastroenterology, Japanese Red Cross Kyoto
Daiichi Hospital, 15-749 Honmachi Higashiyama-ku,
Kyoto 605-0981, Japan
e-mail: nyoshida@koto.kpu-m.ac.jp

N. Yoshida · M. Kuroda · K. Kamada · K. Uchiyama ·
O. Handa · T. Takagi · T. Yoshikawa
Molecular Gastroenterology and Hepatology, Kyoto Prefectural
University of Medicine, Kyoto, Japan

H. Kuramoto
Department of Applied Biology, Kyoto Institute of Technology,
Kyoto, Japan




Fig. 1 Expression of acid- and protease-sensitive nociceptors in the esophageal mucosa. Biopsy samples were taken from the esophageal mucosa at 5 mm from esophageal squamocolumnar junction of healthy control subjects ($n = 24$) and nonerosive reflux disease (NERD) subjects ($n = 24$). Expression of mRNA for transient receptor potential

vanilloid-1 (TRPV1, **a**), acid-sensing ion channel 3 (ASIC3, **b**), and protease-activated receptor 2 (PAR2, **c**) were quantified by real-time reverse transcription-polymerase chain reaction using biopsy mucosal materials and was corrected for that of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA. * $p < 0.05$ vs. normal